

Γ -*KU*-algebras

J. I. BAEK, SAMY M. MOSTAFA, FATEMA F. KAREEM, S. H. HAN, K. HUR

Received 9 October 2023; Revised 1 November 2023; Accepted 9 November 2023

ABSTRACT. In this paper, we introduce the concepts of positive implicative [resp. implicative and commutative] Γ -*KU*-algebras, and obtain their some properties (including characterizations) respectively and some relationships among them. Next, we propose the notions of positive implicative [resp. implicative and commutative] Γ -ideals of a Γ -*KU*-algebra, and deal with their some properties (including characterizations) respectively and some relationships among them. Finally, we define a topological Γ -*KU*-algebra and discuss its various topological structures.

2020 AMS Classification: 03G25, 06F35, 03E72, 08A35

Keywords: Γ -*KU*-algebra, Γ -*KU*-subalgebra, Γ -*KU*-ideal, Topological Γ -*KU*-algebra.

Corresponding Author: Samy M. Mostafa (samymostafa@yahoo.com),
S. H. Han (shhan235@wku.ac.kr)

1. INTRODUCTION

In 1978, Iséki and Tanaka [1] introduced the notion of *BCK*-algebras as a generalization of *I*-algebras proposed by Imai and Iséki [2] in 1966. Iséki [3] defined *BCI*-algebras which is a proper subclass of *BCK*-algebras. Some researchers [4, 5, 6, 7, 8] studied properties of ideals which important role in *BCK*-algebras and *BCI*-algebras respectively. Furthermore, Dudek and Zhang [9] introduced a concept of *BCC*-algebras and discussed relationships between ideals and congruences in *BCC*-algebras. Also, some researchers [10, 11, 12, 13] investigated topological structures on *BCK*-algebras and *BCI*-algebras respectively.

In 2009, Prabpayak and Leerawat [14] defined a *KU*-algebra as new logical algebras and studied properties ideals and congruences in *KU*-algebras. Also, They [15] dealt with isomorphisms in *KU*-algebras. After then, many researchers [16, 17, 18, 19, 20, 21, 22] investigated various properties in *KU*-algebras. Recently, Hur et al. [23] introduced the notion of square root fuzzy sets and obtained some properties of square root fuzzy ideals of a *KU*-algebra. *KU*-algebras were studied by many

mathematicians and applied to group theory, functional analysis, probability theory, topology, graph theory and computer science etc.

In 1981, Sen [24] proposed the notion of Γ -semigroups as a generalization of semigroups. Rao [25] introduced the concept of Γ -groups as generalization of groups and studied its various properties. Also, Rao [26] proposed the notion of Γ -semirings as a generalization of semirings. After then, Kaushik and Moin [27] investigated bi- Γ -ideals in a Γ -semiring, Rao and Venkateswarlu [28] studied some properties related to regular Γ -incline and field Γ -semiring.

In 2022, Saeid et al. [29] introduced the concept of Γ -*BCK*-algebras as a generalization of *BCK*-algebras and dealt with some properties of subalgebras, ideals, closed ideals, normal subalgebras and normal ideals in Γ -*BCK*-algebras and quotient Γ -*BCK*-algebras. After that time, Shi et al. [30] defined positive implicative [resp. implicative and commutative] Γ -*BCK*-algebras and positive implicative [resp. implicative and commutative] Γ -ideals of a Γ -*BCK*-algebra, and studied their some properties respectively and some relationships among them. Also, Shi et al. [31] defined a topological Γ -*BCK*-algebra and studied some of its topological structures.

It is the aim of our study to introduce the notion of Γ -*KU*-algebras as a generalization of *KU*-algebras, and define positive implicative [resp. implicative and commutative] Γ -*KU*-algebras and discuss their some properties (including characterizations) respectively and some relationships among them. Also, we define positive implicative [resp. implicative and commutative] Γ -ideals of a Γ -*KU*-algebra, and obtain their some properties (including characterizations) respectively and some relationships among them. Furthermore, we introduce the concept of topological Γ -*KU*-algebras and study its various topological structures.

2. PRELIMINARIES

We recall some definitions needed in next sections. An algebra $X = (X, *, 0)$ means a nonempty set X together with a binary operation $*$ and a special element 0 .

Definition 2.1 ([14]). An algebra X is called a *KU*-algebra, if it satisfies the following axioms: for any $x, y, z \in X$,

- (KU₁) $(x * y) * [(y * z) * (x * z)] = 0$,
- (KU₂) $x * 0 = 0$,
- (KU₃) $0 * x = x$,
- (KU₄) $x * y = 0$ and $y * x = 0$ imply $x = y$.

In *KU*-algebra X , we define a binary operation \leq on X as follows: for any $x, y \in X$,

$$x \leq y \text{ if and only if } y * x = 0.$$

Definition 2.2. Let X be a *KU*-algebra. Then X is said to be:

- (i) *KU*-positive implicative [19], if $(z * x) * (z * y) = z * (x * y)$ for any $x, y, z \in X$,
- (ii) *KU*-commutative [20], if $(y * x) * x = (x * y) * x$ for any $x, y \in X$,
- (ii) *KU*-implicative [19], if $x = (x * y) * x$ for any $x, y \in X$.

Definition 2.3 (See [14]). Let A be a nonempty set of a *KU*-algebra X . Then A is called a *subalgebra* of X , if $x * y \in A$ for any $x, y \in A$.

Definition 2.4. Let I be a nonempty set of a KU -algebra X . Then

(a) I is called an *ideal* of X [19], if it satisfies the following conditions: for any $x, y \in X$,

$$(I_1) 0 \in I,$$

$$(I_2) x * y \in I \text{ and } x \in I \text{ imply } y \in I.$$

(a) I is called an *ideal* (briefly, KUI) of X [14], if it satisfies the following conditions: for any $x, y, z \in X$,

$$(KUI_1) 0 \in I,$$

$$(KUI_2) x * (y * z) \in I \text{ and } y \in I \text{ imply } x * z \in I.$$

Definition 2.5 ([18]). Let I be a nonempty set of a KU -algebra X . Then I is called a *KU -positive implicative ideal* (briefly, KUPII) of X , if it satisfies the following conditions: for any $x, y, z \in X$,

$$(KUPI_1) 0 \in I,$$

$$(KUPII_2) z * (x * y) \in I \text{ and } z * x \in I \text{ imply } z * y \in I.$$

Definition 2.6 ([18]). Let I be a nonempty set of a KU -algebra X . Then I is called an *KU -implicative ideal* (briefly, KUII) of X , if it satisfies the following conditions: for any $x, y, z \in X$,

$$(KUII_1) 0 \in I,$$

$$(KUII_2) (x * y) * (z * x) \in I \text{ and } z \in I \text{ imply } x \in I.$$

Definition 2.7 ([18]). Let I be a nonempty set of a KU -algebra X . Then I is called a *KU -commutative ideal* (briefly, KUCI) of X , if it satisfies the following conditions: for any $x, y, z \in X$,

$$(KUCI_1) 0 \in I,$$

$$(KUCI_2) y * (z * x) \in I \text{ and } z \in I \text{ imply } [(x * y) * y] * x \in I.$$

Definition 2.8 ([26]). Let X and Γ be two nonempty sets. Then X is called a Γ -*semigroup*, if there is a mapping $f : X \times \Gamma \times X \rightarrow X$, denoted by $f(x, \alpha, y) = x\alpha y$ for each $(x, \alpha, y) \in X \times \Gamma \times X$, such that it satisfies the following condition: for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$,

$$(2.1) \quad x\alpha(y\beta z) = (x\alpha y)\beta z.$$

3. SOME PROPERTIES OF Γ - KU -ALGEBRAS

In this section, we introduce the concept of Γ - KU -algebras and study some of its properties.

Definition 3.1. Let X be a set with a constant 0 and let Γ be a nonempty set. Then X is called a Γ - *KU -algebra*, if there is a mapping $f : X \times \Gamma \times X \rightarrow X$, denoted by $f(x, \alpha, y) = x\alpha y$ for each $(x, \alpha, y) \in X \times \Gamma \times X$, satisfying the following axioms: for any $x, y, z \in X$ and $\alpha, \beta \in \Gamma$,

$$(\Gamma KU_1) (x\alpha y)\beta[(y\alpha z)\beta(x\alpha z)] = 0,$$

$$(\Gamma KU_2) 0\alpha x = x,$$

$$(\Gamma KU_3) x\alpha 0 = 0,$$

$$(\Gamma KU_4) x\alpha y = 0 = y\alpha x \text{ imply } x = y.$$

Remark 3.2. From (ΓKU_1) , (ΓKU_3) and (ΓKU_1) , (ΓKU_2) , we have

$$(3.1) \quad x\alpha x = 0, z\beta(x\alpha z) = 0 \text{ for any } x, z \in X \text{ and any } \alpha, \beta \in \Gamma.$$

We define a binary relation \leq on Γ - KU -algebra X as follows: for any $x, y \in X$ and each $\alpha \in \Gamma$,

$$(3.2) \quad x \leq y \Leftrightarrow y\alpha x = 0.$$

Then we obtain the following properties.

Proposition 3.3. *Let X be a Γ - KU -algebra. Then the following inequalities hold: for any $x, y, z \in X$ and each $\alpha, \beta \in \Gamma$,*

- (1) $(y\alpha z)\beta(x\alpha z) \leq x\alpha y$,
- (2) $0 \leq x$,
- (3) $x \leq y$ and $y \leq x$ imply $x = y$,
- (4) $x \leq x$,
- (5) $x\alpha y \leq y$.

It is clear that for a Γ - KU -algebra X and a fixed $\alpha \in \Gamma$, if we define the operation $*$: $X \times X \rightarrow X$ as follows: for any $x, y \in X$,

$$x * y = x\alpha y,$$

then $(X, *, 0)$ is a KU -algebra and is denoted by X_α .

Example 3.4. (1) Let $X = \{0, 1, 2, 3\}$, let $\Gamma = \{\alpha, \beta, \gamma\}$ and let the ternary operation be defined by the table:

α	0	1	2	3	β	0	1	2	3	γ	0	1	2	3
0	0	1	2	3	0	0	1	2	3	0	0	1	2	3
1	0	0	2	2	1	0	0	2	3	1	0	0	3	3
2	0	0	0	0	2	0	0	0	0	2	0	0	0	0
3	0	0	0	0	3	0	0	0	0	3	0	1	2	0

Table 3.1

Then we can easily check that X is Γ - KU -algebra. Also, X_α, X_β and X_γ can confirm KU -algebras.

(2) Let $X = \{0, 1, 2, 3, 4, 5\}$, let $\Gamma = \{\alpha, \beta, \gamma, \delta, \psi\}$ and let the ternary operation be defined by the table:

Then we can easily check that X is Γ - KU -algebra.

Proposition 3.5. *Let X be a Γ - KI -algebra. Then the followings hold: for any $x, y, z \in X$ and each $\alpha \in \Gamma$,*

- (1) $x \leq y$ implies $y\alpha z \leq x\alpha z$,
- (2) $x \leq y$ and $y \leq z$, then $x \leq z$.

Proof. (1) Suppose $x \leq y$. Then clearly, $y\alpha x = 0$ for each $\alpha \in \Gamma$. Thus by the axiom (ΓKU_1) , we have

$$(y\alpha x)\beta[(x\alpha z)\beta(y\alpha z)] = 0, \text{ i.e., } 0\beta[(x\alpha z)\beta(y\alpha z)] = 0 \text{ for any } \alpha, \beta \in \Gamma.$$

So by the axiom (ΓKU_2) , $(x\alpha z)\beta(y\alpha z) = 0$. Hence by (3.2), $y\alpha z \leq x\alpha z$.

(2) Suppose $x \leq y$ and $y \leq z$. Then by (1), $z\alpha x \leq y\alpha x$. Since $x \leq y$, $y\alpha x = 0$. Thus $z\alpha x \leq 0$. By Proposition 3.3 (2), $0 \leq z\alpha x$. So by Proposition 3.3 (3), $z\alpha x = 0$. Hence $x \leq z$. □

α	0	1	2	3	4	5	α	0	1	2	3	4	5	α	0	1	2	3	4	5
0	0	1	2	3	4	5	0	0	1	2	3	4	5	0	0	1	2	3	4	5
1	0	0	2	3	4	5	1	0	0	3	4	5	1	1	0	0	4	5	1	2
2	0	1	0	3	4	5	2	0	2	0	4	5	1	2	0	3	0	5	1	2
3	0	1	2	0	4	5	3	0	2	3	0	4	1	3	0	3	4	0	1	2
4	0	1	2	3	0	5	4	0	2	3	4	0	1	4	0	3	4	5	0	2
5	0	1	2	3	4	0	5	0	2	3	4	5	0	5	0	3	4	5	1	0

δ	0	1	2	3	4	5	ψ	0	1	2	3	4	5
0	0	1	2	3	4	5	0	0	1	2	3	4	5
1	0	0	5	1	2	3	1	0	0	1	2	3	4
2	0	4	0	1	2	3	2	0	5	0	2	3	4
3	0	4	5	0	2	3	3	0	5	1	0	3	4
4	0	4	5	1	0	3	4	0	5	1	2	0	4
5	0	4	5	1	2	0	5	0	5	1	2	3	0

Table 3.2

From Proposition 3.3 (3), (3.1) and Proposition 3.5 (2), it is obvious that (X, \leq) is a poset with the least element 0.

Proposition 3.6. *Let X be a Γ - KU -algebra. Then the followings hold: for any $x, y, z \in X$ and each $\alpha, \beta \in \Gamma$,*

$$(3.3) \quad z\alpha(y\beta x) = y\alpha(z\beta x).$$

Proof. From the axiom (ΓKU_1) , $(0\alpha z)\beta[(z\alpha x)\beta(0\alpha x)] = 0$. Then by the axioms (ΓKU_2) , $z\beta[(z\alpha x)\beta x] = 0$, i.e.,

$$(3.4) \quad (z\alpha x)\beta x \leq z.$$

Thus by (3.4), Proposition 3.5 (1) and Proposition 3.3 (1), we have

$$(3.5) \quad z\alpha(y\beta x) \leq [(z\alpha x)\beta x]\alpha(y\beta x) \leq y\alpha(z\beta x).$$

Since x, y, z are arbitrary, by interchanging y and z in the equality (3.5), we get

$$(3.6) \quad y\alpha(z\beta x) \leq z\alpha(y\beta x).$$

So the axiom (ΓKU_4) , the identity (3.3) holds. □

The followings are immediate consequences of Proposition 3.6.

Corollary 3.7. *Let X be a Γ - KU -algebra. Then the followings hold: for any $x, y, z \in X$ and each $\alpha, \beta \in \Gamma$,*

- (1) $x\alpha y \leq z$ if and only if $x\alpha z \leq y$,
- (2) $(y\alpha x)\beta x \leq y$.

The following is an immediate consequence of Theorem 3.3 (1) and Corollary 3.7 (1).

Corollary 3.8. *In a Γ - KU -algebra X , the followings hold: for any $x, y, z \in X$ and each $\alpha \in \Gamma$,*

$$(x\alpha y)\beta(x\alpha z) \leq y\alpha z, \text{ i.e., } (y\alpha z)\beta[(x\alpha y)\beta(x\alpha z)] = 0.$$

The following is an immediate consequence of Corollary 3.8.

Corollary 3.9. *In a Γ - KU -algebra X , the followings hold: for any $x, y, z \in X$ and each $\alpha \in \Gamma$,*

$$x \leq y \text{ implies } z\alpha x \leq z\alpha y.$$

We define a binary operation \wedge on a Γ - KU -algebra X as follows: for any $x, y \in X$ and any $\alpha, \beta \in \Gamma$,

$$x \wedge y = (y\alpha x)\beta x.$$

Then it is obvious that $x \wedge y$ is a lower bound of $\{x, y\}$ and $x \wedge x = 0, x \wedge 0 = 0 = 0\alpha x$. However, $x \wedge y \neq y \wedge x$ in general.

Proposition 3.10. *In a Γ - KU -algebra X , the followings hold: for any $x, y \in X$ and each $\alpha \in \Gamma$,*

$$(y \wedge x)\alpha x = y\alpha x.$$

Proof. Since $y \wedge x \leq y$, by Proposition 3.5 (1), we have

$$(3.7) \quad y\alpha x \leq (y \wedge x)\alpha x.$$

On the other hand, by Corollary 3.7 (2), we get

$$(3.8) \quad (y \wedge x)\alpha x = [(x\alpha y)\beta y]\alpha x \leq y\alpha x.$$

Thus $(y \wedge x)\alpha x = y\alpha x$. □

We obtain a characterization of a Γ - KU -algebra.

Theorem 3.11. *Let X be a set with a constant 0 and let Γ be a nonempty set. Then X is a Γ - KU -algebra if and only if it satisfies axioms (ΓKU_1) , (ΓKU_4) and the following condition: for any $x, y \in X$ and any $\alpha, \beta \in \Gamma$,*

$$(3.9) \quad (y\alpha 0)\beta x = x.$$

Proof. (\Rightarrow): The proof is straightforward from the axioms (ΓKU_2) and (ΓKU_3) .

(\Leftarrow): Suppose the necessary conditions hold, let $x \in X$ and let $\alpha, \beta \in \Gamma$. Then from the axiom (ΓKU_3) , we get $(x\alpha 0)\beta[(0\alpha 0)\beta(x\alpha 0)] = 0$. On the other hand, by (3.9), $(x\alpha 0)\beta(x\alpha 0) = 0$. Again by (3.9), $x\alpha 0 = 0$. Thus the axiom (ΓKU_3) holds. By combining (3.9) and the axiom (ΓKU_3) , $0\alpha x = x$. So the axiom (ΓKU_2) holds. Hence X is a Γ - KU -algebra. □

4. SPECIAL Γ - KU -ALGEBRAS

In this section, we define some special Γ - KU -algebras, for example, positive implicative [resp. implicative and commutative] Γ - KU -algebras and obtain some of their properties (including characterizations) respectively and a relationship among them.

Definition 4.1. A Γ - KU -algebra X is said to be *positive implicative*, if it satisfies the following axiom: for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$,

$$(4.1) \quad (z\alpha x)\beta(z\alpha y) = z\beta(x\alpha y).$$

It is obvious that if X is a positive implicative Γ - KU -algebra, then X_α is a positive implicative KU -algebra for each $\alpha \in \Gamma$.

Example 4.2. Let $X = \{0, 1, 2, 3\}$, let $\Gamma = \{\alpha, \beta\}$ and let the ternary operation be defined by the table:

α	0	1	2	3	β	0	1	2	3
0	0	1	2	3	0	0	1	2	3
1	0	0	2	3	1	0	0	2	3
2	0	1	0	3	2	0	0	0	3
3	0	0	2	0	3	0	0	2	0

Table 4.1

Then we can easily check that X is a positive implicative Γ - KU -algebra. Moreover, we confirm that X_α and X_β are positive implicative KU -algebras.

Proposition 4.3. *In a Γ - KU -algebra X , the following holds: for any $x, y \in X$ and any $\alpha, \beta \in \Gamma$,*

$$(4.2) \quad (x\alpha y)\beta((y\alpha x)\beta x) \leq (((x\alpha y)\beta y)\alpha x)\beta x.$$

Proof.
$$\begin{aligned} & [(((x\alpha y)\beta y)\alpha x)\beta x]\alpha[(x\alpha y)\beta((y\alpha x)\beta x)] \\ &= (x\alpha y)\beta[(((x\alpha y)\beta y)\alpha x)\beta x]\alpha((y\alpha x)\beta x) \text{ [By the identity (3.3)]} \\ &\leq (x\alpha y)\beta[(((x\alpha y)\beta y)\alpha((y\alpha x)\beta x))] \text{ [By Corollary 3.7 (2)]} \\ &\leq (x\alpha y)\beta(x\alpha y) \text{ [By Corollary 3.7 (2)]} \\ &= 0. \text{ [By (3.1)]} \end{aligned}$$

Then by Proposition 3.3 (2) and (3), we have

$$[(((x\alpha y)\beta y)\alpha x)\beta x]\alpha[(x\alpha y)\beta((y\alpha x)\beta x)] = 0.$$

Thus the inequality (4.2) holds. □

We have a characterization of a positive implicative Γ - KU -algebra.

Theorem 4.4. *Let X be a Γ - KU -algebra. Then the followings are equivalent:*

- (1) X is positive implicative,
- (2) $x\alpha y = x\alpha(x\beta y)$ for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$.

Proof. (1) \Rightarrow (2): The proof follows from the axiom (ΓKU_2), (3.1) and the identity (4.1).

(2) \Rightarrow (1): Suppose the condition (2) holds, let $x, y, z \in X$ and let $\alpha, \beta \in \Gamma$.

Then we have

$$\begin{aligned} & [z\alpha(x\beta y)]\alpha[(z\beta x)\alpha(z\beta y)] \\ &= [z\alpha(x\beta y)]\alpha[(z\beta x)\alpha(z\beta(z\alpha y))] \text{ [By (2)]} \\ &\leq [z\alpha(x\beta y)]\alpha[x\alpha(z\beta y)] \text{ [By Corollary 3.8]} \\ &= [z\alpha(x\beta y)]\alpha[z\alpha(x\beta y)] \text{ [By Proposition 3.6]} \\ &= 0. \text{ [By (3.1)]} \end{aligned}$$

Thus $(z\beta x)\alpha(z\beta y) \leq z\alpha(x\beta y)$. The proof of the converse inequality is easy. So $(z\beta x)\alpha(z\beta y) = z\alpha(x\beta y)$. Hence X is positive implicative. □

We give another characterization of a Γ - KU -algebra.

Theorem 4.5. *Let X be a Γ - KU -algebra. Then the followings are equivalent: for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$,*

- (1) X is positive implicative,

- (2) $z\alpha(x\beta y) = 0$ implies $(z\alpha x)\beta(z\alpha y) = 0$,
- (3) $y\alpha(y\beta x) = 0$ implies $y\alpha x = 0$.

Proof. (1) \Rightarrow (2): The proof is straightforward.

(2) \Rightarrow (3): Suppose (2) holds and $y\alpha(y\beta x) = 0$ for any $x, y \in X$ and any $\alpha, \beta \in \Gamma$. Then we have

$$\begin{aligned} y\alpha x &= 0\beta(y\alpha x) \text{ [By the axiom } (\Gamma KU_2)] \\ &= (y\alpha y)\beta(y\alpha x) \text{ [By (3.1)]} \\ &= 0. \text{ [By (2)]} \end{aligned}$$

Thus (3) holds.

(3) \Rightarrow (1): Suppose (3) holds. For any $x, y \in X$ and any $\alpha, \beta \in \tau$, let $u = y\beta(y\alpha x)$. Then we have

$$\begin{aligned} y\alpha(y\beta(u\alpha x)) &= y\alpha(u\beta(y\alpha x)) \text{ [By Proposition 3.6]} \\ &= u\alpha(y\beta(y\alpha x)) \text{ [By Proposition 3.6]} \\ &= (y\beta(y\alpha x))\alpha(y\beta(y\alpha x)) \\ &= 0. \text{ [By (3.1)]} \end{aligned}$$

Thus by the hypothesis and (3.1), we get

$$0 = y\beta(u\alpha x) = y\beta(y\beta(y\alpha x)\alpha x) = (y\beta(y\alpha x))\beta(y\alpha x).$$

So $y\alpha x \leq y\beta(y\alpha x)$. On the other hand, from Proposition 3.6 and (3.1), it is obvious that $y\beta(y\alpha x) \leq y\alpha x$. Hence $y\beta(y\alpha x) = y\alpha x$. Therefore by Theorem 4.4, X is positive implicative. \square

Definition 4.6. A Γ - KU -algebra X is said to be *commutative*, if it satisfies the following axiom:

$$(4.3) \quad (y\alpha x)\beta x = (x\alpha y)\beta y, \text{ i.e., } x \wedge y = y \wedge x \text{ for any } x, y \in X \text{ and any } \alpha, \beta \in \Gamma.$$

We can easily see that if X is a commutative Γ - KU -algebra, then X_α is a commutative kU -algebra for each $\alpha \in \Gamma$.

Example 4.7. (1) Let $X = \{0, 1, 2, 3, 4\}$, let $\Gamma = \{\alpha, \beta\}$ and let the ternary operation be defined by the table:

α	0	1	2	3	4	β	0	1	2	3	4
0	0	1	2	3	4	0	0	1	2	3	4
1	0	0	1	1	3	1	0	0	1	1	3
2	0	1	0	3	4	2	0	1	0	3	4
3	0	0	0	0	1	3	0	0	0	0	1
4	0	0	0	0	0	4	0	0	0	0	0

Table 4.2

Then clearly, X is a Γ - KU -algebra but $(2\alpha 3)\beta 3 = 3 \neq 2 = (3\alpha 2)\beta 2$. Thus X is not commutative.

(2) Let $X = \{0, 1, 2, 3\}$, let $\Gamma = \{\alpha, \beta\}$ and let the the ternary operation be defined as the following table:

Then we can easily check that X is commutative Γ - KU -algebra.

The following is an immediate consequence of the axiom (ΓKU_4) and Definition 4.6.

α	0	1	2	3	β	0	1	2	3
0	0	1	2	3	0	0	1	2	3
1	0	0	2	3	1	0	0	2	3
2	0	1	0	2	2	0	1	0	3
3	0	1	0	0	3	0	1	2	0

Table 4.3

Theorem 4.8. *Let X be a Γ - KU -algebra. Then the followings are equivalent: for any $x, y \in X$ and any $\alpha, \beta \in \Gamma$,*

- (1) X is commutative,
- (2) $(y\alpha x)\beta x \leq (x\alpha y)\beta y$,
- (3) $((x\alpha y)\beta y)\alpha((y\alpha x)\beta x) = 0$.

We obtain a characterization of commutative Γ - BCK -algebras.

Theorem 4.9. *Let X be a Γ - KU -algebra. Then the followings are equivalent: for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$,*

- (1) $x \leq z$ and $y\alpha z \leq x\alpha z$ imply $x \leq y$,
- (2) $x, y \leq z$ and $y\alpha z \leq x\alpha z$ imply $x \leq y$,
- (3) $x \leq y$ implies $x = (x\alpha y)\beta y$,
- (4) X is commutative,
- (5) $x \leq y$ implies $((x\alpha y)\beta y)\alpha x = 0$.

Proof. (1) \Rightarrow (2): The proof is clear.

(2) \Rightarrow (3): Suppose $x \leq y$ and let $\alpha, \beta \in \Gamma$. Then by Corollary 3.7 (2), $(x\alpha y)\beta y \leq x$. Moreover, $((x\alpha y)\beta y)\alpha y \leq x\alpha y$. Thus by the hypothesis, $x \leq (x\alpha y)\beta y$. So $x = (x\alpha y)\beta y$.

(3) \Rightarrow (4): Suppose the condition (3) holds, let $x, y \in X$ and let $\alpha, \beta \in \Gamma$. From Corollary 3.7 (2), it is clear that $(x\alpha y)\beta y \leq x$. Then by the hypothesis, we have

$$(4.4) \quad (x\alpha y)\beta y = (((x\alpha y)\beta y)\alpha x)\beta x.$$

On the other hand, we get

$$\begin{aligned} & [(y\alpha x)\beta x]\alpha[(x\alpha y)\beta y] \\ &= [(y\alpha x)\beta x]\alpha[(((x\alpha y)\beta y)\alpha x)\beta x] \text{ [By (4.4)]} \\ &= [(((x\alpha y)\beta y)\alpha x)]\alpha[(y\alpha x)\beta x] \text{ [By Proposition 3.6]} \\ &\leq [(((x\alpha y)\beta y)\alpha x)]\alpha(y\beta x) \text{ [Corollary 3.7 (2)]} \\ &\leq y\alpha[(x\alpha y)\beta y] \text{ [By Proposition 3.3 (1)]} \\ &= (x\alpha y)\alpha(y\beta y) \text{ [By Proposition 3.6]} \\ &= (x\alpha y)\alpha 0 \text{ [By (3.1)]} \\ &= 0 \text{ [By } (\Gamma KU_3)] \end{aligned}$$

Thus $(x\alpha y)\beta y \leq (y\alpha x)\beta x$. So by Theorem 4.8, X is commutative.

(4) \Rightarrow (1): Suppose the condition (4) holds, and suppose $x \leq z$ and $y\alpha z \leq x\alpha z$ for any $x, y, z \in X$ and each $\alpha \in \Gamma$. Then clearly, $z\alpha x = 0$ and $(x\alpha z)\beta(y\alpha z) = 0$ for any $\beta \in \Gamma$. Thus we have

$$\begin{aligned} y\alpha x &= y\alpha(0\beta x) \text{ [By } (\Gamma KU_2)] \\ &= y\alpha[(z\alpha x)\beta x] \text{ [Since } z\alpha x = 0] \\ &= y\alpha[(x\alpha z)\beta z] \text{ [Since } X \text{ is commutative]} \\ &= (x\alpha z)\beta(y\alpha z) \text{ [By Proposition 3.6]} \end{aligned}$$

$$= 0.$$

So $x \leq y$. Hence (1) holds.

(3) \Leftrightarrow (5): The proof is obvious. □

The following is an immediate consequence of Theorem 4.9.

Theorem 4.10. *Let X be a Γ - KU -algebra. Then followings are equivalent: ,*

(1) X is commutative,

(2) $x\alpha(x\beta y) = y\alpha(y\beta(x\alpha(x\beta y)))$ for any $x, y \in X$ and any $\alpha, \beta \in \Gamma$

For a Γ - KU -algebra X and each $x \in X$, the set

$$A(x) = \{y \in X : y \leq x\}$$

is called an *initial section* of x .

Theorem 4.11. *a Γ - KU -algebra X is commutative if and only if for any $x, y \in X$,*

$$A(x) \cap A(y) = A(x \wedge y).$$

Proof. The proof follows from the property of \wedge and Theorem 4.8 (2). □

Definition 4.12. Let X be a Γ - KU -algebra. Then X is said to be *implicative*, if it satisfies the following condition:

$$(4.5) \quad x = (x\alpha y)\beta x \text{ for any } x, y \in X \text{ and any } \alpha, \beta \in \Gamma.$$

It is clear that if X is an implicative Γ - BCK -algebra, then X_α is an implicative BCK -algebra fore each $\alpha \in \Gamma$.

Example 4.13. (1) Let $X = \{0, 1, 2, 3\}$, let $\Gamma = \{\alpha, \beta\}$ and let the the ternary operation be defined as the following table:

α	0	1	2	3	β	0	1	2	3
0	0	1	2	3	0	0	1	2	3
1	0	0	2	2	1	0	0	2	2
2	0	1	0	3	2	0	1	0	3
3	0	0	2	0	3	0	2	2	0

Table 4.4

Then clearly, X is an implicative Γ - KU -algebra.

(2) Let $X = \{0, 1, 2, 3, 4\}$, let $\Gamma = \{\alpha, \beta\}$ and let the the ternary operation be defined as the following table:

α	0	1	2	3	4	β	0	1	2	3	4
0	0	1	2	3	4	0	0	1	2	3	4
1	0	0	2	1	4	1	0	0	2	1	4
2	0	1	0	0	4	2	0	1	0	3	4
3	0	0	0	0	4	3	0	0	1	0	4
4	0	0	0	0	0	4	0	0	0	0	0

Table 4.5

Then X is a Γ - KU -algebra. But it is neither implicative nor commutative.

We obtain a relationship among implicativeness, commutativity and positive implicativeness.

Theorem 4.14. *Let X be a Γ - KU -algebra. Then X is implicative if and only if it is commutative and positive implicative.*

Proof. Suppose X is implicative, let $x, y \in X$ and let $\alpha, \beta \in \Gamma$. Then by Proposition 3.6, we have

$$x\alpha y = [(x\alpha y)\beta x]\beta(x\alpha y) = x\alpha(x\beta y).$$

Thus by Theorem 4.4, X is positive implicative. On the other hand, we get

$$\begin{aligned} (y\alpha x)\beta x &= (y\alpha x)\beta((x\alpha y)\beta x) \text{ [By the hypothesis]} \\ &\leq (x\alpha y)\beta y. \text{ [By Proposition 3.3 (1)]} \end{aligned}$$

So by Theorem 4.8 (1), X is commutative.

Conversely, suppose the necessary conditions hold and let $x, y \in X$ and $\alpha, \beta \in \Gamma$. Then we have

$$\begin{aligned} [(x\alpha y)\beta x]\alpha x &= [x\beta(x\alpha y)]\beta(x\alpha y) \text{ [Since } X \text{ is commutative]} \\ &= [x\beta(x\alpha y)]\beta[x\beta(x\alpha y)] \text{ [By Theorem 4.4 (2)]} \\ &= 0. \text{ [By (3.1)]} \end{aligned}$$

Thus $x \leq (x\alpha y)\beta x$. On the other hand, by Proposition 3.3 (5), $(x\alpha y)\beta x \leq x$. So by Proposition 3.3 (3), $x = (x\alpha y)\beta x$. Hence X is implicative. This completes the proof. \square

Now we provide a sufficient condition of implicative Γ - KU -algebras.

Proposition 4.15. *Let X be a Γ - KU -algebra. Suppose the following holds: for any $x, y \in X$ and any $\alpha, \beta \in \Gamma$,*

$$(4.6) \quad (y\alpha x)\beta[(y\alpha x)\beta x] = (x\alpha y)\beta[(x\alpha y)\beta y].$$

Then X is implicative

Proof. Suppose the condition (4.6) holds, let $x, y \in X$ and let $\alpha, \beta \in \Gamma$. Then we have

$$\begin{aligned} y\alpha x &= 0\alpha[0\beta(y\alpha x)] \text{ [By the axiom } (\Gamma KU_2)] \\ &= [y\beta(x\alpha x)]\alpha[(y\beta(x\alpha x))\beta(y\alpha x)] \text{ [By the axiom } (\Gamma KU_3) \text{ and (3.1)]} \\ &= [x\beta(y\alpha x)]\alpha[(x\beta(y\alpha x))\beta(y\alpha x)] \text{ [By Proposition 3.6]} \\ &= (x\beta y)\alpha[(x\beta y)\beta(y\alpha x)] \text{ [Putting } y\alpha y = y] \\ &= [x\alpha(x\beta y)]\alpha(x\beta y) \text{ [By the condition (2)]} \\ &= [x\alpha(x\beta(x\alpha y))]\alpha[x\beta(x\alpha y)] \text{ [Since } y = x\alpha y] \\ &= (x\alpha y)\alpha[x\beta(x\alpha y)] \text{ [By Proposition 3.6 (6)]} \\ &= [x\alpha(x\beta(x\alpha y))]\alpha y \text{ [By the identity (3.1)]} \\ &= (x\alpha y)\alpha y. \text{ [By Proposition 3.6 (6)]} \end{aligned}$$

Thus X is positive implicative. On the other hand, we have

$$\begin{aligned} x\alpha(x\beta y) &= [x\alpha(x\beta y)]\alpha(x\beta y) \text{ [By Proposition 3.6 (6)]} \\ &= [y\alpha(y\beta x)]\alpha(y\beta x) \text{ [By the condition (2)]} \\ &= y\alpha(y\beta x). \text{ [By Proposition 3.6 (6)]} \end{aligned}$$

So X is commutative. Hence by Theorem 4.14, X is implicative. \square

5. SOME Γ -IDEALS OF Γ - KU -ALGEBRAS

In this section, we introduce the concepts of Γ - KU -ideals, positive implicative Γ - KU -ideals, implicative Γ - KU -ideals and commutative Γ - KU -ideals in Γ - KU -algebras and discuss some of their properties respectively and a relationship among them.

Definition 5.1. Let X be a Γ - KU -algebra and let A be a nonempty subset of X . Then A is called a Γ -subalgebra of X , if it satisfies the following condition:

$$(5.1) \quad x\alpha y \in A \text{ for any } x, y \in A \text{ and for each } \alpha \in \Gamma.$$

Example 5.2. Let X be the Γ - KU -algebra given in Example 3.2 (3), $\{0, 1, 2\}$ is a Γ -subalgebra of X .

Definition 5.3. Let X be Γ - KU -algebra and let I be a nonempty set of X . Then I is called a Γ -ideal (briefly, ΓI) of X , if it satisfies the following conditions: for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$,

$$(\Gamma I_1) \quad 0 \in I,$$

$$(\Gamma I_2) \quad \text{if } x\alpha y \in I \text{ and } x \in I, \text{ then } y \in I.$$

An ideal I is said to be *proper*, if $I \neq X$. It is obvious that X and $\{0\}$ are ideals of X . In particular, X is called a *trivial* Γ -ideal of X .

Example 5.4. (1) Consider the Γ - KU -algebra given in Example 3.4. Then $\{0, 2\}$ is a Γ -ideal but $\{0, 1\}$ not a Γ -ideal of X .

(2) Let X be the commutative Γ - KU -algebra given in Example 4.7 (2). Then we can easily see that X has only two ΓI s $\{0\}$ and X .

The following is an immediate consequence of Definition 5.3.

Proposition 5.5. Let I be a ΓI of a Γ - KU -algebra X and let $x \in I$. If $y \leq x$, then $y \in I$.

Proposition 5.6. Every ΓI of a Γ - KU -algebra X is a Γ -subalgebra of X .

Proof. Let I be a ΓI of X and let $x, y \in I, \alpha \in \Gamma$. Then by Proposition 3.6 and the axiom (ΓKU_3) , $y\beta(x\alpha y) = 0$. Thus $x\alpha y \leq y$. So by Proposition 5.5, $x\alpha y \in I$. Hence I is a Γ -subalgebra of X . \square

Definition 5.7. Let X be Γ - KU -algebra and let $a, b \in X$ and $\alpha \in \Gamma$. Then the subset $A_\alpha(a, b)$ of X is defined as follows:

$$A_\alpha(a, b) = \{x \in X : b\alpha x \leq a\}.$$

It is obvious that $0, a, b \in A_\alpha(a, b)$.

Example 5.8. Let X be the Γ - BCK -algebra in Example 3.4. Then clearly,

$$A_\alpha(1, 2) = X, \quad A_\beta(1, 2) = \{0, 1, 2\} = A_\gamma(1, 2).$$

We have a characterization of ΓI s of a Γ - KU -algebra.

Theorem 5.9. Let I be a nonempty subset of a Γ - KU -algebra X . Then I is a ΓI of X if and only if $A_\alpha(a, b) \subset I$ for any $a, b \in I$ and each $\alpha \in \Gamma$.

Proof. (\Rightarrow): Suppose I is a Γ of X and let $x \in A_\alpha(a, b)$. Then clearly, $b\alpha x \leq a$. Thus by Proposition 5.5, $b\alpha x \in I$. Since $b \in I$ and I is a Γ of X , $x \in I$. So $A_\alpha(a, b) \subset I$.

(\Leftarrow): Suppose the necessary condition holds. Since $I \neq \emptyset$, there is $a \in I$. Then by (3.1), $a\alpha 0 \leq a$. Thus $0 \in A_\alpha(a, a)$. Since $A_\alpha(a, a) \subset I$, $0 \in I$. So the condition (Γ_1) holds. Now let $b\beta a \in I$ and $b \in I$. Then by Corollary 3.7 (2), $(b\beta a)\alpha x \leq b$. Thus $x \in A_\alpha(b\beta a, b) \subset I$. So the condition (Γ_2) holds. Hence I is a Γ of X . \square

The following is an immediate consequence of Theorem 5.9.

Corollary 5.10. *I is a Γ of a Γ - KU -algebra X if and only if for any $a, b \in I$ and any $\alpha, \beta \in \Gamma$, $(b\beta a)\alpha x = 0$ implies $x \in I$.*

Definition 5.11. Let X be Γ - KU -algebra and let I be a nonempty set of X . Then I is called a *positive implicative Γ - KU -ideal* (briefly, $\text{PI}\Gamma\text{KUI}$) of X , if it satisfies the following conditions: for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$,

(Γ_1) $0 \in I$,

($\text{PI}\Gamma\text{KUI}_2$) if $z\alpha(x\beta y) \in I$ and $z\alpha x \in I$, then $z\alpha y \in I$.

It is obvious that X is a $\text{PI}\Gamma\text{KUI}$ of X .

Example 5.12. Let X be the Γ - BCK -algebra given in Example 4.13 (2). Then we can easily check that $\{0, 1, 3\}$ and $\{0, 1, 2, 3\}$ are $\text{PI}\Gamma\text{KUI}$ s of X . Furthermore, $\{0\}$, $\{0, 2\}$ and $\{0, 2, 4\}$ are Γ s but not $\text{PI}\Gamma\text{KUI}$ s of X .

Proposition 5.13. *Every $\text{PI}\Gamma\text{KUI}$ of Γ - KU -algebra X is a Γ of X but the converse is not true.*

Proof. Let I be a $\text{PI}\Gamma\text{KUI}$ of X . Suppose $x\alpha y \in I$ and $x \in I$ for any $x, y \in X$ and each $\alpha \in \Gamma$. Then clearly, $0\beta(x\alpha y) \in I$ and $0\alpha y \in I$. Thus by ($\text{PI}\Gamma\text{KUI}_2$), $x = 0\alpha x \in I$. So I is a Γ of X . See Example 5.12 for the converse. \square

We have a characterization of positive implicative Γ - KU -ideals.

Theorem 5.14. *Let I be a ΓKUI of a Γ - KU -algebra X . Then I is positive implicative if and only if the set $A_a = \{x \in X : a\alpha x \in I \text{ for each } \alpha \in \Gamma\}$ is a Γ of X for each $a \in X$.*

Proof. Suppose I is positive implicative and $x\alpha y \in A_a$, $x \in A_a$ for each $a \in X$ and each $\alpha \in \Gamma$. Then clearly, $a\beta(x\alpha y) \in I$ and $a\alpha y \in I$. Thus by the condition ($\text{PI}\Gamma\text{KUI}_2$), $a\alpha x \in I$. So $x \in A_a$. Hence A_a is a Γ of X .

Now suppose the necessary condition holds, and $z\alpha(x\beta y) \in I$ and $z\alpha y \in I$ for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$. Then clearly, $y\alpha x \in A_z$ and $x \in A_z$. Thus by the hypothesis, $y \in A_z$. So $z\alpha y \in I$. Hence I is positive implicative. \square

The following is an immediate consequence of Theorem 5.14.

Corollary 5.15. *If I is a $\text{PI}\Gamma\text{KUI}$ of a Γ - KU -algebra X , then for each $a \in X$, A_a is the least Γ of X such that $I \cup \{a\} \subset A_a$.*

We obtain a characterization of $\text{PI}\Gamma\text{KUI}$ s.

Theorem 5.16. *Let I be a nonempty subset of a Γ -KU-algebra X . Then the followings are equivalent:*

- (1) I is a P Γ KUI of X ,
- (2) I is a ΓI of X and $y\alpha(y\beta x) \in I$ implies $y\alpha x \in I$ for any $x, y \in X$ and $\alpha, \beta \in \Gamma$,
- (3) I is a ΓI of X and $z\alpha(y\beta x) \in I$ implies $(z\alpha y)\beta(z\alpha x) \in I$ for any $x, y, z \in X$ and $\alpha, \beta \in \Gamma$,
- (4) $0 \in I$, and $z\alpha[y\beta(y\alpha x)] \in I$ and $z \in I$ imply $y\alpha x \in I$ for any $x, y, z \in X$ and $\alpha, \beta \in \Gamma$.

Proof. (1) \Rightarrow (2): Suppose I is a P Γ KUI of X . Then by Proposition 5.13, I is a ΓI of X . Now suppose $y\alpha(y\beta x) \in I$ for any $x, y \in X$ and $\alpha, \beta \in \Gamma$. From (3.1), $y\alpha y = 0 \in I$. Then by (P Γ KUI₂), $x\alpha y \in I$. Thus the condition (2) holds.

(2) \Rightarrow (3): Suppose the condition (2) holds and suppose $z\alpha(y\beta x) \in I$ for any $x, y, z \in X$ and $\alpha, \beta \in \Gamma$. Then we have

$$\begin{aligned} z\alpha[z\alpha((z\beta y)\alpha x)] &= z\beta[(z\beta y)\alpha(z\beta x)] \text{ [By (3.3)]} \\ &\leq z\alpha(y\beta x). \text{ [By Corollaries 3.8 and Corollaries 3.7 (2)]} \end{aligned}$$

Since I is a ΓI of X , by Proposition 5.5, $z\alpha[z\alpha((z\beta y)\alpha x)] \in I$. By the condition (2), $z\alpha[(z\beta y)\alpha x] \in I$. On the other hand, by Proposition 3.6, $z\alpha[(z\beta y)\alpha x] = (z\beta y)\alpha(z\beta x)$. Thus $(z\beta y)\alpha(z\beta x) \in I$. So the condition (3) holds.

(3) \Rightarrow (4): Suppose the condition (3) holds. Then clearly, $0 \in I$. Suppose $z\alpha[y\beta(y\alpha x)] \in I$ and $z \in I$ for any $x, y, z \in X$ and $\alpha, \beta \in \Gamma$. Then by Proposition 3.6, we get

$$z\alpha[y\beta(y\alpha x)] = y\alpha[y\beta(z\alpha x)].$$

Thus $y\alpha[y\beta(z\alpha x)] \in I$. On the other hand, from Proposition 3.6, (3.1) and the condition (3), we have

$$z\beta(y\alpha x) = y\beta(z\alpha x) = (y\alpha y)\beta(z\alpha x) \in I.$$

Since I is a ΓI of X and $z \in I$, $y\alpha x \in I$. So the condition (4) holds.

(4) \Rightarrow (1): Suppose the condition (4) holds. Suppose $x\alpha y \in I$ and $x \in I$ for any $x, y \in X$ and each $\alpha \in \Gamma$. Then by the axiom (Γ KU₂), we get

$$x\alpha y = x\alpha[0\alpha(0\beta y)].$$

Thus $x\alpha[(y\alpha 0)\beta 0] \in I$ and $x \in I$. By the condition (4), $0\beta y \in I$. By the axiom (Γ KU₂), $0\beta y = y$. So $y \in I$. Hence I is a ΓI of X .

Now suppose $z\alpha(x\beta y) \in I$ and $z\alpha x \in I$ for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$. Then from Corollary 3.9 and Proposition 3.6, we have

$$(z\alpha y)\beta[z\alpha(z\alpha x)] \leq y\beta(z\alpha x) = z\beta(y\alpha x).$$

Since $z\alpha(x\beta y) \in I$, $(z\alpha y)\beta[z\alpha(z\alpha x)] \in I$. Since $z\alpha x \in I$, by the condition (4), $y\alpha x \in I$. Thus I is a P Γ KUI of X . This completes the proof. \square

Proposition 5.17. *Let I and J be ΓI s of a Γ -KU-algebra X such that $I \subset J$. If I is positive implicative, then so is J .*

Proof. Suppose $z\beta(x\alpha y) \in J$ and $z\alpha x \in J$ for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$. Let $u = z\beta(x\alpha y)$. Then from Proposition 3.6, (3.1) and the hypothesis, we have

$$z\alpha[x\beta(u\alpha y)] = u\alpha[z\beta(x\alpha y)] = 0 \in I.$$

Since I is positive implicative, by Theorem 5.16 (3), we get

$$(z\alpha x)\beta[z\alpha(u\alpha y)] \in I.$$

On the other hand, by Proposition 3.6, we have

$$(z\alpha x)\beta[z\alpha(u\alpha y)] = u\beta[(z\alpha x)\beta(z\alpha y)] = [z\beta(x\alpha y)]\beta[(z\alpha x)\beta(z\alpha y)].$$

Thus $[z\beta(x\alpha y)]\beta[(z\alpha x)\beta(z\alpha y)] \in I$. Since $I \subset J$, $[z\beta(x\alpha y)]\beta[(z\alpha x)\beta(z\alpha y)] \in J$. Since $z\beta(x\alpha y) \in J$ and J is a ΓI of X , $(z\alpha x)\beta(z\alpha y) \in J$. So by Theorem 5.16 (3), J is positive implicative. \square

From the following Theorem, we can see that in Γ - KU -algebras, the zero ΓI s play important roles.

Theorem 5.18. *Let X be a Γ - KU -algebra. Then the followings are equivalent:*

- (1) X is positive implicative,
- (2) $\{0\}$ is a $\text{PI}\Gamma\text{KUI}$ of X ,
- (3) every ΓI of X is positive implicative,
- (4) the set $A(a) = \{x \in X : x \leq a\}$ is a ΓI of X for each $a \in X$.

Proof. (1) \Rightarrow (2): Suppose X is positive implicative. It is obvious that $\{0\}$ is a ΓI of X . Suppose $y\beta(y\alpha x) \in \{0\}$ for any $x, y \in X$ and any $\alpha, \beta \in \Gamma$. Since X is positive implicative, by Theorem 4.4, $y\alpha x = y\alpha(y\beta x)$. Then by the hypothesis, $y\alpha x \in \{0\}$. Thus by Theorem 5.16 (2), $\{0\}$ is a $\text{PI}\Gamma\text{KUI}$ of X .

(2) \Rightarrow (3): The proof follows from Proposition 5.17.

(3) \Rightarrow (4): Suppose the condition (3) holds and $x\alpha y, y \in A(a)$ for each $a \in X$ and each $\alpha \in \Gamma$. Then clearly, $y\alpha x \leq a$ and $y \leq a$. Thus $a\beta(y\alpha x) = 0 \in \{0\}$ and $a\alpha y = 0 \in \{0\}$ for any $\beta \in \Gamma$. By the hypothesis, $\{0\}$ is positive implicative. So $a\alpha x \in \{0\}$, i.e., $a\alpha x = 0$, i.e., $x \leq a$. Hence $x \in A(a)$. Therefore $A(a)$ is a ΓI of X .

(4) \Rightarrow (1): Suppose the condition (4) holds and $y\beta(y\alpha x) = 0$ for any $x, y \in X$ and any $\alpha, \beta \in \Gamma$. Then clearly, $y\alpha x \leq y$, i.e., $y\alpha x \in A(y)$. By the condition (4), $A(y)$ is a ΓI of X . It is obvious that $y \in A(y)$. Thus $x \in A(y)$. So $y\alpha x = 0$. Hence by Theorem 4.5, X is positive implicative. \square

We have a characterization of a positive implicative Γ - KU -algebra by ΓI s.

Theorem 5.19. *Let X be a Γ - KU -algebra. Then X is positive implicative if and only if A_a is a ΓI of X for each ΓI of X and each $a \in X$.*

Proof. Suppose X is positive implicative, let I be any ΓI of X and let $a \in X$. Then by Theorem 5.18, I is a $\text{PI}\Gamma\text{KUI}$ of X . Thus by Theorem 5.14, A_a is a ΓI of X .

Conversely, suppose the necessary condition holds and let J be any ΓI of X . Suppose $z\alpha(x\beta y) \in J$ and $z\alpha x \in J$ for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$. Consider the set $A_z = \{u \in X : z\alpha u \in J\}$. Then clearly, $x\beta y \in A_z$ and $x \in A_z$. Since A_z is a ΓI of X , $y \in A_z$. Thus $z\alpha y \in J$. So J is positive implicative. Hence by Theorem 5.18, X is positive implicative. \square

Definition 5.20. Let X be Γ - KU -algebra and let I be a nonempty subset of X . Then I is called an *implicative Γ - KU -ideal* (briefly, $\text{I}\Gamma\text{KUI}$) of X , if it satisfies the following conditions: for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$,

$$(\Gamma I_1) 0 \in I,$$

($\Pi\Gamma\text{KUI}_2$) if $z\alpha[(x\beta y)\alpha x] \in I$ and $z \in I$, then $x \in I$.

For any Γ - KU -algebra X , it is obvious that X is always an $\Pi\Gamma\text{KUI}$ of X which is called the *trivial implicative Γ - KU -ideal*.

We can easily show that every ΓI of an implicative Γ - KU -algebra X is implicative.

Example 5.21. Let X be the Γ - KU -algebra given in Example 4.13 (2). Then we can easily check that $\{0, 1, 2, 3\}$ is an $\Pi\Gamma\text{KUI}$ of X . Furthermore, $\{0\}$ is a ΓI of X but not implicative, since $0\alpha[(1\beta 2)\alpha 1] \in \{0\}$ and $0 \in \{0\}$ but $1 \notin \{0\}$.

Proposition 5.22. *Every $\Pi\Gamma\text{KUI}$ is a ΓI but the converse is not true.*

Proof. The proof is straightforward from Definitions 5.3 and 5.20. See Example 5.21) for the converse. \square

Proposition 5.23. *Every $\Pi\Gamma\text{KUI}$ is positive implicative but the converse is not true.*

Proof. Let I be an $\Pi\Gamma\text{KUI}$ of a Γ - KU -algebra X and $z\alpha(y\beta x)$, $z\beta y \in I$ for any x, y in X and any $\alpha, \beta \in \Gamma$. Then we get

$$\begin{aligned} (z\beta y)\alpha[z\alpha(z\beta x)] &\leq y\alpha(z\beta x) \text{ [By Corollary 3.7]} \\ &= z\alpha(y\beta x). \text{ [Proposition 3.6]} \end{aligned}$$

Since $z\alpha(y\beta x) \in I$, by Proposition 5.5, $(z\beta y)\alpha[z\alpha(z\beta x)] \in I$. Since $z\beta y \in I$ and I is a ΓI of X by Proposition 5.22, $z\alpha(z\beta x) \in I$. On the other hand, we have

$$\begin{aligned} [z\alpha(z\beta x)]\alpha(z\beta x) &= z\alpha[(z\alpha(z\beta x))\alpha x] \text{ [By Proposition 3.6]} \\ &= z\alpha(z\beta x) \in I. \text{ [By Proposition 3.10]} \end{aligned}$$

Thus $0\beta[(z\alpha(z\beta x))\alpha(z\beta x)] \in I$. Since $0 \in I$ and I is implicative, $z\beta x \in I$. So I is positive implicative.

In Example 5.12, $\{0, 1, 3\}$ is positive implicative but not implicative. \square

We obtain a condition for a ΓI to become a $\Pi\Gamma\text{KUI}$.

Theorem 5.24. *Let I be a ΓI of a Γ - KU -algebra X . Then I is implicative if and only if the following holds:*

$$(5.2) \quad (x\alpha y)\beta x \in I \text{ implies } x \in I \text{ for any } x, y \in X \text{ and any } \alpha, \beta \in \Gamma.$$

Proof. Suppose I is implicative and $(x\alpha y)\beta x \in I$ for any $x, y \in X$ and any $\alpha, \beta \in \Gamma$. It is obvious that $0\beta[(x\alpha y)\beta x] \in I$ and $0 \in I$. Then by the hypothesis, $x \in I$. Thus (5.2) holds.

The proof of the converse is easy. \square

Now we obtain a condition for a $\Pi\Gamma\text{KUI}$ to become a $\Pi\Gamma\text{KUI}$.

Theorem 5.25. *Let I be a $\Pi\Gamma\text{KUI}$ of a Γ - KU -algebra X . Then I is implicative if and only if the following holds:*

$$(5.3) \quad (x\alpha y)\beta y \in I \text{ implies } (y\alpha x)\beta x \in I \text{ for any } x, y \in X \text{ and any } \alpha, \beta \in \Gamma.$$

Proof. Suppose I is implicative and $(x\alpha y)\beta y \in I$ for any $x, y \in X$ and any $\alpha, \beta \in \Gamma$. Then by Corollary 3.7 (2), $(y\alpha x)\beta x \leq y$. Thus by Proposition 3.5 (1), $x\beta y \leq x\beta[(y\alpha x)\beta x]$. Furthermore, we get

$$\begin{aligned} [x\beta((y\alpha x)\beta x)]\alpha[(y\alpha x)\beta x] &\leq (x\beta y)\alpha[(y\alpha x)\beta x] \text{ [By Proposition 3.5 (1)]} \\ &= (y\alpha x)\beta[(x\alpha y)\beta x]. \text{ [By Proposition 3.6]} \end{aligned}$$

$$\leq (x\alpha y)\beta y. \text{ [By Proposition 3.3 (1)]}$$

Since I is a ΓI of X by the hypothesis and Proposition 5.22, we get

$$0\beta([x\beta((y\alpha x)\beta x)]\alpha[(y\alpha x)\beta x]) \in I.$$

Since $0 \in I$, by the condition (ΓKUI_2) , $y\alpha(x\beta x) \in I$. So (5.3) holds.

Conversely, suppose necessary condition (5.3) holds, and $z\beta[(x\alpha y)\beta x] \in I$ and $z \in I$. Since I is positive implicative, by Proposition 5.13, I is a ΓI of X . Then $(x\alpha y)\beta x \in I$. By Proposition 3.6 (3), we have

$$(x\beta y)\beta[(x\alpha y)\beta y] \leq (x\alpha y)\beta x \in I.$$

Thus $(x\beta y)\beta[(x\alpha y)\beta y] \in I$. Since I is positive implicative, by Theorem 5.16 (2), $(x\alpha y)\beta y \in I$. By the condition (5.3), we get

$$(5.4) \quad (y\alpha x)\beta x \in I.$$

Furthermore, from (3.1) and the axiom (ΓKU_3) , we have

$$z\beta(y\alpha x) \leq y\alpha x \leq (y\alpha x)\beta x \in I.$$

So $z\beta(y\alpha x) \in I$. Since $z \in I$ and I is a ΓI of X , $y\alpha x \in I$. By the condition (5.4), $x \in I$. Hence I is implicative. \square

We obtain a similar consequence to Proposition 5.17.

Proposition 5.26. *If I is an ΓKUI of a Γ - KU -algebra X , then every ΓI containing I is implicative.*

Proof. Suppose I is implicative and let J be any ΓI of X such that $I \subset J$. From Proposition 5.23, it is obvious that I is positive implicative. By Proposition 5.17, J is positive implicative. To prove that I is implicative, it is sufficient to prove that J satisfies the condition (5.3). Suppose $(x\beta y)\alpha y \in J$ for any $x, y \in X$ and any $\alpha, \beta \in \Gamma$ and let $u = (x\beta y)\alpha y$. Then clearly, $u\alpha[(x\beta y)\alpha y] = 0 \in I$. Since I is positive implicative, by Theorem 5.16 (3) and Proposition 3.6, we have

$$[u\alpha(y\beta x)]\alpha(u\beta x) = [y\alpha(u\beta x)]\alpha(u\beta x) \in I.$$

Since I is implicative, by the condition (5.3), $[(u\beta x)\alpha y]\beta y \in I$. Since $I \subset J$, $[(u\beta x)\alpha y]\beta y \in J$. On the other hand, by Corollary 3.7 (2), $[(u\beta x)\alpha y]\beta y \leq u\beta x$ and $(x\beta y)\alpha y \leq x$. Thus we get

$$\begin{aligned} [((u\beta x)\alpha y)\beta y]\alpha[(x\beta y)\alpha y] &\leq (x\beta y)\alpha[(u\beta x)\alpha y] \text{ [By Proposition 3.3 (1)]} \\ &\leq (u\beta x)\alpha x \text{ [By Proposition 3.5 (1)]} \\ &= [(x\beta y)\alpha y]\beta x \text{ [Since } u = x\alpha(x\beta y)\text{]} \\ &\leq (x\beta y)\alpha y \in J. \end{aligned}$$

So $[(u\beta x)\alpha y]\beta y \in J$. Since $[(u\beta x)\alpha y]\beta y \in J$, $(x\beta y)\alpha y \in J$. Hence by Theorem 5.25, J is implicative. \square

Now we obtain a similar consequence of Theorem 5.18.

Theorem 5.27. *Let X be a Γ - KU -algebra. The the followings are equivalent:*

- (1) $\{0\}$ is implicative,
- (2) every ΓI of X is implicative,
- (3) $A(a)$ is implicative for each $a \in X$,

(4) X is implicative.

Proof. (1) \Leftrightarrow (2): The proof follows from Proposition 5.26.

(2) \Leftrightarrow (3): The proof is straightforward from Proposition 5.23 and Theorem 5.18.

(4) \Rightarrow (1): The proof is obvious.

(1) \Rightarrow (4): Suppose $\{0\}$ is implicative. Then by Proposition 5.23, $\{0\}$ is positive implicative. By Theorem 5.18, $A((x\beta y)\alpha x)$ is a Γ I of X for any $x, y, z \in X$. By the hypothesis, $A((x\beta y)\alpha x)$ is implicative. It is clear that $(x\beta y)\alpha x \in A((x\beta y)\alpha x)$. Thus $x \in A((x\beta y)\alpha x)$. So $(x\beta y)\alpha x \leq x$. Note that $x \leq (x\beta y)\alpha x$. Hence $x = (x\beta y)\alpha x$. Therefore X is implicative. \square

Definition 5.28. Let X be Γ - KU -algebra and let I be a nonempty subset of X . Then I is called a *commutative Γ - KU -ideal* (briefly, CFKUI) of X , if it satisfies the following conditions: for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$,

(ΓI_1) $0 \in I$,

(CFKUI_2) if $z\alpha(y\beta x)$, $z \in I$, then $[(x\alpha y)\beta y]\alpha x \in I$.

It is obvious that X is always a CFKUI of a Γ - KU -algebra X which is called the *trivial commutative Γ - KU -ideal*.

Example 5.29. Let X be the Γ - KU -algebra given in Example 4.13 (2). Then we can easily see that $\{0, 4\}$ is commutative but not positive implicative, $\{0, 1, 3\}$ is positive implicative but not commutative and $\{0, 1, 2, 3\}$ is implicative.

Proposition 5.30. Every CFKUI of a Γ - KU -algebra X is a Γ I of X but the converse is not true.

Proof. Let I be any CFKUI of X and $y\alpha x \in I$ and $y \in I$ for any $x, y \in X$ and each $\beta \in \Gamma$. Then clearly, $y\alpha(0\beta x) \in I$ for each $\alpha \in \Gamma$. Since I is commutative, $x = [(x\alpha 0)\beta 0]\alpha x \in I$. Then I is a Γ I of X . See Example 5.29 for the converse. \square

We have an equivalent condition of CFKUI s.

Theorem 5.31. Let X be a Γ - KU -algebra and let I be a Γ I of X . Then I is commutative if and only if it satisfies the following condition:

(5.5) $y\alpha x \in I$ implies $[(x\alpha y)\beta y]\alpha x \in I$ for any $x, y \in X$ any $\alpha, \beta \in \Gamma$.

Proof. Suppose I is commutative and $y\alpha x \in I$ for any $x, y \in X$ and each $\alpha \in \Gamma$. Then clearly, $0\beta(y\alpha x) \in I$ for any $\beta \in \Gamma$ and $0 \in I$. Thus by the condition (CFKUI_2), $[(x\alpha y)\beta y]\alpha x \in I$. So the condition (5.5) holds.

Conversely, suppose the condition (5.5) holds and $z\beta(y\alpha x)$, $z \in I$ for any $x, y, z \in X$ and any $\alpha, \beta \in \Gamma$. Since I is a Γ I of X , $y\alpha x \in I$. Then by the condition (5.5), $[(x\alpha y)\beta y]\alpha x \in I$. Thus I is commutative. \square

We obtain a similar consequence of Theorem 4.14 for Γ I's.

Theorem 5.32. Let X be a Γ - KU -algebra and let I be a nonempty subset of X . Then I is implicative if and only if it is both commutative and positive implicative.

Proof. Suppose I is implicative. Then by Proposition 5.23, I is positive implicative. It is sufficient to prove that I is commutative.

Suppose $y\alpha x \in I$ for any $x, y \in X$ and each $\alpha \in \Gamma$. From (3.1) and the axiom (ΓKU_3) , $[(x\alpha y)\beta y]\alpha x \leq x$ for each $\beta \in \Gamma$. Then $y\alpha x \leq y\beta[(x\alpha y)\beta y]\alpha x$. Let $u = [(x\alpha y)\beta y]\alpha x$. Then we have

$$\begin{aligned} (u\beta y)\alpha u &= [(((x\alpha y)\beta y)\alpha x)\beta y]\alpha [((x\alpha y)\beta y)\alpha x] \\ &\leq (x\alpha y)\alpha [((x\alpha y)\beta y)\alpha x] \\ &= [(x\alpha y)\beta y]\alpha [(x\alpha y)\alpha x] \\ &\leq y\alpha x \in I. \end{aligned}$$

Thus $u\beta(y\alpha u) \in I$. Since I is implicative, by Theorem 5.24, $u \in I$, i.e., $[(x\alpha y)\beta y]\alpha x \in I$. So by Theorem 5.31, I is commutative.

Conversely, suppose the necessary condition holds and $(x\alpha y)\beta x \in I$ for any $x, y \in X$ and each $\alpha \in \Gamma$. It is obvious that

$$(x\alpha y)\beta[(x\alpha y)\beta y] \leq (x\alpha y)\beta x \in I.$$

Then $(x\alpha y)\beta[(x\alpha y)\beta y] \in I$. Since I is positive implicative, by Theorem 5.16 (2), we have

$$(5.6) \quad (x\alpha y)\beta y \in I.$$

Furthermore, by Propositions 3.3 (1) and 3.6, we have

$$y\beta x \leq (x\alpha y)\beta x.$$

Since $(x\alpha y)\beta x \in I$, $y\beta x \in I$, i.e., $y\alpha x \in I$. Since I is commutative, by Theorem 5.31,

$$(5.7) \quad [(x\alpha y)\beta y]\alpha x \in I.$$

Thus by (5.6) and (5.7), $x \in I$. So I is implicative. \square

We obtain a similar consequence of Proposition 5.17 for ΓKU s.

Proposition 5.33. *Let I and J be ΓKU s of a Γ - KU -algebra X such that $I \subset J$. If I is commutative, then so is J .*

Proof. Suppose I is commutative and $y\alpha x \in J$ for any $x, y \in X$ and each $\alpha \in \Gamma$. In order to show that J is commutative, it is sufficient to show that $[(x\alpha y)\beta y]\alpha x \in J$ by using Theorem 5.31. Let $u = y\alpha x$. Then we get

$$y\beta(u\alpha x) = u\beta(y\alpha x) = 0 \in I.$$

Since I is commutative, by Theorem 5.31, we have

$$[((u\alpha x)\beta y)\alpha y]\beta(u\alpha x) \in I.$$

By Proposition 3.6, we have

$$[((u\alpha x)\beta y)\alpha y]\beta(u\alpha x) = u\beta[(((u\alpha x)\beta y)\alpha y)\alpha x] \in I.$$

Since $I \subset J$, $u\beta[(((u\alpha x)\beta y)\alpha y)\alpha x] \in J$. Since J is a ΓI of X and $u \in J$, $[((u\alpha x)\beta y)\alpha y]\alpha x \in J$. On the other hand, from Proposition 3.3 (1), (1) and (ΓKU_3) , we get

$$\begin{aligned} [(((u\alpha x)\beta y)\alpha y)\alpha x]\beta [((x\alpha y)\beta y)\alpha x] &\leq [(x\alpha y)\beta y]\alpha [((u\alpha x)\beta y)\alpha y] \\ &\leq [(u\alpha x)\beta y]\alpha (x\beta y) \\ &\leq x\beta(u\alpha x) \\ &= u\beta(x\alpha x). \\ &= 0 \in J. \end{aligned}$$

Thus $[(x\alpha y)\beta y]\alpha x \in J$. So by Theorem 5.31, J is implicative. \square

Finally, we obtain a characterization of commutative Γ - KU -algebras.

Theorem 5.34. *Let X be a Γ - KU -algebra. The the followings are equivalent:*

- (1) $\{0\}$ is commutative,
- (2) every ΓI of X is commutative,
- (3) X is commutative.

Proof. (1) \Leftrightarrow (2): The proof is clear from Proposition 5.33.

(1) \Leftrightarrow (3): The proof follows from Theorem 4.9. □

6. TOPOLOGICAL STRUCTURES ON Γ - KU -ALGEBRAS

We recall some terms and notations related for a general topology (See [32, 33]). For a subset A of a topological space (X, τ) , the closure and the interior of A are denoted by $cl_\tau(A)$, $cl(A)$ or \bar{A} and $int_\tau(A)$, $int(A)$ or A° . A subfamily \mathcal{B} of τ is called a *base* for τ , if for each $U \in \tau$ either $U = \emptyset$ or there is $\mathcal{B}' \subset \mathcal{B}$ such that $U = \bigcup \mathcal{B}'$. A subset A of X is called a *neighborhood* of $x \in X$, if there is $U \in \tau$ such that $x \in U \subset A$. The set of all neighborhoods of x write as $N_\tau(x)$ or $N(x)$ and $N(x)$ is called the *neighborhood filter* of $x \in X$. A subfamily $\mathcal{N}(x)$ of $N(x)$ is called a *fundamental system of neighborhoods* of x , if for each $U \in N(x)$ there is $V \in \mathcal{N}(x)$ such that $V \subset U$. In fact, $\mathcal{N}(x)$ is a filter base of $N(x)$. Moreover, it is well-known ([32]) that $N_\tau(x)$ satisfies the following properties:

- (N₁) $x \in U$ for each $U \in N_\tau(x)$,
- (N₂) if $U \in N_\tau(x)$ and $U \subset V \subset X$, then $V \in N_\tau(x)$,
- (N₃) if $U_1, U_2 \in N_\tau(x)$, then $U_1 \cap U_2 \in N_\tau(x)$,
- (N₄) if $V \in N_\tau(x)$, there is $W \in N_\tau(x)$ such that $V \in N_\tau(x)$ for each $y \in W$.

Furthermore, it is well-known (Proposition 1.1.2, [32]) that for each $x \in X$ if $\mathcal{B}(x)$ be a set of subsets of X satisfying the properties (N₁)–(N₄), then a unique topology on X such that $\mathcal{B}(x) = N_\tau(x)$, where

$$\tau = \{V \subset X : \forall x \in V, \exists U \in \mathcal{B}(x) \text{ such that } U \subset V\}.$$

Definition 6.1. Let X be a KU -algebra and let τ be a topology on X . Then X is called a *topological KU -algebra* (briefly, TKU -algebra), if $*$: $(X \times X, \tau \times \tau) \rightarrow (X, \tau)$ is continuous, i.e., for any $x, y \in X$ and each $W \in N(x * y)$ there are $U \in N(x)$ and $V \in N(y)$ such that $U * V \subset W$, where $U * V = \{x * y \in X : x \in U, y \in V\}$.

Definition 6.2. Let X be a Γ - KU -algebra and let τ be a topology on X . Then X is called a *topological Γ - KU -algebra* (briefly, $T\Gamma$ - KU -algebra), if a mapping f : $(X, \tau) \times \Gamma \times (X, \tau) \rightarrow (X, \tau)$ is continuous at each $(x, \alpha, y) \in X \times \Gamma \times X$, i.e., for each $\alpha \in \Gamma$, any $x, y \in X$ and each $W \in N(x\alpha y)$ there are $U \in N(x)$ and $V \in N(y)$ such that $U\alpha V \subset W$, where $U\alpha V \subset W = \{x\alpha y : x \in U, y \in V\}$.

It is clear that if X is a $T\Gamma$ - KU -algebra, then X_α is a TKU -algebra for each $\alpha \in \Gamma$.

Example 6.3. (1) Let $X = \{0, 1, 2, 3, 4\}$ be the Γ - KU -algebra given in Example 4.13 (2). Consider the topology τ on X given by:

$$\tau = \{\emptyset, \{4\}, \{0, 1, 2, 3\}, X\}.$$

Then we can easily check that (X, τ) is a $\mathbb{T}\Gamma$ - KU -algebra. Moreover, X_α and X_β are TKU -algebras.

(2) Let $X = \{0, 1, 2, 3\}$ be the Γ - KU -algebra given in Example 3.4 (1). Consider a topology τ on X given by:

$$\tau = \{\emptyset, \{0\}, \{0, 1\}, \{0, 2, 3\}, X\}.$$

Then we can easily see that (X, τ) is a $\mathbb{T}\Gamma$ - KU -algebra.

Proposition 6.4. *Let X be a $\mathbb{T}\Gamma$ - KU -algebra. If $\{0\}$ is open in X , then X is discrete.*

Proof. Let $x \in X$ and let $\alpha \in \Gamma$. Then clearly, $x\alpha x = 0 \in \{0\} \in N(0)$. Thus there are $U, V \in N(x)$ such that $U\alpha V = \{0\}$. Let $W = U \cap V$. Then $W\alpha W \subset U\alpha V = \{0\}$. Thus $W\alpha W = \{0\}$. Since $x \in U \cap V$, $x \in W$. So $W = \{x\}$ and W is open in X . Hence X is discrete. \square

The following is an immediate consequence of Proposition 6.4.

Corollary 6.5. *Let X be a $\mathbb{T}\Gamma$ - KU -algebra. If $\{0\}$ is open in X_α for each $\alpha \in \Gamma$, then X_α is discrete.*

Theorem 6.6. *Let X be a $\mathbb{T}\Gamma$ - KU -algebra. Then $\{0\}$ is closed in X if and only if X is Hausdorff.*

Proof. Suppose $\{0\}$ is closed in X , let $x, y \in X$ such that $x \neq y$ and let $\alpha \in \Gamma$. Then $x\alpha y \neq 0$ or $y\alpha x \neq 0$, say $x\alpha y \neq 0$. Since $\{0\}$ is closed in X and $x\alpha y \neq 0$, $\{0\}^c$ is open in X and $x\alpha y \in \{0\}^c$. Thus $\{0\}^c \in N(x\alpha y)$. Since X is a $\mathbb{T}\Gamma$ - KU -algebra, by Definition 6.2, there are $U \in N(x)$ and $V \in N(y)$ such that $U\alpha V \subset \{0\}^c$. So $U \cap V = \emptyset$. Hence X is Hausdorff.

Conversely, suppose X is Hausdorff and let $x \in \{0\}^c$. Then $x \neq 0$. By the hypothesis, there are $U \in N(x)$ and $V \in N(0)$ such that $U \cap V = \emptyset$. Thus $0 \notin U$. So $U \subset \{0\}^c$. Hence $\{0\}^c$ is open in X . Therefore $\{0\}$ is closed in X . \square

The following is an immediate consequence of Theorem 6.6.

Corollary 6.7. *Let X be a $\mathbb{T}\Gamma$ - KU -algebra. Then $\{0\}$ is closed in X_α if and only if X_α is Hausdorff for each $\alpha \in \Gamma$.*

Proposition 6.8. *Let X be a $\mathbb{T}\Gamma$ - KU -algebra and let A be open in X . If A is a Γ -subalgebra of X , then A is a $\mathbb{T}\Gamma$ - KU -algebra.*

Proof. Let τ be the topology on X and let τ_A be the subspace topology on A with respect to τ . Let $x, y \in A$ and let $\alpha \in \Gamma$. Since A is a Γ -subalgebra of X , $x\alpha y \in A$. Let $W_A \in N_{\tau_A}(x\alpha y)$, where $N_{\tau_A}(x\alpha y)$ denotes the neighborhood of $x\alpha y$ in the subspace (A, τ_A) of (X, τ) . Then there is $W \in N(x\alpha y)$ such that $W_A = A \cap W$. Since X is a $\mathbb{T}\Gamma$ - KU -algebra, there are $U \in N(x)$ and $V \in N(y)$ such that $U\alpha V \subset W$. Thus $U_A = A \cap U \in N_{\tau_A}(x)$ and $V_A = A \cap V \in N_{\tau_A}(y)$. It is clear that

$$U_A\alpha V_A = (A \cap U)\alpha(A \cap V) \subset W \text{ and } U_A\alpha V_A \subset A.$$

So $U_A\alpha V_A \subset A \cap W = W_A$. Hence A is a $\mathbb{T}\Gamma$ - KU -algebra. \square

Corollary 6.9. *Let X be a $\mathbb{T}\Gamma$ - KU -algebra and let A be open in X_α for each $\alpha \in \Gamma$. If A is a Γ -subalgebra of X_α , then A is a TKU -algebra.*

Proposition 6.10. *Let X be a Γ - KU -algebra and let I be open in X . If I is a ΓI of X , then I is closed in X .*

Proof. Let $x \in I^c$ and let $\alpha \in \Gamma$. Since $x\alpha x = 0 \in I$ and I is open, $I \in N(0)$. Since X is a Γ - KU -algebra, there is $U \in N(x)$ such that $U\alpha U \subset I$. Assume that $U \not\subset I^c$. Then there is $y \in X$ such that $y \in U \cap I$. It is obvious that $y\alpha z \in U\alpha U \subset I$ for each $z \in U$. Since I is a ΓI of X and $y \in I$, $z \in I$. Thus $U \subset I$. This is a contradiction. So $U \subset I^c$, i.e., I^c is open in X . Hence I is closed in X . \square

Corollary 6.11. *Let X be a Γ - KU -algebra and let I be open in X_α for each $\alpha \in \Gamma$. If I is a ΓI of X_α , then I is closed in X_α .*

Proposition 6.12. *Let X be a Γ - KU -algebra and let I be a ΓI of X . If $0 \in \text{int}(I)$, then I is open in X .*

Proof. Let $x \in I$ and let $\alpha \in \Gamma$. Since $0 \in \text{int}(I)$ and $x\alpha x = 0 \in I$, there is $W \in N(0) = N(x\alpha x)$ such that $W \subset I$. Since X is a Γ - KU -algebra, there are $U, V \in N(x)$ such that $U\alpha V \subset W \subset I$. It is obvious that $x\alpha y \in U\alpha V \subset I$ for each $y \in U$. Since I is a ΓI of X and $x \in I$, $y \in I$. Thus $U \subset I$. So I is open in X . \square

Corollary 6.13. *Let X be a Γ - KU -algebra and let I be a ΓI of X_α for each $\alpha \in \Gamma$. If $0 \in \text{int}(I)$, then I is open in X_α .*

In Proposition 6.12, when $0 \neq x \in \text{int}(I)$, I need not open in X (See Example 6.14).

Example 6.14. For a set $\Gamma = \{\alpha, \beta\}$, let $X = \{0, 1, 2, 3\}$ be a Γ - KU -algebra with the ternary operation be defined by the table:

α	0	1	2	3	β	0	1	2	3
0	0	1	2	3	0	0	1	2	3
1	0	0	2	2	1	0	0	1	3
2	0	0	0	3	2	0	0	0	3
3	0	1	2	0	3	0	2	1	0

Table 6.1

Consider a topology τ on X given by:

$$\tau = \{\emptyset, \{2\}, \{3\}, \{0, 1\}, \{2, 3\}, \{0, 1, 3\}, X\}.$$

Let $I = \{0, 3\}$. Then clearly, $3 \in \text{int}(I)$. But $I \notin \tau$.

Proposition 6.15. *Let X be Γ - KU -algebra. Then $\bigcap N(0) = \{0\}$ and thus $\bigcap \mathcal{N}(0) = \{0\}$.*

Proof. Assume that $0 \neq x \notin \bigcap N(0)$. Then clearly, there is $U \in N(0)$ such that $0 \in U$ but $x \notin U$. Thus $x \notin \bigcap N(0)$. This is a contradiction. So $\bigcap N(0) = \{0\}$. \square

Proposition 6.16. *Let (X, τ) be a Γ - KU -algebra and let $\mathcal{B}_1, \mathcal{B}_2$ be the families of subsets of X given by:*

$$\mathcal{B}_1 = \{U\alpha x : x \in X, \alpha \in \Gamma, U \in \mathcal{N}(0)\}, \mathcal{B}_2 = \{x\alpha U : x \in X, \alpha \in \Gamma, U \in \mathcal{N}(0)\},$$

where $U\alpha x = \{u\alpha x : u \in U\}$ and $x\alpha U = \{x\alpha u : u \in U\}$. Then \mathcal{B}_1 and \mathcal{B}_2 are bases for τ .

Proof. Let $x \in X$. Since $0 \in U \in \mathcal{N}(0)$, $0\alpha x = x$. Then $\bigcup \mathcal{B}_1 = X$. Suppose $B_1, B_2 \in \mathcal{B}_1$ and $z \in B_1 \cap B_2$. Then there are $U_1, U_2 \in \mathcal{N}(0)$ such that $B_1 = U_1\alpha x$, $B_2 = U_2\alpha x$ and $B_1 \cap B_2 = (U_1 \cap U_2)\alpha x$. Since $z \in B_1 \cap B_2$, there is $y \in U_1 \cap U_2$. Since $U_1, U_2 \in \mathcal{N}(0)$, $U_1 \cap U_2 \in \mathcal{N}(0)$. So there is $V \in \mathcal{N}(0)$ such that $y \in V \subset U_1 \cap U_2$. Hence $z = y\alpha x \in V\alpha x \in \mathcal{B}_1$. Therefore \mathcal{B}_1 is a base for τ . Similarly, we can prove that \mathcal{B}_2 is a base for τ . \square

Now in order to give a filter base on X generating a topology on a Γ - KU -algebra, let us define the subset $U(a)$ of X generated by each $a \in X$ and each subset U of X as follows:

$$U(a) = \{x \in X : x\alpha a \in U, a\alpha x \in U, \alpha \in \Gamma\}.$$

Proposition 6.17. *Let X be a Γ - KU -algebra. Suppose \mathcal{B} is a filter base on X satisfying the following condition:*

- (1) for each $u \in U \in \mathcal{B}$ there is $B \in \mathcal{B}$ such that $B(u) \subset U$,
- (2) for each $u \in U \in \mathcal{B}$ and each $\alpha \in \Gamma$ if $u\alpha x = 0$, then $x \in U$,
- (3) for each $U \in \mathcal{B}$ there is $B \in \mathcal{B}$ such that $B(b) \subset U$ for each $b \in B$, i.e., $B(B) \subset U$.

Then there is a unique topology τ on X such that $\mathcal{B} = \mathcal{N}_\tau(0)$ and (X, τ) is a Π - KU -algebra.

Proof. Let $\tau = \{O \in P(X) : \text{for each } a \in O \text{ there is } B \in \mathcal{B} \text{ such that } B(a) \subset O\}$. Then we can easily prove that τ is a topology on X . To accomplish to the proof, consider the following Claims.

Claim 1: $B(a) \in \tau$. Let $x \in B(a)$. Then $x\alpha a, a\alpha x \in B$ for each $\alpha \in \Gamma$. Thus by the condition (1), there are $B_1, B_2 \in \mathcal{B}$ such that $B_1(x\alpha a) \subset B$ and $B_2(a\alpha x) \subset B$. Since \mathcal{B} is a filter base on X , there is $U \in \mathcal{B}$ such that $U \in B_1 \cap B_2$. Let $x\alpha y, y\alpha x \in U$, i.e., $y \in U(x)$. By Proposition 3.3 (1), we have

$$(x\alpha a)\beta(y\alpha a) \leq y\alpha x, (y\alpha a)\beta(x\alpha a) \leq x\alpha y.$$

Then $(y\alpha x)\beta[(x\alpha a)\beta(y\alpha a)] = 0$, $(x\alpha y)\beta[(y\alpha a)\beta(x\alpha a)] = 0$. By the condition (2), $(x\alpha a)\beta(y\alpha a), (y\alpha a)\beta(x\alpha a) \in U$. Thus we get

$$y\alpha a \in U(x\alpha a) \subset B_1(x\alpha a) \subset B.$$

So $y\alpha a \in B$. Similarly, we can show that $a\alpha y \in U$. Hence $y \in U(a)$, i.e., $U(x) \subset B(a)$. Therefore $B(a) \in \tau$.

Claim 2: $\mathcal{B} = \mathcal{N}_\tau(0)$. Let $A \in \mathcal{B}$ and let $x \in A$. Since X is a Γ - KU -algebra, by the axiom (ΓKU_3), $x\alpha 0 = 0$. By the condition (2), $0 \in A$. By the condition (1), there is $B \in \mathcal{B}$ such that $B(0) \subset A$. Then by Claim 1, $B(0) \in \tau$. Thus $A \in \mathcal{N}_\tau(0)$. So $\mathcal{B} \subset \mathcal{N}_\tau(0)$. Hence by the condition (3), $\mathcal{B} \subset \mathcal{N}_\tau(0)$. It can be easily proved that $\mathcal{N}_\tau(0) \subset \mathcal{B}$. Therefore $\mathcal{B} = \mathcal{N}_\tau(0)$.

Claim 3: A mapping $f : (X, \tau) \times \Gamma \times (X, \tau) \rightarrow (X, \tau)$ is continuous at each $(x, \alpha, y) \in X \times \Gamma \times X$. Let $x, y \in X$, let $\alpha \in \Gamma$ and let $W \in \mathcal{N}_\tau(x\alpha y)$. Since $x\alpha y \in W$, by the condition (1), there is $W' \in \mathcal{B}$ such that $W'(x\alpha y) \subset W$. Since $W' \in \mathcal{B}$, by the condition (3), there is $B \in \mathcal{B}$ such that $B(b) \subset W'$ for each $b \in W'$.

Let $U = B(x)$, $V = B(y)$ and let $u \in U$, $v \in V$. Then we have

$$\begin{aligned} (x\alpha u)\beta[(u\alpha v)\beta(x\alpha y)] &= (u\alpha v)\beta[(x\alpha u)\beta(x\alpha y)] \text{ [By Proposition 3.6]} \\ &\leq (u\alpha v)\beta(u\alpha y) \text{ [By Corollary 3.8]} \\ &\leq v\alpha y. \text{ [By Corollary 3.8]} \end{aligned}$$

Thus $(\alpha y)\beta[(x\alpha u)\beta((u\alpha v)\beta(x\alpha y))] = 0$. Since $v\alpha y \in B$, by the condition (2), $(x\alpha u)\beta[(u\alpha v)\beta(x\alpha y)] \in B$. Similarly, we have $[(u\alpha v)\beta(x\alpha y)]\beta(x\alpha u) \in B$. So we get

$$(u\alpha v)\beta(x\alpha y) \in B(x\alpha u) \subset W', \text{ i.e., } (u\alpha v)\beta(x\alpha y) \in W'.$$

Similarly, $(x\alpha y)\beta(u\alpha v) \in W'$. Hence we have

$$u\alpha v \in W'(x\alpha y), \text{ i.e., } U\alpha V = B(x)\alpha B(y) \subset W'(x\alpha y) \subset W.$$

Therefore f is continuous. The proof of uniqueness for τ is easy. This completes the proof. \square

Example 6.18. (1) Let X be the Γ - KU -algebra and let \mathcal{I} be the collection of all Γ Is of X . Let $x \in I \in \mathcal{I}$. Then clearly, $I(x) \subset I$. Thus \mathcal{I} satisfies the conditions (1) and (3) in Proposition 6.17. Let $y \in I \in \mathcal{I}$ and suppose $y\alpha x = 0$. Then $y\alpha x = 0 \in I$. Thus $x \in I$. So \mathcal{I} satisfies the condition (2) in Proposition 6.17. So \mathcal{I} forms a filter base of X satisfying all the conditions in Proposition 6.17. Hence (X, τ) is a (X, τ) is a $\Gamma\Gamma$ - KU -algebra, where τ is the topology on X generated by \mathcal{I} .

(2) Let $X = \{0, 1, 2, 3\}$ be the Γ - KU -algebra given in Example 4.7 (2). Consider the family \mathcal{B} of subsets of X given by:

$$\mathcal{B} = \{\{0, 1\}, \{0, 2\}, \{0, 3\}, \{0, 1, 2\}, \{0, 1, 3\}, \{0, 2, 3\}\}.$$

Then we can easily check that \mathcal{B} is a filter base on X . Moreover, we have

$$\begin{aligned} \{0, 1\}(0) &= \{0, 1\}(1) = \{0, 1\}, \{0, 1\}(2) = \{2\}, \{0, 1\}(3) = \{3\}, \\ \{0, 2\}(0) &= \{0, 2\}(2) = \{0, 2\}, \{0, 2\}(1) = \{1\}, \{0, 2\}(3) = \{3\}, \\ \{0, 3\}(0) &= \{0, 3\}(3) = \{0, 3\}, \{0, 3\}(1) = \{1\}, \{0, 3\}(2) = \{2\}, \\ \{0, 1, 2\}(0) &= \{0, 1, 2\}(1) = \{0, 1, 2\}(2) = \{0, 1, 2\}, \{0, 1, 2\} = \{3\}, \\ \{0, 1, 3\}(0) &= \{0, 1, 3\}(1) = \{0, 1, 3\}(3) = \{0, 1, 3\}, \{0, 1, 3\}(2) = \{2\}, \\ \{0, 2, 3\}(0) &= \{0, 2, 3\}(2) = \{0, 2, 3\}(3) = \{0, 2, 3\}, \{0, 2, 3\}(1) = \{1\}. \end{aligned}$$

Thus \mathcal{B} is a filter base on X satisfying all the conditions in Proposition 6.17. So the topology τ on X generated by \mathcal{B} is given as follows:

$$\tau = \{\emptyset, \{0, 1\}, \{0, 2\}, \{0, 3\}, \{0, 1, 2\}, \{0, 1, 3\}, \{0, 2, 3\}, X\}.$$

Hence (X, τ) is a $\Gamma\Gamma$ - KU -algebra.

Lemma 6.19. *Let X be a Γ - KU -algebra and let τ be the topology on X generated by \mathcal{B} , where \mathcal{B} is a filter base on satisfying all the conditions in Proposition 6.17. Then for each $B \in \mathcal{B}$ and each $a \in X$,*

- (1) $B(a) \in N_\tau(a)$,
- (2) $B(A) = \bigcup_{a \in A} B(a) \in N_\tau(A)$ for each $A \in P(X)$.

Proof. The proof is straightforward. \square

Proposition 6.20. *Let X be a Γ - KU -algebra and let τ be the topology on X generated by \mathcal{B} , where \mathcal{B} is a filter base on X satisfying all the conditions in Proposition 6.17. Then for each $B \in \mathcal{B}$, $cl_\tau(A) = \bigcap_{B \in \mathcal{B}} B(A)$.*

Proof. Let $x \in cl_\tau(A)$ and let $B \in \mathcal{B}$. By Lemma 6.19 (1), $B(x) \in N_\tau(x)$. Then $B(x) \cap A \neq \emptyset$. Thus there is $a \in A$ such that $a\alpha x$, $x\alpha a \in B$ for each $\alpha \in \Gamma$. So $x \in B(a) \subset B(A)$, i.e., $x \in \bigcap_{B \in \mathcal{B}} B(A)$. Hence $cl_\tau(A) \subset \bigcap_{B \in \mathcal{B}} B(A)$. Conversely, let $x \in \bigcap_{B \in \mathcal{B}} B(A)$. Then $x \in U(A)$ for each $U \in \mathcal{B}$. Thus there is $a \in A$ such that $x \in B(a)$, i.e., $x\alpha a$, $a\alpha x \in B$ for each $\alpha \in \Gamma$. So $a \in B(x)$, i.e., $B(x) \cap A \neq \emptyset$. Hence $x \in cl_\tau(A)$, i.e., $\bigcap_{B \in \mathcal{B}} B(A) \subset cl_\tau(A)$. Therefore $cl_\tau(A) = \bigcap_{B \in \mathcal{B}} B(A)$. \square

Corollary 6.21. *Let (X, τ) be a TT - KU -algebra, where \mathcal{B} is a filter base on X satisfying all the conditions in Proposition 6.17 and τ is the topology on X generated by \mathcal{B} . Then every $B \in \mathcal{B}$ is closed in X , i.e., \mathcal{B} is a collection of clopen subsets of X .*

Proof. Let $B \in \mathcal{B}$. It is obvious that $B(B) \subset B$. Then by Proposition 6.20, $B \subset cl_\tau(B) = \bigcap_{U \in \mathcal{B}} U(B) \subset B(B) \subset B$. Thus $cl_\tau(B) = B$. So B is closed in X . From Proposition 6.17, it is clear that B is open in X . So B is clopen in X . \square

The following shows that every neighborhood of a compact set contains a neighborhood $B(A)$ for some $B \in \mathcal{B}$

Proposition 6.22. *Let A be a compact subset of a TT - KU -algebra. If U is a neighborhood of A , then there is $B \in \mathcal{B}$ such that $A \subset B(A) \subset U$.*

Proof. Suppose U is a neighborhood of A and let $a \in A$. Then there is $B_a \in \mathcal{B}$ such that $B_a \subset U$. Thus by the condition (3), there is $W_a \in \mathcal{B}$ such that $W_a(W_a) \subset B_a$. Since A is a compact subset of X and $A \subset \bigcup_{a \in A} W_a(a)$, there are $a_1, a_2, \dots, a_n \in A$ such that

$$(6.1) \quad A \subset W_{a_1}(a_1) \cup W_{a_2}(a_2) \cup \dots \cup W_{a_n}(a_n).$$

Now let $W = \bigcap_{i=1}^n W_{a_i}$ and let $a \in A$. Then by (6.1), there is $i \in \{1, 2, \dots, n\}$ such that $a \in W_{a_i}(a_i)$. Thus $a\alpha a_i$, $a_i\alpha a \in W_{a_i}$ for each $\alpha \in \Gamma$. Suppose $a\alpha y$, $y\alpha a \in W$ for each $y \in X$. By Proposition 3.3 (1), we have

$$(6.2) \quad (y\alpha a_i)\beta(a\alpha a_i) \leq a\alpha y \in W \text{ for each } \beta \in \Gamma.$$

Then $(y\alpha a_i)\beta(a\alpha a_i) \in W$. Thus we get

$$y\alpha a_i \in W_{a_i}(a\alpha a_i) \subset W_{a_i}(W_{a_i}) \subset B_{a_i}.$$

Similarly, $a_i\alpha y \in B_{a_i}$. So $y \in B_{a_i}(a_i) \subset U$ and $W(a) \subset U$. Hence $W(A) \subset U$. \square

The following is an immediate consequence of Proposition 6.22.

Corollary 6.23. *Let A be a compact subset of a TT - KU -algebra and let F is closed in X . If $A \cap F = \emptyset$, then there is $B \in \mathcal{B}$ such that $B(A) \cap B(F) = \emptyset$.*

7. CONCLUSIONS

By proposing positive implicative [resp. implicative and commutative] Γ - KU -algebras, we obtained some of their properties respectively and a relationship among them (See Theorem 4.14). Also, by defining positive implicative [resp. implicative and commutative] Γ - KU -ideals of a Γ - KU -algebra, we studied their various properties respectively and a relationship among them (See Theorem 5.32). Moreover, we discussed some topological structures on a Γ - KU -algebra.

In the future, we will use our proposed Γ - KU -algebras to address quotient Γ - KU -algebras, homomorphism problems, graph theory and Zariski topological structures. Furthermore, we want to study some ideals of a Γ - KU -algebra in the sense of the fuzzy set theory.

REFERENCES

- [1] K. Iséki and S. Tanaka, An introduction to the theory of BCK -algebras, *Math. Japon.* 23 (1) (1978) 1–26.
- [2] Y. Imai and K. Iséki, On axiom systems of propositional calculi, XIV, *Proceedings of the Japan Academy* 42 (1966) 19–22.
- [3] K. Iséki, On BCI -algebras, *Math. Seminar Notes* 8 (1980) 125–130.
- [4] K. Iséki and S. Tanaka, Ideal theory of BCK -algebras, *Math. Japon.* 23 (1978) 1–26.
- [5] J. Meng, Ideals in BCK -algebras, *Pure Appl. Math.* 2 (1986) 68–76.
- [6] J. Meng, Commutative ideals in BCK -algebras, *Pure Appl. Math.* 9 (1991) 49–53.
- [7] J. Meng, On ideals in BCK -algebras, *Math. Japon.* 40 (1994) 143–154.
- [8] C. H. Hoo, Filters and ideals in BCI -algebras, *Math. Japon.* 36 (1991) 987–997.
- [9] W. A. Dudek and X. Zhang, On ideals and congruences in BCC -algebras, *Czechoslovak Math. Journal* 48 (123) (1998) 21–29.
- [10] Dong Soo Lee and Dong Neun Ryu, Notes on topological BCK -algebras, *Scientiae Mathematicae* 1 (2) (1998) 231–235.
- [11] T. Roudabri and L. Torkzadeh, A topology on BCK -algebras via left and right stabilizers, *Iranian Journal of Mathematical Sciences and Informatics* 4 (2) (2009) 1–18.
- [12] Y. B. Jun, X. L. Xin and D. S. Lee, On topological BCI -algebras, *Inform. Sci.* 116 (1999) 253–261.
- [13] S. A. Ahn and S. H. Kwon, Topological properties in BCI -algebras, *Comm. Korean Math. Soc.* 23 (2) (2008) 169–1278.
- [14] C. Prabpayak and U. Leerawat, On ideals and congruences in KU -algebras, *Scientia Magna* 5 (1) (2009) 54–57.
- [15] C. Prabpayak and U. Leerawat, On Isomorphisms of KU -algebras, *Scientia Magna* 5 (3) (2009) 25–31.
- [16] Samy M. Mostafa, Mokhtar A. Abd-Elnaby and Moustafa M. M. Yousef, Fuzzy ideals of KU -algebras, *International Mathematical Forum* 6 (63) (2011) 3139–3149.
- [17] Naveed Yaqoob, Samy M. Mostafa, and Moin A. Ansari, On cubic KU -ideals of KU -algebras, *ISRN Algebra* 2013 (2013), Article ID 935905 10 pages, <http://dx.doi.org/10.1155/2013/935905>.
- [18] Samy M. Mostafa and Fatema F. Kareem, N -fold commutative KU -algebras, *International Journal of Algebra* 8 (6) (2014) 267–275.
- [19] Samy M. Mostafa, Abdelaziz E. Radwan, Fayza A. Ibrahim and Fatema F. Kareem, Topology spectrum of a KU -algebra, *Journal of New Theory* (8) (2015) 78–91.
- [20] Samy M. Mostafa, Abdelaziz E. Radwan, Fayza A. Ibrahim and Fatema F. Kareem, The graph of a commutative KU -algebra, *Algebra Letters* 2015 (1) (2015) 1–18.
- [21] Samy M. Mostafa, R. A. K. Omar and O. W. Abd El-Baseer, Sub implicative ideals of KU -Algebras, *International Journal of Modern Science and Technology* 2 (5) (2017) 105–116.
- [22] Ali N. A. Koam, Azeem Haider and Moin A. Ansari, Pseudo-metric on KU -algebras, *Korean J. Math.* 27 (1) (2019) 131–140, <https://doi.org/10.11568/kjm.2019.27.1.131>

- [23] Kul Hur, Jong Il Baek, Samy M. Mostafa, Ola W. Abd El-Baseer, Square root fuzzy sub-implicative ideals of KU -algebras, To be published in AFMI.
- [24] M. K. Sen, On Γ -semigroup, Proc. of Inter. Con. of Alg. and its Appl. Decker Publ-icaiton, New York (1981) 301–308.
- [25] M. Murali Krishna Rao, Γ -Group, Bulletin Int. Math. Virtual Inst. 10 (1) (2020) 51–58.
- [26] M. Murali Krishna Rao, Γ -semirings-I, Southeast Asian Bull. of Math. 19 (1) (1995) 49–54.
- [27] J. P. Kaushik and Moin Khan, On bi- Γ -ideal in Γ -Semirings, Int. J. Contemp. Math. Sciences 3 (26) (2008) 1255–1260.
- [28] M. Murali Krishna Rao and B. Venkateswarlu, Regular Γ -incline and field Γ -semiring, Novi Sad J. of Math. 45 (2) (2015) 155–171.
- [29] A. Borumand Saeid, M. Murali Krishna Rao and K. Rajendra Kumar, Γ -BCK-algebras, JMMRC 11 3 (2022) 133–145.
- [30] D. L. Shi, J. I. Baek, S. H. Han, M. Cheong, K. Hur, A study on Γ -BCK-algebras, To be accepted in AFMI.
- [31] D. L. Shi, J. I. Baek, M. Cheong, S. H. Han, Kul Hur, Topological structures on Γ -BCK-algebras, To be submitted.
- [32] N. Bourbaki, General Topology Part 1, Addison-Wesley Publishing Company 1966.
- [33] C. Wayne Patty, Foundations of Topology, PWS Publishing Company 1993. 347–354.

J. I. BAEK (jibaek@wku.ac.kr)

School of Big Data and Financial Statistics, Wonkwang University, Korea

SAMY M. MOSTAFA (samymostafa@yahoo.com)

Department of Mathematics, Faculty of Education, Ain Shams University Roxy, Cairo, Egypt

FATEMA F. KAREEM (fa_sa20072000@yahoo.com)

Department of Mathematics, Ibn-Al-Haitham College of Education University of Baghdad, Iraq

S. H. HAN (shhan235@wku.ac.kr)

Department of Applied Mathematics, Wonkwang University, Korea

K. HUR (kulhur@wku.ac.kr)

Department of Applied Mathematics, Wonkwang University, Korea