

## $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy structures of a $BCC$ -ideal of $BCC$ -algebras and their correlation coefficient

SAMY M. MOSTAFA, AMANY M. MENSRAWY, OLA WAGEEH ABD EL-BASEER,  
D. L. SHI, K. HUR

Received 5 September 2023; Revised 2 December 2023; Accepted 14 December 2023

**ABSTRACT.** A picture fuzzy set is one of the generalizations of Atanassove's (IFSs) fuzzy set. Under this environment, in this manuscript, we familiarize a new type of extensions of fuzzy sets called cubic root fuzzy sets (briefly,  $\sqrt[3]{\cdot}$ -Fuzzy sets) and Fermatean fuzzy sets to contrast  $(3, 3, \sqrt[3]{\cdot})$ -picture sets. We introduce the notion of  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy  $BCC$ -ideals of  $BCC$ -algebras. After then, we study the homomorphic image and inverse image of  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy  $BCC$ -ideals under homomorphism of  $BCC$ -algebras. Moreover, the Cartesian product of  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy  $BCC$ -ideals of  $BCC$ -algebras is given. Finally, we introduce the concept of correlation for  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy sets, which is a new extension of the correlation of Atanassove's IFSs and investigated several properties.

2020 AMS Classification: 06F35, 03G25, 08A72

**Keywords:**  $BCC$ -algebra, Fuzzy  $BCC$ -ideal,  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy  $BCC$ -ideal, The pre-image of a  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy  $BCC$ -ideal under a homomorphism of  $BCC$ -algebras, The Cartesian product of  $(3, 3, \sqrt[3]{\cdot})$ -picture  $BCC$ -ideal, Correlation coefficient for  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy sets.

**Corresponding Author:** Ola W. Abd El-Baseer ([olawagih@yahoo.com](mailto:olawagih@yahoo.com))  
D. L. Shi ([shidali2589@wku.ac.kr](mailto:shidali2589@wku.ac.kr))

### 1. INTRODUCTION

In 1978, Iséki and Tanaka [1] introduced the notion of  $BCK$ -algebras. Iséki [2] introduced the concept of a  $BCI$ -algebra which is a generalization of  $BCK$ -algebra. Since then, numerous mathematical papers have been written investigating the algebraic properties of  $BCK/BCI$ -algebras and their relationship with other structures including lattices and Boolean algebras. There is a great deal of literature which has been produced on the theory of  $BCK/BCI$ -algebras. In particular, the emphasis

seems to have been put on the ideal theory of  $BCK/BCI$ -algebras. Iséki posed an interesting problem (solved by Wron'ski [3]) whether the class of  $BCK$ -algebras is a variety. In connection with this problem, Komori [4] introduced a notion of  $BCC$ -algebras, and Dudek [5] redefined the concept of  $BCC$ -algebras by using a dual form of the ordinary definition in the sense of Komori. In [6], Dudek and Zhang introduced the notion of  $BCC$ -ideals of  $BCC$ -algebras and described connections between such ideals and congruences. In 1956, Zadeh [7] introduced the concept of fuzzy sets. At present, this concept has been applied to many mathematical branches. The idea of "intuitionistic fuzzy set" was first published by Atanassov [8] as a generalization of the notion of fuzzy sets. In 1991, Xi [9] applied fuzzy sets to  $BCI$ ,  $BCK$ ,  $MV$ -algebras. Dudek and Jun [10] considered the fuzzification of  $BCC$ -ideals of  $BCC$ -algebras. In 2013, including the measure of neutral membership and generalizing the notion of intuitionistic fuzzy set, the concept of picture fuzzy set was initiated by Cuong [11] (See [12, 13] for further research). After the initiation of the picture fuzzy set, different types of research works in the context of picture fuzzy set. Yager [14, 15] put forward the concept of Pythagorean fuzzy sets which is a generalization of intuitionistic fuzzy sets, and it is a more powerful tool to solve uncertain problems. Ibrahim et al. [16] defined a new generalization of Pythagorean fuzzy sets called  $(3; 2)$ -fuzzy sets. The main advantage of  $(3; 2)$ -fuzzy sets is that they can characterize more vague cases than Pythagorean fuzzy sets, which can be exploited in many decision-making problems. In 2019, Senapati et al. [17] initiated Fermatean fuzzy sets. Fermatean fuzzy set is characterized by membership and non membership grade restricted that cube sum of its membership grade and non membership grade is less than or equal to one. Correlation coefficient is one of the hot research topics in IFS theory, and it has received a lot of attention from various researchers (See [18, 19, 20, 21, 22, 23, 24]). It is widely used in statistical analysis and engineering sciences. For instance, Gerstenkorn and Manko [18] introduced the correlation coefficients of Atanassov's IFSs. In this paper, we introduce the notion of  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy  $BCC$ -ideals of  $BCC$ -algebras and we study the homomorphic image and inverse image of  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy  $BCC$ -ideals under a homomorphism of  $BCC$ -algebras. Moreover, the Cartesian product of  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy  $BCC$ -ideal of  $BCC$ -algebras is given. Finally, we introduce the concept of correlation for  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy sets, which is a new extension of the correlation of Atanassov's IFSs and investigated several properties.

## 2. PRELIMINARIES

Now we review some definitions and properties that will be useful in our results.

**Definition 2.1** (See [5, 25, 26]). An algebra system  $(X, *, 0)$  of type  $(2, 0)$  is called a  $BCC$ -algebra, if it satisfies the following axioms: for any  $x, y, z \in X$ ,

- (BCC<sub>1</sub>)  $[(x * y) * (z * y)] * (x * z) = 0$ ,
- (BCC<sub>2</sub>)  $x * 0 = x$ ,
- (BCC<sub>3</sub>)  $x * x = 0$ ,
- (BCC<sub>4</sub>)  $0 * x = 0$ ,
- (BCC<sub>5</sub>)  $x * y = 0$  and  $y * x = 0$  imply  $x = y$ .

On a *BCC*-algebra  $X$ , we can define a partial order  $\leq$  on  $X$  by: for any  $x, y \in X$ ,

$$x \leq y \Leftrightarrow x * y = 0.$$

Any *BCK*-algebra is a *BCC*-algebra, but there are *BCC*-algebras which are not *BCK*-algebras (See Dudek [25]). Note that a *BCC*-algebra is a *BCK*-algebra if and only if

$$(x * y) * z = (x * z) * y \text{ for any } x, y, z \in X.$$

In what follows,  $X$  will denote a *BCC*-algebra unless otherwise specified.

**Result 2.2** (See [25]). *The followings hold: for any  $x, y, z \in X$ ,*

- (1)  $(x * y) * x = 0$ ,
- (2)  $x \leq y$  implies  $x * z \leq y * z$ ,
- (3)  $x \leq y$  implies  $z * y \leq z * x$ ,
- (4)  $(x * y) * (z * y) \leq x * z$ .

For any  $x, y \in X$ , we denote  $x \wedge y = y * (y * x)$ .

We will refer to is a *KU*-algebra unless otherwise indicated.

**Definition 2.3** ([25]). Let  $I$  be a nonempty subset of  $X$ . Then  $I$  is called an *BCK-ideal* of  $X$ , if it satisfies the following axioms: for any  $x, y \in X$ ,

- (BCKI<sub>1</sub>)  $0 \in I$ ,
- (BCKI<sub>2</sub>)  $x * y, y \in I$  imply  $x \in I$ .

**Definition 2.4** ([25]). Let  $I$  be a nonempty subset of  $X$ . Then  $I$  is called an *BCC-ideal* of  $X$ , if it satisfies the following axioms: for any  $x, y, z \in X$ ,

- (BCCI<sub>1</sub>)  $0 \in I$ ,
- (BCCI<sub>2</sub>)  $(x * y) * z, y \in I$  imply  $x * z \in I$ .

**Example 2.5** (See [5]). Let  $X = \{0, 1, 2, 3\}$  be a in which the operation  $*$  is defined as follows:

$*$	0	1	2	3
0	0	0	0	0
1	1	0	1	0
2	2	2	0	0
3	3	3	1	0

Table 2.1

Then we can easily check that  $(X, *, 0)$  is a *BCC*-algebra.

**Definition 2.6** (See[8]). Let  $X$  be a nonempty set of  $X$ . Then a mapping  $A = (A^\in, A^\notin)X \rightarrow [0, 1] \times [0, 1]$  is called an *intuitionistic fuzzy set* (briefly, *IFS*) in  $X$ , if  $0 \leq A^\in(x) + A^\notin(x)$  for each  $x \in X$ . In this case,  $A^\in(x)$  [resp.  $A^\notin(x)$ ] is called the *degree of membership* [resp. *nonmembership*] of  $A$  at  $x \in X$ .

**Definition 2.7** ([11, 12]). Let  $X$  be a nonempty universe set. Then a mapping  $A = \langle A_p, A_I, A_N \rangle : X \rightarrow [0, 1] \times [0, 1] \times [0, 1]$  is called a *picture fuzzy set* (briefly, *PFS*) in  $X$ , if  $0 \leq A_p(x) + A_I(x) + A_N(x) \leq 1$  for each  $x \in X$ . In this case,  $A_p(x)$  [resp.  $A_I(x)$  and  $A_N(x)$ ] is called the *positive-membership* [resp. the *neutral-membership* and the *negativ-emembership*] degree of  $A$  at  $x \in X$ .

**Definition 2.8.** Let  $X$  be a nonempty set and let  $A = (A^\in, A^\notin) : X \rightarrow [0, 1] \times [0, 1]$  be a mapping. Then  $A$  is called:

- (i) a Pythagorean fuzzy set in  $X$  (See [14]), if  $0 \leq A^{\in 2}(x) + A^{\notin 2}(x) \leq 1$  for each  $x \in X$ ,
- (ii) an  $n$ -Pythagorean fuzzy set in  $X$  (See [27]), if  $0 \leq A^{\in n}(x) + A^{\notin n}(x) \leq 1$  for each  $x \in X$ ,
- (iii) a Fermatean fuzzy set in  $X$  (See [17]), if  $0 \leq A^{\in 3}(x) + A^{\notin 3}(x) \leq 1$  for each  $x \in X$ ,
- (iv) a  $(3; 2)$ -fuzzy set in  $X$  (See [16, 28, 29]), if  $0 \leq A^{\in 3}(x) + A^{\notin 2}(x) \leq 1$  for each  $x \in X$ ,
- (v) an  $(n; m)$ -fuzzy set in  $X$  (See [29]), if  $0 \leq A^{\in n}(x) + A^{\notin m}(x) \leq 1$  for each  $x \in X$ ,
- (vi) an  $SR$ -fuzzy set in  $X$  (See [30]), if  $0 \leq A^{\in 2}(x) + \sqrt{A^{\notin}(x)} \leq 1$  for each  $x \in X$ ,
- (vii) a  $CR$ -fuzzy set in  $X$  (See [31]), if  $0 \leq A^{\in 3}(x) + \sqrt[3]{A^{\notin}(x)} \leq 1$  for each  $x \in X$ .

### 3. $(3, 3, \sqrt[3]{\cdot})$ -PICTURE FUZZY $BCC$ -IDEALS OF $BCC$ -ALGEBRAS

First of all, we introduce a new notion called  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy set in a nonempty set  $X$  and study some of its properties. Next, we propose the concepts of subalgebras and ideals of  $BCC$ -algebras and deal with some of their properties.

**Definition 3.1.** Let  $X$  be a nonempty set. Then a mapping  $\bar{A} = \langle A_P, A_I, A_N \rangle : X \rightarrow [0, 1] \times [0, 1] \times [0, 1]$  is called a  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy set in  $X$  (briefly,  $(3, 3, \sqrt[3]{\cdot})$ -PFS), if  $0 \leq A_P^3(x) + A_I^3(x) + \sqrt[3]{A_N(x)} \leq 1$  for each  $x \in X$ . The whole [resp. empty]  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy set in  $X$ , denoted by  $\bar{1}$  [resp.  $\bar{0}$ ], is a  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy set in  $X$  defined as follows: for each  $x \in X$ ,

$$\bar{1}(x) = \langle 1, 0, 0 \rangle \text{ [resp. } \bar{0}(x) = \langle 0, 0, 1 \rangle].$$

We will denote the set of all  $(3, 3, \sqrt[3]{\cdot})$ -PFSs in  $X$  as  $PFS_{(3,3,\sqrt[3]{\cdot})}(X)$ .

**Example 3.2.** Let  $X = \{0, a, b, c, d\}$  be a reference set in which the picture fuzzy set  $A = \langle A_P, A_I, A_N \rangle$  is given by the following table:

$X$	0	$a$	$b$	$c$	$d$
$A_P(x)$	0.4	0.3	0.3	0.2	0.1
$A_I(x)$	0.1	0.1	0.2	0.2	0.3
$A_N(x)$	0.006	0.004	0.002	0.001	0.001

Table 3.1

Then we obtain the  $(3, 3, \sqrt[3]{\cdot})$ -PFS  $\bar{A}$  in  $X$  given by the following table:

$X$	0	$a$	$b$	$c$	$d$
$A_P^3(x)$	0.064	0.027	0.027	0.008	0.001
$A_I^3(x)$	0.001	0.001	0.008	0.008	0.027
$\sqrt[3]{A_N(x)}$	0.182	0.159	0.126	0.1	0.1

Table 3.2

**Definition 3.3.** Let  $X$  be a nonempty set and let  $\bar{A}, \bar{B} \in PFS_{(3,3, \sqrt[3]{\cdot})}(X)$ . Then the inclusion  $\bar{A} \subset \bar{B}$  and the equality  $\bar{A} = \bar{B}$  as follows:

- (i)  $\bar{A} \subset \bar{B}$  iff  $A_P^3(x) \leq B_P^3(x)$ ,  $A_I^3(x) \geq B_I^3(x)$ ,  $\sqrt[3]{A_N(x)} \leq \sqrt[3]{B_N(x)}$  for each  $x \in X$ ,
- (ii)  $\bar{A} = \bar{B}$  iff  $\bar{A} \subset \bar{B}$ ,  $\bar{B} \subset \bar{A}$ .

**Definition 3.4.** Let  $X$  be a nonempty set, let  $\bar{A}, \bar{B} \in PFS_{(3,3, \sqrt[3]{\cdot})}(X)$  and let  $(\bar{A}_j)_{j \in J} \subset PFS_{(3,3, \sqrt[3]{\cdot})}(X)$ , where  $J$  denotes an index set. Then the complement  $\bar{A}^c = \langle A_P^c, A_I^c, A_N^c \rangle$  of  $\bar{A}$ , the intersection  $\bar{A} \cap \bar{B}$  and the union  $\bar{A} \cup \bar{B}$  of  $\bar{A}$  and  $\bar{B}$ , the intersection  $\bigcap_{j \in J} \bar{A}_j$  and the union  $\bigcup_{j \in J} \bar{A}_j$  of  $(\bar{A}_j)_{j \in J}$  are  $(3, 3, \sqrt[3]{\cdot})$ -PFSs in  $X$  defined respectively as follows: for each  $x \in X$ ,

- (i)  $\bar{A}^c(x) = \langle 1 - A_N(x), 1 - A_I(x), 1 - A_P(x) \rangle$ ,
- (ii)  $(\bar{A} \cap \bar{B})(x) = \langle (A_P \cap B_P)(x), (A_I \cup B_I)(x), (A_N \cap B_N)(x) \rangle$ , where  $(A_P \cap B_P)^3(x) = A_P^3(x) \wedge B_P^3(x)$ ,  $(A_I \cup B_I)^3(x) = A_I^3(x) \vee B_I^3(x)$ ,  $\sqrt[3]{(A_N \cap B_N)(x)} = \sqrt[3]{A_N(x)} \wedge \sqrt[3]{B_N(x)}$ ,
- (iii)  $(\bar{A} \cup \bar{B})(x) = \langle (A_P \cup B_P)(x), (A_I \cap B_I)(x), (A_N \cup B_N)(x) \rangle$ , where  $(A_P \cup B_P)^3(x) = A_P^3(x) \vee B_P^3(x)$ ,  $(A_I \cap B_I)^3(x) = A_I^3(x) \wedge B_I^3(x)$ ,  $\sqrt[3]{(A_N \cup B_N)(x)} = \sqrt[3]{A_N(x)} \vee \sqrt[3]{B_N(x)}$ ,
- (iv)  $(\bigcap_{j \in J} \bar{A}_j)(x) = \langle (\bigcap_{j \in J} A_{P,j})(x), (\bigcup_{j \in J} A_{I,j})(x), (\bigcap_{j \in J} A_{N,j})(x) \rangle$ , where  $(\bigcap_{j \in J} A_{P,j})^3(x) = \bigwedge_{j \in J} A_{P,j}^3(x)$ ,  $(\bigcup_{j \in J} A_{I,j})^3(x) = \bigvee_{j \in J} A_{I,j}^3(x)$ ,  $\sqrt[3]{(\bigcap_{j \in J} A_{N,j})(x)} = \bigwedge_{j \in J} \sqrt[3]{A_{N,j}(x)}$ ,
- (v)  $(\bigcup_{j \in J} \bar{A}_j)(x) = \langle (\bigcup_{j \in J} A_{P,j})(x), (\bigcap_{j \in J} A_{I,j})(x), (\bigcup_{j \in J} A_{N,j})(x) \rangle$ , where  $(\bigcup_{j \in J} A_{P,j})^3(x) = \bigvee_{j \in J} A_{P,j}^3(x)$ ,  $(\bigcap_{j \in J} A_{I,j})^3(x) = \bigwedge_{j \in J} A_{I,j}^3(x)$ ,  $\sqrt[3]{(\bigcup_{j \in J} A_{N,j})(x)} = \bigvee_{j \in J} \sqrt[3]{A_{N,j}(x)}$ .

**Proposition 3.5.** Let  $X$  be a nonempty set, let  $\bar{A} \in PFS_{(3,3, \sqrt[3]{\cdot})}(X)$  and let  $C(\bar{A}) = \langle C(A_P), C(A_I), C(A_N) \rangle$  be defined by: for each  $x \in X$ ,

$$C(A_P)(x) = A_N(x), \quad C(A_N)(x) = A_P(x),$$

$$C(A_I)(x) = \sqrt[3]{1 - \left( \sqrt[3]{A_P(x)} + A_N^3(x) + \sqrt[3]{A_N(x)} \right)}.$$

Then  $C(\bar{A}) \in PFS_{(3,3, \sqrt[3]{\cdot})}(X)$ .

*Proof.* Let  $x \in X$ . Then we have

$$\begin{aligned} & C(A_P)^3(x) + C(A_I)^3(x) + \sqrt[3]{C(A_N)(x)} \\ &= A_N^3(x) + \left( 1 - \left( \sqrt[3]{A_P(x)} + A_N^3(x) + \sqrt[3]{A_N(x)} \right) \right) + \sqrt[3]{A_P(x)} \\ &= 1 - \sqrt[3]{A_N(x)}. \end{aligned}$$

Since  $0 \leq 1 - \sqrt[3]{A_N(x)} \leq 1$ ,  $0 \leq C(A_P)^3(x) + C(A_I)^3(x) + \sqrt[3]{C(A_N)(x)} \leq 1$ . Thus  $C(\bar{A}) \in PFS_{(3,3, \sqrt[3]{\cdot})}(X)$ .  $\square$

The following is an immediate consequence of Definition 3.4.

**Lemma 3.6.** Let  $X$  be a nonempty sets and let  $\bar{A}, \bar{B}, \bar{C} \in PFS_{(3,3, \sqrt[3]{\cdot})}(X)$ . Then

- (1)  $\bar{A} \cap \bar{A} = \bar{A}, \bar{A} \cup \bar{A} = \bar{A},$
- (2)  $\bar{A} \cap \bar{B} = \bar{B} \cap \bar{A}, \bar{A} \cup \bar{B} = \bar{B} \cup \bar{A},$
- (3)  $(\bar{A} \cap \bar{B}) \cap \bar{C} = \bar{A} \cap (\bar{B} \cap \bar{C}), (\bar{A} \cup \bar{B}) \cup \bar{C} = \bar{A} \cup (\bar{B} \cup \bar{C}),$
- (4)  $\bar{A} \cap (\bar{B} \cup \bar{C}) = (\bar{A} \cap \bar{B}) \cup (\bar{A} \cap \bar{C}), \bar{A} \cup (\bar{B} \cap \bar{C}) = (\bar{A} \cup \bar{B}) \cap (\bar{A} \cup \bar{C}),$
- (5)  $(\bar{A} \cap \bar{B})^c = \bar{A}^c \cup \bar{B}^c, (\bar{A} \cup \bar{B})^c = \bar{A}^c \cap \bar{B}^c,$
- (6)  $(\bar{A}^c)^c = \bar{A},$
- (7)  $\bar{A} \cap \bar{A}^c \neq \bar{0}, \bar{A} \cup \bar{A}^c \neq \bar{1}$  in general (See Example 3.7).

**Example 3.7.** Consider  $\bar{A} = \langle A_P, A_I, A_N \rangle = \langle 0.04, 0.2, 0.006 \rangle \in PFS_{(3,3, \sqrt[3]{\cdot})}(X)$ .

Then clearly,  $\bar{A}^c = \langle 0.994, 0.8, 0.96 \rangle$ . Thus we have: for each  $x \in X$ ,

$$(A_P \cup A_P^c)^3(x) = 0.04^3 \vee 0.994^3 = 0.997 \neq 1,$$

$$(A_P \cap A_P^c)^3(x) = 0.04^3 \wedge 0.994^3 = 0.000064 \neq 0.$$

So  $A_P \cup A_P^c \neq \bar{1}, A_P \cap A_P^c \neq \bar{0}$ .

**Definition 3.8.** Let  $\bar{A}$  be a  $(3, 3, \sqrt[3]{\cdot})$ -PFS in a  $BCC$ -algebra  $X$ . Then  $\bar{A}$  is called a  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy subalgebra (briefly,  $(3, 3, \sqrt[3]{\cdot})$ -PFSA) of  $X$ , if it satisfies the following conditions: for any  $x, y \in X$ ,

$$(PFSA_1) A_P^3(x * y) \geq A_P^3(x) \wedge A_P^3(y),$$

$$(PFSA_2) A_I^3(x * y) \leq A_I^3(x) \vee A_I^3(y),$$

$$(PFSA_2) \sqrt[3]{A_N(x * y)} \geq \sqrt[3]{A_N(x)} \wedge \sqrt[3]{A_N(y)}.$$

**Example 3.9.** Let  $X = \{0, 1, 2, 3\}$  be a set with the binary operation  $*$  defined by the following table:

$*$	0	1	2	3
0	0	0	0	0
1	1	0	1	0
2	2	2	0	0
3	3	3	1	0

Table 3.3

Then we can easily check that  $(X, *, 0)$  is a  $BCC$ -algebra. Define  $A_P, A_I, A_N$  as the following table:

$X$	0	1	2	3
$A_P(x)$	0.04	0.03	0.02	0.01
$A_I(x)$	0.02	0.02	0.02	0.01
$A_N(x)$	0.006	0.004	0.002	0.001

Table 3.4

Then we obtain the following table:

Thus we can easily see that  $\bar{A} = \langle A_P, A_I, A_N \rangle$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFSA of  $X$ .

X	0	1	2	3
$A_P^3(x)$	0.064	0.027	0.008	0.001
$A_I^3(x)$	0.008	0.008	0.027	0.027
$\sqrt[3]{A_N(x)}$	0.1817	0.1587	0.1260	0.1

Table 3.5

**Lemma 3.10.** If  $\bar{A} = \langle A_P, A_I, A_N \rangle$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFSA of a BCC-algebra  $X$ , then the following inequalities hold: for each  $x \in X$ ,

$$A_P^3(0) \geq A_P^3(x), A_I^3(0) \leq A_I^3(x), \sqrt[3]{A_N(0)} \geq \sqrt[3]{A_N(x)}.$$

*Proof.* Suppose  $\bar{A}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFSA of  $X$  and let  $x \in X$ . Then we have

$$\begin{aligned} A_P^3(0) &= A_P^3(x * x) \text{ [By the axiom (BCC}_3\text{)]} \\ &\geq A_P^3(x) \wedge A_P^3(x) \text{ [By the condition (PFSA}_1\text{)]} \\ &= A_P^3(x), \\ A_I^3(0) &= A_I^3(x * x) \\ &\leq A_I^3(x) \vee A_I^3(x) \text{ [By the condition (PFSA}_2\text{)]} \\ &= A_I^3(x), \\ \sqrt[3]{A_N(0)} &= \sqrt[3]{A_N(x * x)} \\ &\geq \sqrt[3]{A_N(x)} \wedge \sqrt[3]{A_N(x)} \text{ [By the condition (PFSA}_3\text{)]} \\ &= \sqrt[3]{A_N(x)}. \end{aligned}$$

Thus the results hold. □

**Definition 3.11.** Let  $\bar{A}$  be a  $(3, 3, \sqrt[3]{\cdot})$ -PFSA of a BCC-algebra  $X$ . Then  $\bar{A}$  is called a  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy BCK-ideal (briefly,  $(3, 3, \sqrt[3]{\cdot})$ -PFBCKI) of  $X$ , if it satisfies the following conditions: for any  $x, y \in X$ ,

$$\begin{aligned} \text{(PFBCKI}_1\text{)} \quad &A_P^3(0) \geq A_P^3(x), A_I^3(0) \leq A_I^3(x), \sqrt[3]{A_N(0)} \geq \sqrt[3]{A_N(x)}, \\ \text{(PFBCCI}_2\text{)} \quad &A_P^3(x) \geq A_P^3(x * y) \wedge A_P^3(y), \\ \text{(PFBCCI}_3\text{)} \quad &A_I^3(x) \leq A_I^3(x * y) \vee A_I^3(y), \\ \text{(PFBCCI}_4\text{)} \quad &\sqrt[3]{A_N(x)} \geq \sqrt[3]{A_N(x * y)} \wedge \sqrt[3]{A_N(y)}. \end{aligned}$$

**Definition 3.12.** Let  $\bar{A}$  be a  $(3, 3, \sqrt[3]{\cdot})$ -PFSA of a BCC-algebra  $X$ . Then  $\bar{A}$  is called a  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy BCC-ideal (briefly,  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI) of  $X$ , if it satisfies the following conditions: for any  $x, y, z \in X$ ,

$$\begin{aligned} \text{(PFBCCI}_1\text{)} \quad &A_P^3(0) \geq A_P^3(x), A_I^3(0) \leq A_I^3(x), \sqrt[3]{A_N(0)} \geq \sqrt[3]{A_N(x)}, \\ \text{(PFBCCI}_2\text{)} \quad &A_P^3(x * z) \geq A_P^3((x * y) * z) \wedge A_P^3(y), \\ \text{(PFBCCI}_3\text{)} \quad &A_I^3(x * z) \leq A_I^3((x * y) * z) \vee A_I^3(y), \\ \text{(PFBCCI}_4\text{)} \quad &\sqrt[3]{A_N(x * z)} \geq \sqrt[3]{A_N((x * y) * z)} \wedge \sqrt[3]{A_N(y)}. \end{aligned}$$

**Example 3.13.** Let  $X$  be the BCC-algebra and  $\bar{A}$  be the  $(3, 3, \sqrt[3]{\cdot})$ -PFS in  $X$  given in Example 3.9. Then we can easily check that  $\bar{A}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X$ .

**Lemma 3.14.** Let  $\bar{A}$  be a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of a BCC-algebra  $X$ . If  $x \leq y$  in  $X$ , then  $A_P^3(x) \geq A_P^3(y)$ ,  $A_I^3(x) \leq A_I^3(y)$ ,  $\sqrt[3]{A_N(x)} \geq \sqrt[3]{A_N(y)}$ .

*Proof.* Let  $x, y \in X$  such that  $x \leq y$ . Then clearly,  $x * y = 0$ . Since  $\bar{A}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X$ , by the condition (PFBCCI<sub>1</sub>),

$$(3.1) \quad A_P^3(0) \geq A_P^3(x), A_I^3(0) \leq A_I^3(x), \sqrt[3]{A_N(0)} \geq \sqrt[3]{A_N(x)}.$$

Thus we have

$$\begin{aligned}
 A_P^3(x) &= A_P^3(x * 0) \text{ [By the axiom (BCC}_2\text{)]} \\
 &\geq A_P^3((x * y) * 0) \wedge A_P^3(y) \text{ [By the condition (PFBCCI}_2\text{)]} \\
 &= A_P^3(0 * 0) \wedge A_P^3(y) \text{ [Since } x * y = 0\text{]} \\
 &= A_P^3(0) \wedge A_P^3(y) \text{ [By the axiom (BCC}_3\text{)]} \\
 &= A_P^3(y), \text{ [By (3.1)]} \\
 A_I^3(x) &= A_I^3(x * 0) \\
 &\leq A_I^3((x * y) * 0) \vee A_I^3(y) \text{ [By the condition (PFBCCI}_3\text{)]} \\
 &= A_I^3(0 * 0) \vee A_I^3(y) \\
 &= A_I^3(0) \vee A_I^3(y) \\
 &= A_I^3(y), \\
 \sqrt[3]{A_N(x)} &= \sqrt[3]{A_N(x * 0)} \\
 &\geq \sqrt[3]{A_N((x * y) * 0)} \wedge \sqrt[3]{A_N(y)} \text{ [By the condition (PFBCCI}_4\text{)]} \\
 &= \sqrt[3]{A_N(0 * 0)} \wedge \sqrt[3]{A_N(y)} \\
 &= \sqrt[3]{A_N(0)} \wedge \sqrt[3]{A_N(y)} \\
 &= \sqrt[3]{A_N(y)}.
 \end{aligned}$$

Thus the results hold.  $\square$

**Lemma 3.15.** *Let  $\bar{A}$  be a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of a BCC-algebra  $X$ . If  $x * y \leq z$  in  $X$ , then the following inequalities:*

$$A_P^3(x) \geq A_P^3(z) \wedge A_P^3(y), \quad A_I^3(x) \leq A_I^3(z) \vee A_I^3(y), \quad \sqrt[3]{A_N(x)} \geq \sqrt[3]{A_N(z)} \wedge \sqrt[3]{A_N(y)}.$$

*Proof.* Let  $x, y, z \in X$  such that  $x * y \leq z$ . Then by Lemma 3.14, we get

$$(3.2) \quad A_P^3(x * y) \geq A_P^3(z), \quad A_I^3(x * y) \leq A_I^3(z), \quad \sqrt[3]{A_N(x * y)} \geq \sqrt[3]{A_N(z)}.$$

Thus we have

$$\begin{aligned}
 A_P^3(x) &= A_P^3(x * 0) \text{ [By the axiom (BCC}_2\text{)]} \\
 &\geq A_P^3((x * y) * 0) \wedge A_P^3(y) \text{ [By the condition (PFBCCI}_2\text{)]} \\
 &= A_P^3(x * y) \wedge A_P^3(y) \text{ [By the axiom (BCC}_2\text{)]} \\
 &\geq A_P^3(z) \wedge A_P^3(y), \text{ [By the axiom (3.2)]} \\
 A_I^3(x) &= A_I^3(x * 0) \\
 &\leq A_I^3((x * y) * 0) \vee A_I^3(y) \text{ [By the condition (PFBCCI}_3\text{)]} \\
 &= A_I^3(x * y) \vee A_I^3(y) \\
 &\geq A_I^3(z) \vee A_I^3(y), \\
 \sqrt[3]{A_N(x)} &= \sqrt[3]{A_N(x * 0)} \\
 &\geq \sqrt[3]{A_N((x * y) * 0)} \wedge \sqrt[3]{A_N(y)} \text{ [By the condition (PFBCCI}_4\text{)]} \\
 &= \sqrt[3]{A_N(x * y)} \wedge \sqrt[3]{A_N(y)} \\
 &\geq \sqrt[3]{A_N(z)} \wedge \sqrt[3]{A_N(y)}.
 \end{aligned}$$

Thus the inequalities hold.  $\square$

**Proposition 3.16.** *Every  $(3, 3, \sqrt[3]{\cdot})$ -PFBCKI of a BCK-algebra  $X$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X$ .*

*Proof.* Let  $\bar{A}$  be a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCKI of a BCK-algebra  $X$ . Then clearly,  $\bar{A}$  satisfies the condition (PFBCCI<sub>1</sub>). Let  $x, y, z \in X$ . Since  $X$  is a BCK-algebra,

$$(x * y) * z = (x * z) * y \text{ for any } x, y, z \in X.$$



Then we have

$$\begin{aligned} A_P^3(x * z) &\geq A_P^3((x * z) * y) \wedge A_P^3(y) \text{ [By the condition (PFBCKI}_2\text{)]} \\ &= A_P^3((x * y) * z) \wedge A_P^3(y), \\ A_I^3(x * z) &\leq A_I^3((x * z) * y) \vee A_I^3(y) \text{ [By the condition (PFBCKI}_3\text{)]} \\ &= A_I^3((x * y) * z) \vee A_I^3(y), \\ \sqrt[3]{A_N(x * z)} &\leq \sqrt[3]{A_N((x * z) * y)} \wedge \sqrt[3]{A_N(y)} \text{ [By the condition (PFBCKI}_4\text{)]} \\ &= \sqrt[3]{A_N((x * y) * z)} \wedge \sqrt[3]{A_N(y)}. \end{aligned}$$

Thus  $\bar{A}$  satisfies the conditions (PFBCCL<sub>2</sub>), (PFBCCL<sub>3</sub>) and (PFBCCL<sub>4</sub>). So  $\bar{A}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X$ .  $\square$

**Proposition 3.17.** *Every  $(3, 3, \sqrt[3]{\cdot})$ -PFBCKI of a BCC-algebra  $X$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFSA of  $X$ .*

*Proof.* Let  $\bar{A}$  be a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCKI of a BCC-algebra  $X$  and let  $x, y \in X$ . Then by Result 2.2 (1) and the condition (PFBCKI<sub>1</sub>), we get

$$(3.3) \quad A_P^3(((x * y) * x) = A_P^3(0) \geq A_P^3(x),$$

$$(3.4) \quad A_I^3(((x * y) * x) = A_I^3(0) \leq A_I^3(x),$$

$$(3.5) \quad \sqrt[3]{A_N((x * y) * x)} = \sqrt[3]{A_N(0)} \geq \sqrt[3]{A_N(x)}.$$

Thus we have

$$\begin{aligned} A_P^3(x * y) &\geq A_P^3((x * y) * x) \wedge A_P^3(x) \text{ [By the condition (PFBCKI}_2\text{)]} \\ &= A_P^3(0) \wedge A_P^3(x) \text{ [By Result 2.2 (1)]} \\ &= A_P^3(x) \text{ [By (3.3)]} \\ &\geq A_P^3(x) \wedge A_P^3(y), \\ A_I^3(x * y) &\leq A_I^3((x * y) * x) \vee A_I^3(x) \text{ [By the condition (PFBCKI}_3\text{)]} \\ &= A_I^3(0) \vee A_I^3(x) \\ &= A_I^3(x) \text{ [By (3.4)]} \\ &\geq A_I^3(x) \vee A_I^3(y), \\ \sqrt[3]{A_N(x * y)} &\geq \sqrt[3]{A_N((x * y) * x)} \wedge \sqrt[3]{A_N(x)} \text{ [By the condition (PFBCKI}_4\text{)]} \\ &= \sqrt[3]{A_N(0)} \wedge \sqrt[3]{A_N(x)} \text{ (1)} \\ &= \sqrt[3]{A_N(x)} \text{ [By (3.4)]} \\ &\geq \sqrt[3]{A_N(x)} \wedge \sqrt[3]{A_N(y)}. \end{aligned}$$

So  $\bar{A}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFSA of  $X$ .  $\square$

**Proposition 3.18.** *Let  $(\bar{M}_j)_{j \in J}$  be a family of  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of a BCC-algebra  $X$ . Then  $\bigcap_{j \in J} \bar{M}_j$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X$ .*

*Proof.* Let  $\bar{M} = \bigcap_{j \in J} \bar{M}_j$  and let  $x \in X$ . Then we have

$$M_P^3(0) = \bigwedge_{j \in J} M_{P,j}^3(0) \geq \bigwedge_{j \in J} M_{P,j}^3(x) = M_P^3(x),$$

$$M_I^3(0) = \bigvee_{j \in J} M_{I,j}^3(0) \leq \bigvee_{j \in J} M_{I,j}^3(x) = M_I^3(x),$$

$$\sqrt[3]{M_N(0)} = \bigwedge_{j \in J} \sqrt[3]{A_{N,j}(0)} \geq \bigwedge_{j \in J} \sqrt[3]{A_{N,j}(x)} = \sqrt[3]{M_N(x)}.$$

Thus  $\overline{M}$  satisfies the condition (PFBCCI<sub>1</sub>).

Now let  $x, y, z \in X$ . Then we get

$$\begin{aligned} M_P^3(x * z) &= \bigwedge_{j \in J} M_{P,j}^3(x * z) \\ &\geq \bigwedge_{j \in J} [M_{P,j}^3((x * y) * z) \wedge M_{P,j}^3(y)] \text{ [By (PFBCCI}_2\text{)]} \\ &= (\bigwedge_{j \in J} [M_{P,j}^3((x * y) * z)]) \wedge (\bigwedge_{j \in J} M_{P,j}^3(y)) \\ &= M_P^3((x * y) * z) \wedge M_P^3(y), \\ M_I^3(x * z) &= \bigvee_{j \in J} M_{I,j}^3(x * z) \\ &\leq \bigvee_{j \in J} [M_{I,j}^3((x * y) * z) \vee M_{I,j}^3(y)] \text{ [By (PFBCCI}_3\text{)]} \\ &= (\bigvee_{j \in J} [M_{I,j}^3((x * y) * z)]) \vee (\bigvee_{j \in J} M_{I,j}^3(y)) \\ &= M_I^3((x * y) * z) \vee M_I^3(y), \\ \sqrt[3]{M_N(x * z)} &= \bigwedge_{j \in J} \sqrt[3]{M_{N,j}(x * z)} \\ &\geq \bigwedge_{j \in J} [\sqrt[3]{M_{N,j}((x * y) * z)} \wedge \sqrt[3]{M_{N,j}(y)}] \text{ [By (PFBCCI}_2\text{)]} \\ &= (\bigwedge_{j \in J} [\sqrt[3]{M_{N,j}((x * y) * z)}]) \wedge (\bigwedge_{j \in J} \sqrt[3]{M_{N,j}(y)}) \\ &= \sqrt[3]{M_N((x * y) * z)} \wedge \sqrt[3]{M_N(y)}. \end{aligned}$$

Thus  $\overline{M}$  satisfies the conditions (PFBCCI<sub>2</sub>), (PFBCCI<sub>3</sub>) and (PFBCCI<sub>4</sub>). So  $\overline{M}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X$ .  $\square$

#### 4. THE IMAGE (PRE-IMAGE) OF A $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI UNDER A HOMOMORPHISM OF $BCC$ -ALGEBRAS

**Definition 4.1.** Let  $(X, *, 0)$  and  $(Y, *', 0')$  be two  $BCC$ -algebras. Then a mapping  $f : X \rightarrow Y$  is called a *homomorphism*, if  $f(x * y) = f(x) *' f(y)$  for any  $x, y \in X$ . Note that if  $f : X \rightarrow Y$  is a homomorphism of  $BCC$ -algebras, then  $f(0) = 0'$ .

**Definition 4.2.** Let  $X$  and  $Y$  be two nonempty set, and let  $\overline{A}$  be a  $(3, 3, \sqrt[3]{\cdot})$ -PFS in  $X$  and let  $\overline{B}$  be a  $(3, 3, \sqrt[3]{\cdot})$ -PFS in  $Y$ .

(i) The *image* of  $\overline{A}$ , denoted by  $f(\overline{A})$ , is a  $(3, 3, \sqrt[3]{\cdot})$ -PFS in  $Y$  defined as follows: for each  $y \in Y$ ,

$$(4.1) \quad f(\overline{A})(y) = \begin{cases} \langle \bigwedge_{x \in f^{-1}(y)} A_P^3(x), \bigvee_{x \in f^{-1}(y)} A_I^3(x), \bigwedge_{x \in f^{-1}(y)} \sqrt[3]{A_N(x)} \rangle & \text{if } f^{-1}(y) \neq \emptyset \\ \langle 0, 1, 0 \rangle & \text{otherwise.} \end{cases}$$

(ii) The *pre-image* of  $\overline{B}$ , denoted by  $f^{-1}(\overline{B}) = \langle f^{-1}(B_P), f^{-1}(B_I), f^{-1}(A_N) \rangle$ , is a  $(3, 3, \sqrt[3]{\cdot})$ -PFS in  $X$  defined as follows: for each  $x \in X$ ,

$$(4.2) \quad f^{-1}(\overline{B})(x) = \langle B_P^3(f(x)), B_I^3(f(x)), \sqrt[3]{B_N(f(x))} \rangle.$$

In fact,  $(f^{-1}(B_P))^3(x) = B_P^3(f(x))$ ,  $(f^{-1}(B_I))^3(x) = B_I^3(f(x))$ ,  $\sqrt[3]{f^{-1}(A_N)(x)} = \sqrt[3]{B_N(f(x))}$ .

**Proposition 4.3.** Let  $f : X \rightarrow Y$  be a homomorphism of  $BCC$ -algebras. If  $\overline{B}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $Y$ , then  $f^{-1}(\overline{B})$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X$ .

*Proof.* Suppose  $\overline{B}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $Y$  and let  $x \in X$ . Then we have

$$\begin{aligned} (f^{-1}(B_P))^3(0) &= B_P^3(f(0)) \\ &= B_P^3(0) \text{ [Since } f \text{ is a homomorphism]} \\ &\geq B_P^3(f(x)) \text{ [By the condition (PFBCCI}_1\text{)]} \\ &= (f^{-1}(B_P))^3(x). \end{aligned}$$

Similarly, we get  $(f^{-1}(B_P))^3(0 \leq (f^{-1}(B_P))^3(x)$  and  $\sqrt[3]{f^{-1}(A_N)(0)} \geq \sqrt[3]{f^{-1}(A_N)(x)}$ . Thus  $f^{-1}(\overline{B})$  satisfies the condition (PFBCCI<sub>1</sub>).

Now let  $x, y, z \in X$ . Then we get

$$\begin{aligned} (f^{-1}(B_P))^3(x * z) &= B_P^3(f(x * z)) \\ &= B_P^3(f(x) * f(z)) \text{ [Since } f \text{ is a homomorphism]} \\ &\geq B_P^3((f(x) * f(y)) * f(z)) \wedge B_P^3(f(y)) \\ &\quad \text{[By the condition (PFBCCI}_2\text{)]} \\ &= B_P^3(f((x * y) * z)) \wedge B_P^3(f(y)) \\ &\quad \text{[Since } f \text{ is a homomorphism]} \\ &= (f^{-1}(B_P))^3((x * y) * z) \wedge (f^{-1}(B_P))^3(z), \\ (f^{-1}(B_I))^3(x * z) &= B_I^3(f(x * z)) \\ &= B_I^3(f(x) * f(z)) \\ &\leq B_I^3((f(x) * f(y)) * f(z)) \vee B_I^3(f(y)) \\ &\quad \text{[By the condition (PFBCCI}_3\text{)]} \\ &= B_I^3(f((x * y) * z)) \vee B_I^3(f(y)) \\ &= (f^{-1}(B_I))^3((x * y) * z) \vee (f^{-1}(B_I))^3(z), \\ \sqrt[3]{f^{-1}(B_N)(x * z)} &= \sqrt[3]{B_N(f(x * z))} \\ &= \sqrt[3]{B_N(f(x) * f(z))} \\ &\geq \sqrt[3]{B_N((f(x) * f(y)) * f(z))} \wedge \sqrt[3]{B_N(f(y))} \\ &\quad \text{[By the condition (PFBCCI}_4\text{)]} \\ &= \sqrt[3]{B_N(f((x * y) * z))} \wedge \sqrt[3]{B_N(f(y))} \\ &= \sqrt[3]{f^{-1}(B_N)((x * y) * z)} \wedge \sqrt[3]{f^{-1}(B_N)(z)}. \end{aligned}$$

Thus  $f^{-1}(\overline{B})$  satisfies the conditions (PFBCCI<sub>2</sub>), (PFBCCI<sub>3</sub>) and (PFBCCI<sub>4</sub>). So  $f^{-1}(\overline{B})$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X$ .  $\square$

**Proposition 4.4.** *Let  $f : X \rightarrow Y$  be an epimorphism of BCC-algebras and let  $\overline{B}$  be a  $(3, 3, \sqrt[3]{\cdot})$ -PFS in  $Y$ . If  $f^{-1}(\overline{B})$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X$ , then  $\overline{B}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $Y$ .*

*Proof.* Suppose  $f^{-1}(\overline{B})$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X$  and let  $a \in Y$ . Since  $f$  is surjective, there is  $x \in X$  such that  $a = f(x)$ . Then we have

$$\begin{aligned} B_P^3(0) &= B_P^3(f(0)) \text{ [Since } f \text{ is a homomorphism]} \\ &= (f^{-1}(B_P))^3(0) \\ &\geq (f^{-1}(B_P))^3(x) \text{ [By the condition (PFBCCI}_1\text{)]} \\ &= B_P^3(f(x)) \\ &= B_P^3(a), \\ B_I^3(0) &= B_I^3(f(0)) \\ &= (f^{-1}(B_I))^3(0) \\ &\leq (f^{-1}(B_I))^3(x) \\ &= B_I^3(f(x)) \\ &= B_I^3(a), \end{aligned}$$

$$\begin{aligned} \sqrt[3]{B_N(0)} &= \sqrt[3]{B_N(f(0))} \\ &= \sqrt[3]{f^{-1}(B_N)(0)} \\ &\geq \sqrt[3]{f^{-1}(B_N)(x)} \\ &= \sqrt[3]{B_N(f(x))} \\ &= \sqrt[3]{B_N(a)}. \end{aligned}$$

Thus  $\bar{B}$  satisfies the condition (PFBCCI<sub>1</sub>).

Now let  $a, b, c \in X$ . Then clearly, there are  $x, y, z \in X$  such that  $a = f(x)$ ,  $b = f(y)$ ,  $c = f(z)$ . Thus we get

$$\begin{aligned} B_P^3(a * c) &= B_P^3((f(x) * f(z))) \\ &= B_P^3(f(x * z)) \text{ [Since } f \text{ is a homomorphism]} \\ &= (f^{-1}(B_P))^3(x * z) \\ &\geq (f^{-1}(B_P))^3((x * y) * z) \wedge (f^{-1}(B_P))^3(y) \\ &\text{ [By the condition (PFBCCI}_2\text{)]} \\ &= B_P^3(f((x * y) * z)) \wedge B_P^3(f(y)) \\ &= B_P^3(((f(x) * f(y)) * f(z)) \wedge B_P^3(f(y)) \\ &\text{ [Since } f \text{ is a homomorphism]} \\ &= B_P^3((a * b) * c) \wedge B_P^3(b), \end{aligned}$$

$$\begin{aligned} B_I^3(a * c) &= B_I^3((f(x) * f(z))) \\ &= B_I^3(f(x * z)) \\ &= (f^{-1}(B_I))^3(x * z) \\ &\leq (f^{-1}(B_I))^3((x * y) * z) \vee (f^{-1}(B_I))^3(y) \\ &\text{ [By the condition (PFBCCI}_3\text{)]} \\ &= B_I^3(f((x * y) * z)) \vee B_I^3(f(y)) \\ &= B_I^3(((f(x) * f(y)) * f(z)) \vee B_I^3(f(y)) \\ &= B_I^3((a * b) * c) \vee B_I^3(b), \end{aligned}$$

$$\begin{aligned} \sqrt[3]{B_N(a * c)} &= \sqrt[3]{B_N((f(x) * f(z)))} \\ &= \sqrt[3]{B_N(f(x * z))} \\ &= \sqrt[3]{f^{-1}(A_N)(x * z)} \\ &\geq \sqrt[3]{f^{-1}(A_N)((x * y) * z)} \wedge \sqrt[3]{f^{-1}(A_N)(y)} \\ &\text{ [By the condition (PFBCCI}_4\text{)]} \\ &= \sqrt[3]{B_N(f((x * y) * z))} \wedge \sqrt[3]{B_N(f(y))} \\ &= \sqrt[3]{B_N(((f(x) * f(y)) * f(z)) \wedge \sqrt[3]{B_N(f(y))} \\ &= \sqrt[3]{B_N((a * b) * c)} \wedge \sqrt[3]{B_N(b)}. \end{aligned}$$

Thus  $\bar{B}$  satisfies the conditions (PFBCCI<sub>2</sub>), (PFBCCI<sub>3</sub>) and (PFBCCI<sub>4</sub>). So  $\bar{B}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $Y$ . □

### 5. THE PRODUCT OF $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI

**Definition 5.1.** Let  $X, Y$  be two nonempty sets, let  $\bar{A}$  be a  $(3, 3, \sqrt[3]{\cdot})$ -PFS in  $X$  and let  $\bar{B}$  be a  $(3, 3, \sqrt[3]{\cdot})$ -PFS in  $Y$ . Then the Cartesian product  $\bar{A} \times \bar{B} = \langle A_P \times B_P, A_I \times B_I, A_N \times B_N \rangle$  of  $\bar{A}$  and  $\bar{B}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFS in  $X \times Y$  defined as follows: for each  $(x, y) \in X \times Y$ ,

$$\begin{aligned} (A_P \times B_P)^3(x, y) &= A_P^3(x) \wedge B_P^3(y), \quad (A_I \times B_I)^3(x, y) = A_I^3(x) \vee B_I^3(y), \\ \sqrt[3]{(A_N \times B_N)(x, y)} &= \sqrt[3]{A_N(x)} \wedge \sqrt[3]{B_N(y)}. \end{aligned}$$

**Remark 5.2.** Let  $X$  and  $Y$  be two  $BCC$ -algebras. We define  $*$  on  $X \times Y$  by: for any  $(x, y), (u, v) \in X \times Y$ ,

$$(x, y) * (u, v) = (x * u, y * v).$$

Then clearly,  $(X \times Y, *, (0, 0))$  is a  $BCC$ -algebra.

**Proposition 5.3.** Let  $X$  be a  $BCC$ -algebra. If  $\bar{A}$  and  $\bar{B}$  are two  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X$ , then  $\bar{A} \times \bar{B}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X \times X$ .

*Proof.* Let  $(x, y) \in X \times X$ . Then by the condition (PFBCCI<sub>1</sub>), we have

$$(A_P \times B_P)^3(0, 0) = A_P(0) \wedge B_P(0) \geq A_P(x) \wedge B_P(y) = (A_P \times B_P)^3(x, y),$$

$$(A_I \times B_I)^3(0, 0) = A_I(0) \vee B_I(0) \leq A_I(x) \vee B_I(y) = (A_I \times B_I)^3(x, y),$$

$$\sqrt[3]{(A_N \times B_N)(0, 0)} = \sqrt[3]{A_N(0) \wedge B_N(0)} \geq \sqrt[3]{A_N(x) \wedge B_N(y)} = \sqrt[3]{(A_N \times B_N)(x, y)}.$$

Thus  $\bar{A} \times \bar{B}$  satisfies the condition (PFBCCI<sub>1</sub>).

Now let  $(x_1, x_2), (y_1, y_2), (z_1, z_2) \in X \times X$ . Then we get

$$\begin{aligned} & (A_P \times B_P)^3(((x_1, x_2) * (y_1, y_2)) * (z_1, z_2)) \wedge (A_P \times B_P)^3(y_1, y_2) \\ &= (A_P \times B_P)^3((x_1 * y_1) * z_1, (x_2 * y_2) * z_2) \wedge (A_P \times B_P)^3(y_1, y_2) \\ &= [A_P((x_1 * y_1) * z_1) \wedge B_P((x_2 * y_2) * z_2)] \wedge [A_P(y_1) \wedge B_P(y_2)] \\ &= [A_P((x_1 * y_1) * z_1) \wedge A_P(y_1)] \wedge [B_P((x_2 * y_2) * z_2) \wedge B_P(y_2)] \\ &\leq A_P(x_1 * z_1) \wedge B_P(x_2 * z_2) \\ &= (A_P \times B_P)^3(x_1 * z_1, x_2 * z_2), \end{aligned}$$

$$\begin{aligned} & (A_I \times B_I)^3(((x_1, x_2) * (y_1, y_2)) * (z_1, z_2)) \vee (A_I \times B_I)^3(y_1, y_2) \\ &= (A_I \times B_I)^3((x_1 * y_1) * z_1, (x_2 * y_2) * z_2) \vee (A_I \times B_I)^3(y_1, y_2) \\ &= [A_I((x_1 * y_1) * z_1) \vee B_I((x_2 * y_2) * z_2)] \vee [A_I(y_1) \vee B_I(y_2)] \\ &= [A_I((x_1 * y_1) * z_1) \vee A_I(y_1)] \vee [B_I((x_2 * y_2) * z_2) \vee B_I(y_2)] \\ &\geq A_I(x_1 * z_1) \vee B_I(x_2 * z_2) \\ &= (A_I \times B_I)^3(x_1 * z_1, x_2 * z_2), \end{aligned}$$

$$\begin{aligned} & \sqrt[3]{(A_N \times B_N)((x_1, x_2) * (y_1, y_2)) * (z_1, z_2)} \wedge \sqrt[3]{(A_N \times B_N)(y_1, y_2)} \\ &= \sqrt[3]{(A_N \times B_N)((x_1 * y_1) * z_1, (x_2 * y_2) * z_2)} \wedge \sqrt[3]{(A_N \times B_N)(y_1, y_2)} \\ &= [\sqrt[3]{A_N((x_1 * y_1) * z_1)} \wedge \sqrt[3]{B_N((x_2 * y_2) * z_2)}] \wedge [\sqrt[3]{A_N(y_1)} \wedge \sqrt[3]{B_N(y_2)}] \\ &= [\sqrt[3]{A_N((x_1 * y_1) * z_1)} \wedge \sqrt[3]{A_N(y_1)}] \wedge [\sqrt[3]{B_N((x_2 * y_2) * z_2)} \wedge \sqrt[3]{B_N(y_2)}] \\ &\leq \sqrt[3]{A_N(x_1 * z_1)} \wedge \sqrt[3]{B_N(x_2 * z_2)} \\ &= \sqrt[3]{(A_N \times B_N)(x_1 * z_1, x_2 * z_2)}. \end{aligned}$$

Thus  $\bar{A} \times \bar{B}$  satisfies the conditions (PFBCCI<sub>2</sub>), (PFBCCI<sub>3</sub>) and (PFBCCI<sub>4</sub>). So  $\bar{A} \times \bar{B}$  is a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI of  $X \times X$ .  $\square$

## 6. CORRELATION COEFFICIENTS FOR $(3, 3, \sqrt[3]{\cdot})$ -PICTURE FUZZY SETS

Correlation plays an important role in statistics, engineering sciences and so on (See [2, 27, 32]). In this section, we propose some correlation coefficients for any two  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy sets.

**Definition 6.1.** Let  $X = \{x_1, x_2, \dots, x_n\}$  be a universe of discourse, and let  $\bar{A}$  and  $\bar{B}$  be two  $(3, 3, \sqrt[3]{\cdot})$ -PFSs.

(i) The *informational*  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy energy of  $\bar{A}$ , denoted by  $E_{(3,3, \sqrt[3]{\cdot})}(A)$ , is defined as follows: for each  $i \in \{1, 2, \dots, n\}$ ,

$$E_{(3,3, \sqrt[3]{\cdot})}(A) = \sum_{i=1}^n \left[ (A_P^3(x_i))^2 + (A_I^3(x_i))^2 + \left( \sqrt[3]{A_N(x_i)} \right)^2 \right].$$

(ii) The *correlation* between  $\bar{A}$  and  $\bar{B}$ , denoted by  $C_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B})$ , is defined as follows: for each  $i \in \{1, 2, \dots, n\}$ ,

$$C_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) = \sum_{i=1}^n (\bar{A} \times \bar{B})(x_i, x_i), \text{ i.e.,}$$

$$C_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) = \sum_{i=1}^n \left[ (A_P^3(x_i) \times B_P^3(x_i)) + (A_I^3(x_i) \times B_I^3(x_i)) + \left( \sqrt[3]{A_N(x_i)} \times \sqrt[3]{B_N(x_i)} \right) \right].$$

(iii) The *correlation coefficient* between  $\bar{A}$  and  $\bar{B}$ , denoted by  $\kappa_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B})$ , is defined as follows: for each  $i \in \{1, 2, \dots, n\}$ ,

$$\kappa_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) = \frac{C_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B})}{\sqrt{E_{(3,3, \sqrt[3]{\cdot})}(A) \cdot E_{(3,3, \sqrt[3]{\cdot})}(B)}}, \text{ i.e.,}$$

$$\kappa_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) = \frac{\sum_{i=1}^n [(A_P^3(x_i) \times B_P^3(x_i)) + (A_I^3(x_i) \times B_I^3(x_i)) + (\sqrt[3]{A_N(x_i)} \times \sqrt[3]{B_N(x_i)})]}{\sqrt{\sum_{i=1}^n [(A_P^3(x_i))^2 + (A_I^3(x_i))^2 + (\sqrt[3]{A_N(x_i)})^2] \cdot \sum_{i=1}^n [(B_P^3(x_i))^2 + (B_I^3(x_i))^2 + (\sqrt[3]{B_N(x_i)})^2]}}.$$

On the other hand, by using idea of Xu [33], we can suggest an alternative form of the correlation coefficient  $\kappa(\bar{A}, \bar{B})$  between two  $\bar{A}$  and  $\bar{B}$  as follows: for each  $i \in \{1, 2, \dots, n\}$ ,

$$\kappa(\bar{A}, \bar{B}) = \frac{\sum_{i=1}^n [(A_P^3(x_i) \times B_P^3(x_i)) + (A_I^3(x_i) \times B_I^3(x_i)) + (\sqrt[3]{A_N(x_i)} \times \sqrt[3]{B_N(x_i)})]}{\sqrt{\sum_{i=1}^n [(A_P^3(x_i))^2 + (A_I^3(x_i))^2 + (\sqrt[3]{A_N(x_i)})^2] \vee \sum_{i=1}^n [(B_P^3(x_i))^2 + (B_I^3(x_i))^2 + (\sqrt[3]{B_N(x_i)})^2]}}.$$

The following is an immediate consequence of Definition 6.1.

**Proposition 6.2.** Let  $\bar{A}$  and  $\bar{B}$  be two  $(3, 3, \sqrt[3]{\cdot})$ -PFSs in a universe of discourse  $X = \{x_1, x_2, \dots, x_n\}$ . Then

- (1)  $C_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{A}) = E_{(3,3, \sqrt[3]{\cdot})}(A)$ ,
- (2)  $C_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) = C_{(3,3, \sqrt[3]{\cdot})}(\bar{B}, \bar{A})$ .

**Proposition 6.3.** Let  $\bar{A}$  and  $\bar{B}$  be two  $(3, 3, \sqrt[3]{\cdot})$ -PFSs in a universe of discourse  $X = \{x_1, x_2, \dots, x_n\}$ . Then

- (1)  $\kappa_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) = \kappa_{(3,3, \sqrt[3]{\cdot})}(\bar{B}, \bar{A})$ ,
- (2)  $0 \leq \kappa_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) \leq 1$ ,
- (3) if  $\bar{A} = \bar{B}$ , then  $\kappa_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) = 1$ .

*Proof.* (1) The proof is straightforward.

(2) It is clear that  $0 \leq \kappa_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B})$ . We prove that  $\kappa_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) \leq 1$ . Recall the well-known Cauchy-Schwarz inequality.

$$(6.1) \quad \left( \sum_{i=1}^n a_i b_i \right)^2 \leq \left( \sum_{i=1}^n a_i^2 \right) \cdot \left( \sum_{i=1}^n b_i^2 \right) \text{ for any } a_i, b_i \in \mathbb{R}.$$

Then by (5.1), we get

$$\begin{aligned} & \left( C_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) \right)^2 \\ &= \left( \sum_{i=1}^n \left[ (A_P^3(x_i) \times B_P^3(x_i)) + (A_I^3(x_i) \times B_I^3(x_i)) + \left( \sqrt[3]{A_N(x_i)} \times \sqrt[3]{B_N(x_i)} \right) \right] \right)^2 \\ &\leq \left( \sum_{i=1}^n \left[ (A_P^3(x_i))^2 + (A_I^3(x_i))^2 + \left( \sqrt[3]{A_N(x_i)} \right)^2 \right] \right) \\ &\quad \cdot \left( \sum_{i=1}^n \left[ (A_P^3(x_i))^2 + (A_I^3(x_i))^2 + \left( \sqrt[3]{A_N(x_i)} \right)^2 \right] \right) \\ &= E_{(3,3, \sqrt[3]{\cdot})}(A) \cdot E_{(3,3, \sqrt[3]{\cdot})}(B). \end{aligned}$$

Thus  $\kappa_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) \leq 1$ . So  $0 \leq \kappa_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) \leq 1$ .

(3) The proof is straightforward. □

**Example 6.4.** Let  $\bar{A}$  and  $\bar{B}$  be two  $(3, 3, \sqrt[3]{\cdot})$ -PFSs in the universe of discourse  $X = \{0, 1, 2, 3\}$  defined by the following tables:

$X$	0	1	2	3
$A_P(x)$	0.04	0.03	0.02	0.01
$A_I(x)$	0.02	0.02	0.03	0.03
$A_N(x)$	0.006	0.004	0.002	0.001
$A_P^3(x)$	0.000064	0.000027	0.000008	0.000001
$A_I^3(x)$	0.000008	0.000008	0.000027	0.000027
$\sqrt[3]{A_N(x)}$	0.181712	0.15874	0.12599	0.1

Table 6.1

$X$	0	1	2	3
$B_P(x)$	0.04	0.03	0.02	0.01
$B_I(x)$	0.01	0.02	0.03	0.04
$B_N(x)$	0.003	0.002	0.002	0.001
$B_P^3(x)$	0.000064	0.000027	0.000008	0.000001
$B_I^3(x)$	0.000001	0.000008	0.000027	0.000064
$\sqrt[3]{B_N(x)}$	0.144225	0.125992	0.125992	0.1

Table 6.2

Then we have

$$E_{(3,3, \sqrt[3]{\cdot})}(A) = 0.496409, \quad E_{(3,3, \sqrt[3]{\cdot})}(B) = 0.566444.$$

Thus  $C_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) = 0.0711$ . So we have

$$(6.2) \quad \kappa_{(3,3, \sqrt[3]{\cdot})}(\bar{A}, \bar{B}) = 0.053306,$$

$$(6.3) \quad \kappa(\bar{A}, \bar{B}) = 0.09451.$$

Hence from (6.2) and (6.3), we see that  $(3, 3, \sqrt[3]{\cdot})$ -PFSs  $\bar{A}$  and  $\bar{B}$  are not correlated.

## 7. CONCLUSIONS

We discussed some properties of  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCIs of  $BCC$ -algebras. The image and the pre-image of  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCIs of  $BCC$ -algebras under a homomorphism of  $BCC$ -algebras were defined and how the image and the pre-image of  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCIs of  $BCC$ -algebras become a  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCI were studied. Moreover, the product of  $(3, 3, \sqrt[3]{\cdot})$ -PFBCCIs was established. Finally, we introduced the concept of correlations for  $(3, 3, \sqrt[3]{\cdot})$ -PFSs, which is a new extension of the correlation of Atanassov's IFSs and investigated its some properties. The main purpose of our future work is to investigate  $(3, 3, \sqrt[3]{\cdot})$ -picture fuzzy topologies and  $(3, 3, \sqrt[3]{\cdot})$ - picture fuzzy  $m$ - polar of  $BCC$ -ideals of  $BCC$ -algebras.

**Acknowledgements.** The authors greatly appreciate the referees for their valuable comments and suggestions for improving the paper.

## REFERENCES

- [1] K. Iséki and S. Tanaka, An introduction to the theory of  $BCK$ -algebras, *Mathematica Japonica* 23 (1) (1978) 1–26.
- [2] K. Iséki, On  $BCI$ -algebras, *Mathematics Seminar Notes* 8 (1) (1980) 125–130.
- [3] A. Wron' ski,  $BCK$ -algebras do not form a variety, *Math. Japon.* 28 (2) (1983) 211–213.
- [4] Y. Komori, The class of  $BCC$ -algebras is not a variety, *Math. Japonica* 29 (1984) 391–394.
- [5] W. A. Dudek, A new characterization of ideals in  $BCC$ -algebras, *Novi Sad J. Math.* 29 (1999) 139–145.
- [6] W. A. Dudek and X. H. Zhang, On ideals and congruences in  $BCC$ -algebras, *Czechoslovak Mathematical Journal* 48 (1) (1998) 21–29.
- [7] L. A. Zadeh, Fuzzy sets, *Information and Control* 8 (1965) 338–353.
- [8] K. Atanassov, Intuitionistic fuzzy sets, *Fuzzy Sets and Systems* 20 (1986) 87–96.
- [9] O. G. Xi, Fuzzy  $BCK$ -algebra, *Math. Japon.* 36 (1991) 935–942.
- [10] W. A. Dudek and Y. B. Jun, Normalizations of fuzzy  $BCC$ -ideals in  $BCC$ -algebras, *Math. Moravica* 3 (1999) 17–24.
- [11] B. C. Cuong, Picture fuzzy sets, *Journal of Computer Science and Cybernetics*, 30 (2014) 409–420.
- [12] B. C. Cuong and V. Kreinovich, Picture fuzzy sets ÔÇô a new concept for computational intelligence problems, *Proceedings of the Third World Congress on Information and Communication Technologies WIICT* (2013).
- [13] B. C. Cuong and P. V. Hai, Some fuzzy logic operators for picture fuzzy sets, *Seventh International Conference on Knowledge and Systems Engineering* (2015).
- [14] R. R. Yager, Pythagorean fuzzy subsets, in *Proceedings of the 2013 joint IFSA world congress and NAFIPS annual meeting (IFSA/NAFIPS)*, IEEE, Edmonton, Canada (2013) 57–61.
- [15] R. R. Yager, Pythagorean membership grades in multi-criteria decision making, *Technical report MII-3301 Machine Intelligence Institute, Iona College, New Rochelle, NY* 2013.
- [16] H. Z. Ibrahim, T. M. Al-shamiand and O. G. Elbarbary,  $(3, 2)$ -Fuzzy sets and their applications to topology and computational optimal choices, *Intelligence and Neuroscience* 2021 (2021), Article ID 1272266, 14 pages.
- [17] T. Senapati and R. R. Yager, Fermatean fuzzy sets, *Journal of Ambient Intelligence and Humanized Computing* 11 (2020) 663–674. (2020)
- [18] T. Gerstenkorn and J. Manko, Correlation of intuitionistic fuzzy sets, *Fuzzy sets Systems* 44 (1991) 39–43.
- [19] H. Bustince and P. Burillo, Correlation of interval-valued intuitionistic fuzzy sets, *Fuzzy Sets and Systems* 74 (1995) 237–244.
- [20] D. H. Hong and S. Y. Hwang, (1995) Correlation of intuitionistic fuzzy sets in probability spaces, *Fuzzy Sets and Systems* 75 (1995) 77–81.



- [21] D. H. Hong,(1998) A note on correlation of interval-valued intuitionistic fuzzy sets, *Fuzzy Sets and Systems* 95 (1998) 113–117.
- [22] W. L. Hung, (2001) Using statistical viewpoint in developing correlation of intuitionistic fuzzy sets. *Int. J. Uncertain Fuzziness Knowl Based Syst.* 9 (2001) 509–516.
- [23] W. L. Hung and J. W. Wu, (2002) Correlation of intuitionistic fuzzy sets by centroid method, *Inform. Sci.* 144 (2002) 219–225.
- [24] H. B. Mitchell (2004) A correlation coefficient for intuitionistic fuzzy sets, *Int. J. Intell. Syst.* 19 (2004) 483–490.
- [25] W. A. Dudek, On proper *BCC*-algebras, *Bull. Inst. Math. Academia Sinica* 20 (1992) 137–150.
- [26] W. A. Dudek and Y. B. Jun, Fuzzy *BCC*-ideals in *BCC*-algebras, *Math. Montisnigri* 10 (1999) 21–30.
- [27] A. Bryniarska, The *n*-Pythagorean fuzzy sets, *Symmetry* 2020, 12,1772.
- [28] S. S. Ahn, H. S. Kim, S. Z. Song and Y. B. Jun, The (2,3)-fuzzy set and its application in *BCK*-algebras and *BCI*-algebras *J. Math. Computer Sci.* 2 (2022) 118–130.
- [29] Y. B. Jun and K. Hur, The (*m*; *n*)-fuzzy set and its application in *BCK*-algebras, *Anna. Fuzzy Math. Inform.* 24 (1) (2022) 17–29.
- [30] T. M. Al-shami, A. I. EL-Maghrabi, H. Z. Ibrahim and A. A. Azzam, *SR*-fuzzy sets and their weighted aggregated operators in application to decision-making *Journal of Function Spaces* 2022 2022), Article ID 3653225, 14 pages.
- [31] Y. A. Salih and Hariwan Z. Ibrahim, *CR*-fuzzy sets and their applications, *J. Math. Computer Sci.* 28 (2023) 171–181.
- [32] D. A. Chiang and N. P. Lin. Correlation of fuzzy sets *Fuzzy Sets Systems* 102 (1999) 221–226.
- [33] Z. S. Xu, On correlation measures of intuitionistic fuzzy sets, *Lect. Notes Comput. Sci.* 4224 (2006) 15–24.

SAMY M. MOSTAFA (samymostafa@yahoo.com)

Department of Mathematics, Faculty of Education, Ain Shams University, Roxy, Cairo, Egypt

AMANY M. MENSRAWY (amaniminshawi@edu.asu.edu.eg)

Department of Mathematics, Faculty of Education, Ain Shams University, Roxy, Cairo, Egypt

OLA W. ABD EL-BASEER (olawagih@yahoo.com)

Department of Mathematics, Faculty of Education, Ain Shams University, Roxy, Cairo, Egypt

D. L. SHI (shidali2589@wku.ac.kr)

School of Big Data and Financial Statistics, Wonkwang University, Korea

KUL HUR (kulhur@wku.ac.kr )

Department of Applied Mathematics, Wonkwang University, Korea