

α - b -regularity in a fuzzy topological space

ANJANA BHATTACHARYYA

Received 1 November 2023; Revised 24 November 2023; Accepted 30 December 2023

ABSTRACT. This paper deals with a new type of fuzzy separation axiom, viz., fuzzy α - b -regular space by introducing fuzzy α - b -open set as a basic tool. This newly defined class of sets is strictly larger than that of fuzzy open set as well as fuzzy preopen set, fuzzy semiopen set, fuzzy α -open set and fuzzy β -open set. Also, we introduce new type of fuzzy compact space and a strong form of fuzzy T_2 -space. However, three different types of functions are introduced and studied. Also the mutual relationships of these functions are established. Lastly some applications of these functions on the spaces introduced here are established.

2020 AMS Classification: 54A40, 03E72

Keywords: Fuzzy α - b -open set, Fuzzy regular open set, Fuzzy α - b - r -continuous function, Fuzzy α - b -continuity, Fuzzy almost α - b -continuity, Fuzzy extremally disconnected space.

Corresponding Author: Anjana Bhattacharyya (anjanabhattacharyya@hotmail.com)

1. INTRODUCTION

Fuzzy α -open set was introduced in [1]. Using this concept as a basic tool, here we introduce fuzzy α - b -open set. After introducing fuzzy continuous function in [2], different types of fuzzy continuous-like functions were introduced and studied. Using the concept of fuzzy regular closed set [3], here we introduce fuzzy α - b - r -continuous function, fuzzy α - b -continuity, and fuzzy almost α - b -continuity. Fuzzy regular space was introduced in [4]. Here we introduce fuzzy α - b -regular space, the class of which is strictly larger than that of fuzzy regular space. It is shown that in this space fuzzy open set and fuzzy α - b -open set coincide. Again fuzzy compact space was introduced by Chang [2]. Here we introduce fuzzy α - b -compactness which is weaker than fuzzy compactness. Also fuzzy α - b - T_2 -space was introduced, the class of which is strictly larger than that of fuzzy T_2 -space [4]. Recently, new types of fuzzy sets, viz., fuzzy soft set and fuzzy octahedron set are

introduced and studied. A new branch of fuzzy topology is developed using these types of fuzzy sets. In this context we have to mention [5, 6, 7, 8, 9, 10].

2. PRELIMINARIES

We recall the concepts related to fuzzy sets introduced by Zadeh [11]. A *fuzzy set* A in a nonempty set X is a mapping from X into the closed interval $I = [0, 1]$, i.e., $A \in I^X$. The *support* of a fuzzy set A , denoted by $\text{supp}A$, is defined by $\text{supp}A = \{x \in X : A(x) \neq 0\}$. The fuzzy set with the singleton support $\{x\} \subseteq X$ and the value t ($0 < t \leq 1$) will be denoted by x_t . 0_X and 1_X are the constant fuzzy sets taking values 0 and 1 respectively in X . The *complement* of a fuzzy set A in X , denoted by $1_X \setminus A$, is defined by $(1_X \setminus A)(x) = 1 - A(x)$ for each $x \in X$. For any two fuzzy sets A, B in X , $A \leq B$ means $A(x) \leq B(x)$ for all $x \in X$, while AqB means there exists $x \in X$ such that $A(x) + B(x) > 1$ and A is said to be *quasi-coincident* (q-coincident, for short) *with* B [12]. The negation of these two statements will be denoted by $A \not\leq B$ and $A \not q B$ respectively.

Throughout the paper, (X, τ) or simply by X we shall mean a fuzzy topological space (fts, for short) in the sense of Chang [2]. For a fuzzy set A in a fts X , clA and $\text{int}A$ stand for the *fuzzy closure* and the *fuzzy interior* of A in X as follows [2]:

$$clA = \bigwedge \{F : A \leq F, F \text{ is a closed set in } X\},$$

$$\text{int}A = \bigvee \{G : G \leq A, G \text{ is an open set in } X\}.$$

For a fts X , $A \in I^X$ is said to be *fuzzy regular open* [3] (resp. *fuzzy semiopen* [3], *fuzzy preopen* [13], *fuzzy α -open* [1], *fuzzy β -open* [14]), if $A = \text{int}(clA)$ (resp. $A \leq cl(\text{int}A)$, $A \leq \text{int}(clA)$, $A \leq \text{int}(cl(\text{int}A))$, $A \leq cl(\text{int}(clA))$). The complement of a fuzzy regular open (resp. fuzzy semiopen, fuzzy preopen, fuzzy α -open, fuzzy β -open) set is called a *fuzzy regular closed* (resp. *fuzzy semiclosed*, *fuzzy preclosed*, *fuzzy α -closed*, *fuzzy β -closed*) *set*. The smallest fuzzy semiclosed (resp., fuzzy preclosed, fuzzy α -closed, fuzzy β -closed) set containing a fuzzy set A in X is called the *fuzzy semiclosure* (resp. *fuzzy preclosure*, *fuzzy α -closure*, *fuzzy β -closure*) of A , denoted by $sclA$ (resp. $pclA$, αclA , βclA). It is well-known that A is fuzzy semiclosed (resp. fuzzy preclosed, fuzzy α -closed, fuzzy β -closed) in a fts X if and only if $A = sclA$ (resp. $A = pclA$, $A = \alpha clA$, $A = \beta clA$). The collection of all fuzzy regular open (resp. fuzzy semiopen, fuzzy preopen, fuzzy α -open, fuzzy β -open) sets in X is denoted by $FRO(X)$ (rssp., $FSO(X)$, $FPO(X)$, $F\alpha O(X)$, $F\beta O(X)$) and the collection of all fuzzy regular closed (resp., fuzzy semiclosed, fuzzy preclosed, fuzzy α -closed, fuzzy β -closed) sets in X is denoted by $FRC(X)$ (rssp., $FSC(X)$, $FPC(X)$, $F\alpha C(X)$, $F\beta C(X)$). For a fuzzy open set A in X , $sclA = \text{int}(clA)$ [15].

3. FUZZY α -b-OPEN SET : SOME PROPERTIES

In this section, we first introduce fuzzy α -b-open set and then establish the mutual relationships of this newly defined set with the sets defined in [1, 3, 13, 14].

First we recall some definitions from [16] for ready references.

Definition 3.1 ([16]). Let (X, τ) be an fts and $A \in I^X$. A fuzzy point x_α in X is said to be a *fuzzy θ -semicluster point* of A , if $clUqA$ for all $U \in FSO(X)$ with $x_\alpha q U$.

The union of all fuzzy θ -semiclosed points of A is called the *fuzzy θ -semiclosure* of A and is denoted by $\theta\text{-scl}A$. It is obvious that A is fuzzy θ -semiclosed in X if and only if $A = \theta\text{-scl}A$.

Definition 3.2 ([16]). Let (X, τ) be an fts and $A \in I^X$. Then the *r-kernel* of A , denoted by $r\text{-Ker}A$, is defined as follows :

$$r\text{-Ker}A = \bigwedge \{U : U \in FRO(X), A \leq U\}.$$

Let us now introduce the following concept.

Definition 3.3. A fuzzy set A in an fts (X, τ) is said to be *fuzzy α -b-open* in X , if $A \leq cl(\alpha int(clA))$. The complement of a fuzzy α -b-open set is said to be *fuzzy α -b-closed* in X . The collection of all fuzzy α -b-open (resp. fuzzy α -b-closed) sets in an fts X is denoted by $FabO(X)$ (resp. $FabC(X)$).

Remark 3.4. The union of any two fuzzy α -b-open sets is also so. But intersection of any two fuzzy α -b-open sets may not be so, as it seen from the following example.

Example 3.5. Let $X = \{a, b\}$ and let $\tau = \{0_X, 1_X, A\}$, where $A(a) = 0.5, A(b) = 0.6$. Then (X, τ) is an fts. Consider two fuzzy sets B, C in X defined by:

$$B(a) = 0.4, B(b) = 0.6, C(a) = 0.6, C(b) = 0.4.$$

Then clearly, $B, C \in FabO(X)$. Let $D = B \wedge C$. Then $D(a) = D(b) = 0.4$. Thus

$$cl(\alpha int(clD)) = cl(\alpha int(1_X \setminus A)) = cl0_X = 0_X \not\geq D.$$

So $D \notin FabO(X)$.

From Example 3.5, we can conclude that the set of all fuzzy α -b-open sets in an fts X does not form a fuzzy topology.

Remark 3.6. It is clear from definitions that fuzzy open set, fuzzy semiopen set, fuzzy preopen set, fuzzy α -open set, fuzzy β -open set implies fuzzy α -b-open set, but the reverse implications are not necessarily true follow from the following example.

Example 3.7. Let $X = \{a, b\}$ and let $\tau = \{0_X, 1_X, A\}$, where $A(a) = 0.5, A(b) = 0.4$. Then (X, τ) is an fts. Here $F\alpha O(X) = \{0_X, 1_X, U\}$, where $A \not\prec U \not\leq 1_X \setminus A$. Consider a fuzzy set B in X defined by $B(a) = B(b) = 0.5$. Then clearly, $B \notin \tau, B \notin FPO(X), B \notin F\alpha O(X)$. But $cl(\alpha int(clB)) = 1_X \setminus A \geq B$. Thus $B \in FabO(X)$.

Next consider the fuzzy set C in X defined by $C(a) = C(b) = 0.4$. Then clearly, $C \notin FSO(X)$, but $C \in FabO(X)$.

As $\tau \subseteq F\alpha O(X)$, clearly fuzzy α -b-open set may not necessarily fuzzy β -open set.

Theorem 3.8. Let (X, τ) be an fts. Then the union of any collection of fuzzy α -b-open sets in X is fuzzy α -b-open in X .

Proof. Let $\mathcal{G} = \{G_\alpha : \alpha \in \Lambda\}$ be any collection of fuzzy α -b-open sets in X and let $\alpha \in \Lambda$. Then clearly, $G_\alpha \leq cl(\alpha int(clG_\alpha))$. Also, $G_\alpha \leq \bigvee_{\alpha \in \Lambda} G_\alpha$. Thus $clG_\alpha \leq cl(\bigvee_{\alpha \in \Lambda} G_\alpha)$. So $G_\alpha \leq cl(\alpha int(clG_\alpha)) \leq cl(\alpha int(cl(\bigvee_{\alpha \in \Lambda} G_\alpha)))$. Taking union on both sides, $\bigvee_{\alpha \in \Lambda} G_\alpha \leq cl(\alpha int(cl(\bigvee_{\alpha \in \Lambda} G_\alpha)))$. Hence $\bigvee_{\alpha \in \Lambda} G_\alpha$ is fuzzy α -b-open in X . \square

Definition 3.9. Let (X, τ) be an fts and $A \in I^X$. Then fuzzy α - b -closure of A , denoted by $abclA$, is defined by

$$abclA = \bigwedge \{U \in I^X : A \leq U, U \in FabC(X)\}$$

and fuzzy α - b -interior of A , denoted by $abintA$, is defined by

$$abintA = \bigvee \{G \in I^X : G \leq A, G \in FabO(X)\}.$$

Note 3.10. By Remark 3.4, we can conclude that for any fuzzy set A in an fts X , $abclA$ is fuzzy α - b -closed and $abintA$ is fuzzy α - b -open. Again, if $A \in FabC(X)$, then $A = abclA$ and if $A \in FabO(X)$, then $A = abintA$.

Result 3.11. Let (X, τ) be an fts. Then the following statements are true:

- (1) for any fuzzy point x_t in X and any $U \in I^X$, $x_t \in abclU$ implies that for any $V \in FabO(X)$ with $x_t qV$, $V qU$,
- (2) for any two fuzzy sets U, V , where $V \in FabO(X)$, if $U \not qV$, then $abclU \not qV$.

Proof. (1) Let $x_t \in abclU$ and $V \in FabO(X)$ with $x_t qV$. Then $x_t \notin 1_X \setminus V \in FabC(X)$. Thus $U \not \leq 1_X \setminus V$. So $U qV$.

(2) Assume that $abclU qV$ but $U \not qV$. Then there exists $x \in X$ such that $(abclU)(x) + V(x) > 1$. Thus $V(x) + t > 1$, where $t = (abclU)(x)$. So $x_t \in abclU$, where $x_t qV$, $V \in FabO(X)$. By (1), $V qU$. This is a contradiction. \square

Result 3.12. Let (X, τ) be an fts and $A \in I^X$. Then the following statements are true:

- (1) $abcl(1_X \setminus A) = 1_X \setminus abintA$,
- (2) $1_X \setminus abclA = abint(1_X \setminus A)$.

Proof. (1) Let $x_t \in abcl(1_X \setminus A)$. Assume that $x_t \notin 1_X \setminus abintA$. Then $x_t q abintA$. Thus there exists $U \in FabO(X)$ with $U \leq A$ such that $x_t qU$. Since $x_t \in abcl(1_X \setminus A)$, by Result 3.11 (1), $U q(1_X \setminus A)$. So $A q(1_X \setminus A)$. This is a contradiction. Hence

$$(3.1) \quad abcl(1_X \setminus A) \leq 1_X \setminus abintA.$$

Conversely, let $x_t \in 1_X \setminus abintA$. Then we have

$$1 - abintA(x) \geq t \Rightarrow x_t \not q abintA \Rightarrow x_t \not q U,$$

$$(3.2) \quad \text{where } U \in FabO(X) \text{ with } U \leq A.$$

Let $V \in FabC(X)$ with $1_X \setminus A \leq V$. Then $1_X \setminus V \leq A$, where $1_X \setminus V \in FabO(X)$. Thus by (3.2), we get

$$x_t \not q(1_X \setminus V) \Rightarrow x_t \in V \Rightarrow x_t \in abcl(1_X \setminus A).$$

So we have

$$(3.3) \quad 1_X \setminus abintA \leq abcl(1_X \setminus A).$$

Hence combining (3.1) and (3.3), we get the result.

- (2) Writing $1_X \setminus A$ for A in (1), we get the proof. \square

Let us now recall the following Lemma from [16] for ready references.

Lemma 3.13 ([16]). *Let (X, τ) be an fts and $A \in I^X$. Then the following statements hold:*

- (1) *for any $A \in FRO(X)$, $\theta\text{-scl}A = A$,*
- (2) *for any $A \in F\beta O(X)$, $clA = \alpha clA$,*
- (3) *for any $A \in FSO(X)$, $clA = pclA$,*
- (4) *for any $A \in \tau$, $sclA = \theta\text{-scl}A$.*

4. FUZZY α - b - r -CONTINUOUS FUNCTION : SOME CHARACTERIZATIONS

In this section, we first introduce fuzzy α - b - r -continuous function and characterize it in several ways. Afterwards, two new types of functions, viz., fuzzy α - b -continuous function and fuzzy almost α - b -continuous function are introduced. The mutual relationships of these three functions are established here.

Definition 4.1. Let (X, τ) and (Y, τ_1) be two fts's. Then $f : X \rightarrow Y$ is called a *fuzzy α - b - r -continuous function*, if $f^{-1}(A) \in FabC(X)$ for all $A \in FRO(Y)$.

Theorem 4.2. *Let (X, τ) and (Y, τ_1) be two fts's and $f : X \rightarrow Y$ be a function. Then the following statements are equivalent:*

- (1) *f is fuzzy α - b - r -continuous,*
- (2) *$f^{-1}(A) \in FabO(X)$ for all $A \in FRC(Y)$,*
- (3) *$f(abcl_\tau U) \leq r\text{-ker}(f(U))$ for all $U \in I^X$,*
- (4) *$abcl_\tau(f^{-1}(A)) \leq f^{-1}(r\text{-ker}(A))$ for all $A \in I^Y$,*
- (5) *$abcl_\tau(f^{-1}(R)) \leq f^{-1}(\theta\text{-scl}_{\tau_1} R)$ for all $R \in \tau_1$,*
- (6) *$abcl_\tau(f^{-1}(R)) \leq f^{-1}(scl_{\tau_1} R)$ for all $R \in \tau_1$,*
- (7) *$abcl_\tau(f^{-1}(R)) \leq f^{-1}(int_{\tau_1}(cl_{\tau_1} R))$ for all $R \in \tau_1$,*
- (8) *$f^{-1}(int_{\tau_1}(cl_{\tau_1} A)) \in FabC(X)$ for all $A \in \tau_1$,*
- (9) *$f^{-1}(cl_{\tau_1}(int_{\tau_1} F)) \in FabO(X)$ for all $F \in \tau_1^c$,*
- (10) *$f^{-1}(cl_{\tau_1} U) \in FabO(X)$ for all $U \in F\beta O(Y)$,*
- (11) *$f^{-1}(cl_{\tau_1} U) \in FabO(X)$ for all $U \in FSO(Y)$,*
- (12) *$f^{-1}(int_{\tau_1}(cl_{\tau_1} U)) \in FabC(X)$ for all $U \in FPO(Y)$,*
- (13) *$f^{-1}(\alpha cl_{\tau_1} U) \in FabO(X)$ for all $U \in F\beta O(Y)$,*
- (14) *$f^{-1}(pcl_{\tau_1} U) \in FabO(X)$ for all $U \in FSO(Y)$.*

Proof. (1) \Leftrightarrow (2): Obvious.

(2) \Rightarrow (3): Let $U \in I^X$ and suppose y_t is a fuzzy point in Y with $y_t \notin r\text{-ker}(f(U))$. Then there exists $V \in FRO(Y)$ such that $f(U) \leq V$ and $y_t \notin V$, i.e., $V(y) < t$. Thus $y_t q(1_Y \setminus V) \in FRC(Y)$ and $1_Y \setminus f(U) \geq 1_Y \setminus V$. So $f(U) \not\leq q(1_Y \setminus V)$, i.e., $U \not\leq f^{-1}(1_Y \setminus V)$. By (2), $f^{-1}(1_Y \setminus V) = 1_X \setminus f^{-1}(V) \in FabO(X)$. By Result 3.11 (2), $abcl_\tau U \not\leq q(1_X \setminus f^{-1}(V))$. This implies that $abcl_\tau U \leq f^{-1}(V)$, i.e., $f(abcl_\tau U) \leq V$ implies that $1_Y \setminus f(abcl_\tau U) \geq 1_Y \setminus V$. Hence we have

$$1 - f(abcl_\tau U)(y) \geq 1 - V(y) > 1 - t, \text{ i.e., } t > f(abcl_\tau U)(y), \text{ i.e., } y_t \notin f(abcl_\tau U).$$

Therefore $f(abcl_\tau U) \leq r\text{-ker}(f(U))$.

(3) \Rightarrow (4): Let $A \in I^Y$. Then $f^{-1}(A) \in I^X$. Thus by (3), we get

$$f(abcl_\tau f^{-1}(A)) \leq r\text{-ker}(f(f^{-1}(A))) \leq r\text{-ker}(A).$$

So $abcl_\tau(f^{-1}(A)) \leq f^{-1}(r\text{-ker}(A))$.

(4) \Rightarrow (1): Let $A \in FRO(Y)$. By (4), $\alpha bcl_\tau(f^{-1}(A)) \leq f^{-1}(r\text{-ker}(A)) = f^{-1}(A)$. But $f^{-1}(A) \leq \alpha bcl_\tau(f^{-1}(A))$. Thus $f^{-1}(A) = \alpha bcl_\tau(f^{-1}(A))$. So $f^{-1}(A) \in FabC(X)$. Hence f is a fuzzy α - b - r -continuous function.

(5) \Leftrightarrow (6): Follows from Lemma 3.13 (4).

(6) \Leftrightarrow (7): Obvious

(7) \Rightarrow (1): Let $A \in FRO(Y)$. Then by (7), we get

$$\alpha bcl_\tau(f^{-1}(A)) \leq f^{-1}(int_{\tau_1}(cl_{\tau_1}A)) = f^{-1}(A).$$

Thus $f^{-1}(A) \in FabC(X)$. So f is a fuzzy α - b - r -continuous function.

(1) \Rightarrow (7): Let $A \in \tau_1$. Then $int_{\tau_1}(cl_{\tau_1}A) \in FRO(Y)$. Then by (1), $f^{-1}(int_{\tau_1}(cl_{\tau_1}A)) \in FabC(X)$. Thus we have

$$\alpha bcl_\tau(f^{-1}(A)) \leq \alpha bcl_\tau(f^{-1}(int_{\tau_1}(cl_{\tau_1}A))) = f^{-1}(int_{\tau_1}(cl_{\tau_1}A)).$$

(1) \Rightarrow (8): Let $A \in \tau_1$. Then $int_{\tau_1}(cl_{\tau_1}A) \in FRO(Y)$. Thus by (1), $f^{-1}(int_{\tau_1}(cl_{\tau_1}A)) \in FabC(X)$.

(8) \Rightarrow (1): Let $A \in FRO(Y)$. Then $A \in \tau_1$. Thus by (8), $f^{-1}(A) = f^{-1}(int_{\tau_1}(cl_{\tau_1}A)) \in FabC(X)$.

(2) \Rightarrow (9): Let $F \in \tau_1^c$. Then $cl_{\tau_1}int_{\tau_1}F \in FRC(Y)$. Thus by (2), $f^{-1}(cl_{\tau_1}(int_{\tau_1}F)) \in FabO(X)$.

(9) \Rightarrow (2): Let $F \in FRC(Y)$. Then by (9), $f^{-1}(F) = f^{-1}(cl_{\tau_1}(int_{\tau_1}F)) \in FabO(X)$.

(2) \Rightarrow (10): Let $U \in F\beta O(Y)$. Then $U \leq cl_{\tau_1}(int_{\tau_1}(cl_{\tau_1}U)) \leq cl_{\tau_1}U$. Thus

$$cl_{\tau_1}U \leq cl_{\tau_1}(cl_{\tau_1}(int_{\tau_1}(cl_{\tau_1}U))) = cl_{\tau_1}(int_{\tau_1}(cl_{\tau_1}U)) \leq cl_{\tau_1}(cl_{\tau_1}U) = cl_{\tau_1}U.$$

So $cl_{\tau_1}U = cl_{\tau_1}(int_{\tau_1}(cl_{\tau_1}U))$. Hence $cl_{\tau_1}U \in FRC(Y)$. Therefore by (2), $f^{-1}(cl_{\tau_1}U) \in FabO(X)$.

(10) \Rightarrow (6): Since $FSO(Y) \subseteq F\beta O(Y)$, by (10), $f^{-1}(cl_{\tau_1}U) \in FabO(X)$ for all $U \in FSO(Y)$.

(11) \Rightarrow (12): Let $U \in FPO(Y)$. Then $U \leq int_{\tau_1}(cl_{\tau_1}U)$. We claim that $int_{\tau_1}(cl_{\tau_1}U) \in FRO(Y)$. Note that the following inequalities hold:

$$int_{\tau_1}(cl_{\tau_1}U) \leq int_{\tau_1}(cl_{\tau_1}(int_{\tau_1}(cl_{\tau_1}U))) \leq int_{\tau_1}(cl_{\tau_1}U).$$

Then $int_{\tau_1}(cl_{\tau_1}U) = int_{\tau_1}(cl_{\tau_1}(int_{\tau_1}(cl_{\tau_1}U)))$. Thus $1_Y \setminus int_{\tau_1}(cl_{\tau_1}U) \in FRC(Y)$. So $1_Y \setminus int_{\tau_1}(cl_{\tau_1}U) \in FSO(Y)$. By (11), $f^{-1}(cl_{\tau_1}(1_Y \setminus int_{\tau_1}(cl_{\tau_1}U))) \in FabO(X)$. Hence we have

$$1_X \setminus f^{-1}(int_{\tau_1}(int_{\tau_1}(cl_{\tau_1}U))) = 1_X \setminus f^{-1}(int_{\tau_1}(cl_{\tau_1}U)) \in FabO(X).$$

Therefore $f^{-1}(int_{\tau_1}(cl_{\tau_1}U)) \in FabC(X)$.

(12) \Rightarrow (1): Let $U \in FRO(Y)$. Then $U \in FPO(Y)$. By (12), $f^{-1}(int_{\tau_1}(cl_{\tau_1}U)) \in FabC(X)$. Thus $f^{-1}(U) = f^{-1}(int_{\tau_1}(cl_{\tau_1}U)) \in FabC(X)$. So (1) holds.

(10) \Leftrightarrow (13): The proof follows from Lemma 3.13 (2).

(11) \Leftrightarrow (14): The proof follow from Lemma 3.13 (3). □

Theorem 4.3. Let (X, τ) and (Y, τ_1) be two fts's and $f : X \rightarrow Y$ be a function. Consider the following statements:

- (a) for each fuzzy point x_t in X and each $A \in FSO(Y)$ with $f(x_t)qA$, there exists $U \in FabO(X)$ with x_tqU , $f(U) \leq cl_{\tau_1}A$,
- (b) $f(\alpha bcl_\tau P) \leq \theta\text{-scl}_{\tau_1}(f(P))$ for all $P \in I^X$,

- (c) for each fuzzy point x_t in X and each $A \in FSO(Y)$ with $f(x_t) \in A$, there exists $U \in F\alpha bO(X)$ such that $x_t \in U$ and $f(U) \leq cl_{\tau_1} A$,
- (d) $f^{-1}(A) \leq \alpha bint_{\tau}(f^{-1}(cl_{\tau_1} A))$ for all $A \in FSO(Y)$,
- (e) $\alpha bcl_{\tau}(f^{-1}(R)) \leq f^{-1}(\theta-scl_{\tau_1} R)$ for all $R \in I^Y$,
- (f) f is a fuzzy α - b - r -continuous function.

Then we have:

- (1) (a), (b), (c), (d) and (e) are equivalent,
- (2) (e) implies (f).

Proof. (1) (a) \Rightarrow (b): Let $P \in I^X$ and x_t be any fuzzy point in X such that $x_t \in \alpha bcl_{\tau} P$ and let $G \in FSO(Y)$ with $f(x_t)qG$. By (a), there exists $U \in F\alpha bO(X)$ with x_tqU , $f(U) \leq cl_{\tau_1} G$. As $x_t \in \alpha bcl_{\tau} P$, by Result 3.11 (1), UqP . Then $f(U)qf(P)$. Thus $f(P)qcl_{\tau_1} G$. So $f(x_t) \in \theta-scl_{\tau_1}(f(P))$. Hence $f(\alpha bcl_{\tau} P) \leq \theta-scl_{\tau_1}(f(P))$.

(b) \Rightarrow (e): Let $R \in I^Y$. Then $f^{-1}(R) \in I^X$. Thus by (b), we have

$$f(\alpha bcl_{\tau}(f^{-1}(R))) \leq \theta-scl_{\tau_1}(f(f^{-1}(R))) \leq \theta-scl_{\tau_1} R.$$

So $\alpha bcl_{\tau}(f^{-1}(R)) \leq f^{-1}(\theta-scl_{\tau_1} R)$.

(e) \Rightarrow (a): Let x_t be any fuzzy point in X and $A \in FSO(Y)$ with $f(x_t)qA$. Since $cl_{\tau_1} A \not/q(1_Y \setminus cl_{\tau_1} A)$, by definition, $f(x_t) \notin \theta-scl_{\tau_1}(1_Y \setminus cl_{\tau_1} A)$. Then $x_t \notin f^{-1}(\theta-scl_{\tau_1}(1_Y \setminus cl_{\tau_1} A))$. By (e), $x_t \notin \alpha bcl_{\tau}(f^{-1}(1_Y \setminus cl_{\tau_1} A))$. Thus there exists $U \in F\alpha bO(X)$ with x_tqU such that $U \not/qf^{-1}(1_Y \setminus cl_{\tau_1} A)$. So $f(U) \not/q(1_Y \setminus cl_{\tau_1} A)$. Hence $f(U) \leq cl_{\tau_1} A$.

(a) \Rightarrow (d): Let $A \in FSO(Y)$ and x_t be any fuzzy point in X such that $x_tqf^{-1}(A)$. Then $f(x_t)qA$. By (a), there exists $U \in F\alpha bO(X)$ with x_tqU such that $f(U) \leq cl_{\tau_1} A$. Thus $x_tqU \leq f^{-1}(cl_{\tau_1} A)$. So $x_tqU = \alpha bint_{\tau} U \leq \alpha bint_{\tau}(f^{-1}(cl_{\tau_1} A))$. Since $\alpha bint_{\tau}(f^{-1}(cl_{\tau_1} A))$ is the union of all fuzzy α - b -open sets in X contained in $f^{-1}(cl_{\tau_1} A)$, $x_tq\alpha bint_{\tau}(f^{-1}(cl_{\tau_1} A))$. Hence $f^{-1}(A) \leq \alpha bint_{\tau}(f^{-1}(cl_{\tau_1} A))$.

(d) \Rightarrow (a): Let x_t be any fuzzy point in X and $A \in FSO(Y)$ with $f(x_t)qA$. Then by (d), $x_tqf^{-1}(A) \leq \alpha bint_{\tau}(f^{-1}(cl_{\tau_1} A))$. Thus there exists $U \in F\alpha bO(X)$ with x_tqU such that $U \leq f^{-1}(cl_{\tau_1} A)$. So $f(U) \leq cl_{\tau_1} A$.

(c) \Rightarrow (d): Let $A \in FSO(Y)$ and x_t be any fuzzy point in X such that $x_t \in f^{-1}(A)$. Then $f(x_t) \in A$. By (c), there exists $U \in F\alpha bO(X)$ with $x_t \in U$ such that $f(U) \leq cl_{\tau_1} A$. Thus $U \leq f^{-1}(cl_{\tau_1} A)$. So $x_t \in U = \alpha bint_{\tau} U \leq \alpha bint_{\tau}(f^{-1}(cl_{\tau_1} A))$. Hence $f^{-1}(A) \leq \alpha bint_{\tau}(f^{-1}(cl_{\tau_1} A))$.

(d) \Rightarrow (c): Let x_t be any fuzzy point in X and $A \in FSO(Y)$ with $f(x_t) \in A$. Then by (d), $x_t \in f^{-1}(A) \leq \alpha bint_{\tau}(f^{-1}(cl_{\tau_1} A))$. Thus there exists $U \in F\alpha bO(X)$ with $x_t \in U$ such that $U \leq f^{-1}(cl_{\tau_1} A)$. So $f(U) \leq cl_{\tau_1} A$.

(2) Suppose (e) holds and let $A \in FRO(Y)$. Then by (e), we get

$$\alpha bcl_{\tau}(f^{-1}(A)) \leq f^{-1}(\theta-scl_{\tau_1} A) = f^{-1}(A).$$

Thus by Lemma 3.13 (1), $f^{-1}(A) \in F\alpha bC(X)$. So f is a fuzzy α - b - r -continuous function. \square

Theorem 4.4. Let (X, τ) and (Y, τ_1) be two fts's and $f : X \rightarrow Y$ be a function satisfying $\alpha bcl_{\tau}(f^{-1}(R)) \leq f^{-1}(\theta-scl_{\tau_1} R)$, for all $R \in I^Y$. Then the following statements hold:

- (1) $\alpha bcl_{\tau}(f^{-1}(R)) \leq f^{-1}(\theta-scl_{\tau_1} R)$ for all $R \in FSO(Y)$
- (2) $\alpha bcl_{\tau}(f^{-1}(R)) \leq f^{-1}(\theta-scl_{\tau_1} R)$ for all $R \in FPO(Y)$,
- (3) $\alpha bcl_{\tau}(f^{-1}(R)) \leq f^{-1}(\theta-scl_{\tau_1} R)$ for all $R \in F\beta O(Y)$.

Proof. Obvious. □

Definition 4.5. Let (X, τ) and (Y, τ_1) be two fts's and $f : X \rightarrow Y$ be a function. Then f is said to be fuzzy

- (i) α - b -continuous, if $f^{-1}(A) \in FabO(X)$ for all $A \in \tau_1$,
- (ii) almost α - b -continuous, if $f^{-1}(A) \in FabO(X)$ for all $A \in FRO(Y)$.

Let us now recall the following definition from [2] for ready references.

Definition 4.6 ([2]). Let (X, τ) and (Y, τ_1) be two fts's and $f : X \rightarrow Y$ be a function. Then f is said to be a *fuzzy continuous function*, if $f^{-1}(U) \in \tau$ for all $U \in \tau_1$.

Remark 4.7. It is clear from definitions that:

(1) fuzzy continuity \Rightarrow fuzzy α - b -continuity \Rightarrow fuzzy almost α - b -continuity, but reverse implications are not necessarily true, in general, follow from the next examples,

(2) fuzzy α - b - r -continuity is an independent concept of fuzzy continuity, fuzzy α - b -continuity and fuzzy almost α - b -continuity, follow from the next examples.

Example 4.8. Fuzzy continuity, fuzzy α - b -continuity and fuzzy almost α - b -continuity $\not\Rightarrow$ fuzzy α - b - r -continuity.

Let $X = \{a, b\}$, $\tau_1 = \{0_X, 1_X, A, B\}$, $\tau_2 = \{0_X, 1_X, B\}$, where $A(a) = A(b) = 0.5, B(a) = B(b) = 0.4$. Then (X, τ_1) and (X, τ_2) are fts's. Consider the identity function $i : (X, \tau_1) \rightarrow (X, \tau_2)$. Then clearly, i is fuzzy continuous and thus fuzzy α - b -continuous as well as fuzzy almost α - b -continuous function. Now $1_X \setminus B \in FRC(X, \tau_2)$. $i^{-1}(1_X \setminus B) = 1_X \setminus B$. Then we get

$$cl_{\tau_1}(\alpha int_{\tau_1}(cl_{\tau_1}(1_X \setminus B))) = A \not\supseteq 1_X \setminus B \Rightarrow 1_X \setminus B \notin FabO(X, \tau_1).$$

Thus i is not a fuzzy α - b - r -continuous function.

Example 4.9. Fuzzy α - b - r -continuity, fuzzy α - b -continuity and fuzzy almost α - b -continuity $\not\Rightarrow$ fuzzy continuity.

Let $X = \{a, b\}$, $\tau_1 = \{0_X, 1_X\}$, $\tau_2 = \{0_X, 1_X, A\}$, where $A(a) = A(b) = 0.5$. Then (X, τ_1) and (X, τ_2) are fts's. Consider the identity function $i : (X, \tau_1) \rightarrow (X, \tau_2)$. Since every fuzzy set in (X, τ_1) is fuzzy α - b -open in (X, τ_1) , i is fuzzy α - b - r -continuous, fuzzy α - b -continuous and fuzzy almost α - b -continuous. Since $A \in \tau_2$, $i^{-1}(A) = A \notin \tau_1$. Then i is not a fuzzy continuous function.

Example 4.10. Fuzzy α - b - r -continuity, fuzzy almost α - b -continuity $\not\Rightarrow$ fuzzy α - b -continuity.

Let $X = \{a, b\}$, $\tau_1 = \{0_X, 1_X, A, B\}$, $\tau_2 = \{0_X, 1_X, C\}$, where $A(a) = A(b) = 0.4, B(a) = B(b) = 0.5, C(a) = 0.5, C(b) = 0.6$. Then (X, τ_1) and (X, τ_2) are fts's. Consider the identity function $i : (X, \tau_1) \rightarrow (X, \tau_2)$. Now $C \in \tau_2$, $i^{-1}(C) = C$. Then $cl_{\tau_1}(\alpha int_{\tau_1}(cl_{\tau_1}C)) = B \not\supseteq C$. Thus $C \notin FabO(X, \tau_1)$. So i is not fuzzy α - b -continuous. Since $0_X, 1_X \in FRO(X, \tau_2)$ only, i is a fuzzy α - b - r -continuous function and a fuzzy almost α - b -continuous function.

Example 4.11. Fuzzy α - b - r -continuity $\not\Rightarrow$ fuzzy almost α - b -continuity.

Let $X = \{a, b\}$, $\tau_1 = \{0_X, 1_X, A\}$, $\tau_2 = \{0_X, 1_X, B\}$, where $A(a) = 0.5, A(b) = 0.6, B(a) = 0.5, B(b) = 0.3$. Then (X, τ_1) and (X, τ_2) are fts's. Consider the identity function $i : (X, \tau_1) \rightarrow (X, \tau_2)$. Now $B \in FRO(X, \tau_2)$, $i^{-1}(B) = B$. Then $int_{\tau_1}(\alpha cl_{\tau_1}(int_{\tau_1} B)) = 0_X \leq B$. Thus $B \in FabC(X, \tau_1)$. So i is a fuzzy α - b - r -continuous function. But $cl_{\tau_1}(\alpha int_{\tau_1}(cl_{\tau_1} B)) = 0_X \not\leq B$, i.e., $B \notin FabO(X, \tau_1)$. Hence i is not a fuzzy almost α - b -continuous function.

Definition 4.12 ([17]). An fts (X, τ) is said to be *fuzzy extremally disconnected*, if the closure of every fuzzy open set in X is fuzzy open in X .

Theorem 4.13. Let (X, τ) and (Y, τ_1) be two fts's and $f : X \rightarrow Y$ be a function. If (Y, τ_1) is a fuzzy extremally disconnected space, then f is a fuzzy α - b - r -continuous function if and only if f is a fuzzy almost α - b -continuous function.

Proof. First suppose that f is fuzzy α - b - r -continuous function and let $U \in FRO(Y)$. Then $U = int_{\tau_1}(cl_{\tau_1} U)$. As Y is fuzzy extremally disconnected, $cl_{\tau_1} U \in \tau_1$. Thus $U = int_{\tau_1} cl_{\tau_1} U = cl_{\tau_1} U = cl_{\tau_1} int_{\tau_1} U$. So $U \in FRC(Y)$. By the hypothesis, $f^{-1}(U) \in FabO(X)$. Hence f is a fuzzy almost α - b -continuous function.

Conversely, suppose f is a fuzzy almost α - b -continuous function and let $U \in FRC(Y)$. As Y is a fuzzy extremally disconnected space, $U \in FRO(Y)$. Then By the hypothesis, $f^{-1}(U) \in FabO(X)$. Thus f is a fuzzy α - b - r -continuous function. \square

Remark 4.14. Composition of two fuzzy α - b - r -continuous (resp. fuzzy α - b -continuous and fuzzy almost α - b -continuous) functions need not be so, as it seen from the following examples.

Example 4.15. Let $X = \{a, b\}$, $\tau_1 = \{0_X, 1_X, A, B\}$, $\tau_2 = \{0_X, 1_X\}$, $\tau_3 = \{0_X, 1_X, B\}$, where $A(a) = A(b) = 0.5, B(a) = B(b) = 0.4$. Then (X, τ_1) , (X, τ_2) and (X, τ_3) are fts's. Consider two identity functions $i_1 : (X, \tau_1) \rightarrow (X, \tau_2)$, $i_2 : (X, \tau_2) \rightarrow (X, \tau_3)$. Clearly, i_1 and i_2 are fuzzy α - b - r -continuous functions. Let $i_3 = i_2 \circ i_1$. Now $1_X \setminus B \in FRC(X, \tau_3)$, $i_3^{-1}(1_X \setminus B) = 1_X \setminus B$. Then we have

$$cl_{\tau_1}(\alpha int_{\tau_1}(cl_{\tau_1}(1_X \setminus B))) = A \not\leq 1_X \setminus B.$$

Thus $1_X \setminus B \notin FabO(X, \tau_1)$. So i_3 is not a fuzzy α - b - r -continuous function.

Example 4.16. Let $X = \{a, b\}$, $\tau_1 = \{0_X, 1_X, A\}$, $\tau_2 = \{0_X, 1_X\}$, $\tau_3 = \{0_X, 1_X, B\}$, where $A(a) = 0.5, A(b) = 0.6, B(a) = 0.5, B(b) = 0.3$. Then (X, τ_1) , (X, τ_2) and (X, τ_3) are fts's. Consider two identity functions $i_1 : (X, \tau_1) \rightarrow (X, \tau_2)$ and $i_2 : (X, \tau_2) \rightarrow (X, \tau_3)$. Clearly, i_1 and i_2 are fuzzy α - b -continuous and thus fuzzy almost α - b -continuous functions. Let $i_3 = i_2 \circ i_1$. Now $B \in \tau_3$ as well as $B \in FRO(X, \tau_3)$. $i_3^{-1}(B) = B$. Now $cl_{\tau_1}(\alpha int_{\tau_1}(cl_{\tau_1} B)) = 0_X \not\leq B \Rightarrow B \notin FabO(X, \tau_1) \Rightarrow i_3$ is not a fuzzy α - b -continuous and also fuzzy almost α - b -continuous function.

5. FUZZY α - b -REGULAR, α - b -COMPACT AND α - b - T_2 -SPACES

In this section, new types of separation axioms and compactness are introduced and studied. Then the mutual relationships of these spaces with the spaces defined in [2, 4] are established.

Definition 5.1. An fts (X, τ) is called an α -*b-regular space*, if for each fuzzy point x_t in X and each fuzzy α -*b-closed set* F with $x_t \notin F$, there exist a fuzzy open set U and a fuzzy α -*b-open set* V in X such that $x_t q U$, $F \leq V$ and $U \not q V$.

Theorem 5.2. For an fts (X, τ) , the following statements are equivalent:

- (1) X is fuzzy α -*b-regular*,
- (2) for each fuzzy point x_t in X and each fuzzy α -*b-open set* U in X with $x_t q U$, there exists a fuzzy open set V in X such that $x_t q V \leq \alpha bcl V \leq U$,
- (3) for each fuzzy α -*b-closed set* F in X , $\bigwedge \{clV : F \leq V, V \in FabO(X)\} = F$,
- (4) for each fuzzy set G in X and each fuzzy α -*b-open set* U in X such that $G q U$, there exists a fuzzy open set V in X such that $G q V$ and $\alpha bcl V \leq U$.

Proof. (1) \Rightarrow (2): Let x_t be a fuzzy point in X and let U be a fuzzy α -*b-open set* in X with $x_t q U$. Then $x_t \notin 1_X \setminus U \in FabC(X)$. By (1), there exist a fuzzy open set V and a fuzzy α -*b-open set* W in X such that $x_t q V$, $1_X \setminus U \leq W$ and $V \not q W$. Thus $x_t q V \leq 1_X \setminus W \leq U$. So $x_t q V \leq \alpha bcl V \leq \alpha bcl(1_X \setminus W) = 1_X \setminus W \leq U$.

(2) \Rightarrow (1): Let F be a fuzzy α -*b-closed set* in X and let x_t be a fuzzy point in X with $x_t \notin F$. Then $x_t q(1_X \setminus F) \in FabO(X)$. By (2), there exists a fuzzy open set V in X such that $x_t q V \leq \alpha bcl V \leq 1_X \setminus F$. Put $U = 1_X \setminus \alpha bcl V$. Then $U \in FabO(X)$ and $x_t q U$, $F \leq U$ and $U \not q V$.

(2) \Rightarrow (3): Let F be fuzzy α -*b-closed set* in X . Then we get

$$F \leq \bigwedge \{clV : F \leq V, V \in FabO(X)\}.$$

Conversely, let $x_t \notin F \in FabC(X)$. Then $F(x) < t$. Thus $x_t q(1_X \setminus F)$, where $1_X \setminus F \in FabO(X)$. By (2), there exists a fuzzy open set U in X such that $x_t q U \leq \alpha bcl U \leq 1_X \setminus F$. Put $V = 1_X \setminus \alpha bcl U$. Then $F \leq V$ and $U \not q V$. Thus $x_t \notin clV$. So $\bigwedge \{clV : F \leq V, V \in FabO(X)\} \leq F$. Hence $\bigwedge \{clV : F \leq V, V \in FabO(X)\} = F$.

(3) \Rightarrow (2): Let V be any fuzzy α -*b-open set* in X and let x_t be any fuzzy point in X with $x_t q V$. Then $V(x) + t > 1$, i.e., $x_t \notin (1_X \setminus V)$, where $1_X \setminus V \in FabC(X)$. By (3), there exists $G \in FabO(X)$ such that $1_X \setminus V \leq G$ and $x_t \notin clG$. Thus there exists a fuzzy open set U in X with $x_t q U$ such that $U \not q G$. So $U \leq 1_X \setminus G \leq V$. Hence $x_t q U \leq \alpha bcl U \leq \alpha bcl(1_X \setminus G) = 1_X \setminus G \leq V$.

(3) \Rightarrow (4): Let G be any fuzzy set in X and let U be any fuzzy α -*b-open set* in X with $G q U$. Then there exists $x \in X$ such that $G(x) + U(x) > 1$. Let $G(x) = t$. Then $x_t q U \Rightarrow x_t \notin 1_X \setminus U$, where $1_X \setminus U \in FabC(X)$. By (3), there exists $W \in FabO(X)$ such that $1_X \setminus U \leq W$ and $x_t \notin clW$. Thus $(clW)(x) < t$. So $x_t q(1_X \setminus clW)$. Let $V = 1_X \setminus clW$. Then V is fuzzy open set in X and $V(x) + t > 1$. Thus $V(x) + G(x) > 1$. So $V q G$ and $\alpha bcl V = \alpha bcl(1_X \setminus clW) \leq \alpha bcl(1_X \setminus W) = 1_X \setminus W \leq U$.

(4) \Rightarrow (2): Obvious. □

Note 5.3. It is clear from Theorem 5.2 that in a fuzzy α -*b-regular space*, every fuzzy α -*b-closed set* is fuzzy closed and hence every fuzzy α -*b-open set* is fuzzy open. As a result, in a fuzzy α -*b-regular space*, the collection of all fuzzy closed (resp., fuzzy open) sets and fuzzy α -*b-closed* (resp., fuzzy α -*b-open*) sets coincide.

Definition 5.4. Let A be a fuzzy set in X . A collection \mathcal{U} of fuzzy sets in X is called a *fuzzy cover* of A , if $\sup\{U(x) : U \in \mathcal{U}\} = 1$ for each $x \in \text{supp}A$ [18]. In particular, if $A = 1_X$, we get the definition of fuzzy cover of X [2].

Definition 5.5. A fuzzy cover \mathcal{U} of a fuzzy set A in X is said to *have a finite subcover* \mathcal{U}_0 , if \mathcal{U}_0 is a finite subcollection of \mathcal{U} such that $\bigcup \mathcal{U}_0 \geq A$, i.e., \mathcal{U}_0 is also a fuzzy cover of A [18]. In particular, if $A = 1_X$, we get $\bigcup \mathcal{U}_0 = 1_X$ [2].

Definition 5.6. A fuzzy set A in an fts (X, τ) is said to be *fuzzy compact* [18], if every fuzzy covering \mathcal{U} of A by fuzzy open sets in X has a finite subcovering \mathcal{U}_0 of \mathcal{U} . In particular, if $A = 1_X$, we get the definition of fuzzy compact space [2].

Definition 5.7. An fts (X, τ) is said to be *fuzzy s-closed* [19] (resp. *fuzzy nearly compact* [17]), if every fuzzy covering of X by fuzzy regular closed (resp. fuzzy regular open) sets of X contains a finite subcovering.

Let us now introduce the following concept.

Definition 5.8. A fuzzy set A in an fts (X, τ) is called *fuzzy α -b-compact*, if every fuzzy covering of A by fuzzy α -b-open sets of X has a finite subcovering. In particular, if $A = 1_X$, we get the definition of fuzzy α -b-compact space.

Remark 5.9. It is clear from above discussion that fuzzy α -b-compact space is fuzzy compact. But the converse is not necessarily true follows from the next example.

Example 5.10. Let $X = \{a\}$, $\tau = \{0_X, 1_X\}$. The clearly (X, τ) is a fuzzy compact space. Here every fuzzy set is fuzzy α -b-open set in X . Consider the fuzzy cover $\mathcal{U} = \{U_n(a) : n \in \mathbf{N}\}$, where $U_n(a) = \{\frac{n}{n+1} : n \in \mathbf{N}\}$. Then \mathcal{U} is a fuzzy α -b-open cover of X . But it does not have any subcovering of X . Thus X is not fuzzy α -b-compact space.

Theorem 5.11. *Every fuzzy α -b-closed set A in a fuzzy α -b-compact space X is fuzzy α -b-compact.*

Proof. Let A be a fuzzy α -b-closed set in a fuzzy α -b-compact space X and let \mathcal{U} be a fuzzy covering of A by fuzzy α -b-open sets in X . Then $\mathcal{V} = \mathcal{U} \cup (1_X \setminus A)$ is a fuzzy α -b-open covering of X . By the hypothesis, there exists a finite subcollection \mathcal{V}_0 of \mathcal{V} which also covers X . If \mathcal{V}_0 contains $1_X \setminus A$, we omit it and get a finite subcovering of A . Consequently, A is fuzzy α -b-compact. \square

Let us now recall the following definition from [4] for ready references.

Definition 5.12 ([4]). Let (X, τ) be an fts. Then X is said to be a *fuzzy T_2 -space*, if for each pair of distinct fuzzy points x_α, y_β : when $x \neq y$, there exist fuzzy open sets U_1, U_2, V_1, V_2 such that $x_\alpha \in U_1, y_\beta q V_1$ and $U_1 \not q V_1$ and $x_\alpha q U_2, y_\beta \in V_2$ and $U_2 \not q V_2$; when $x = y, \alpha < \beta$ (say), there exist fuzzy open sets U, V in X such that $x_\alpha \in U, y_\beta q V$ and $U \not q V$.

Now we introduce the following concept.

Definition 5.13. Let (X, τ) be an fts. Then X is said to be a *fuzzy α -b- T_2 -space*, if for each pair of distinct fuzzy points x_α, y_β : when $x \neq y$, there exist fuzzy α -b-open sets U_1, U_2, V_1, V_2 such that $x_\alpha \in U_1, y_\beta q V_1$ and $U_1 \not q V_1$ and $x_\alpha q U_2, y_\beta \in V_2$ and $U_2 \not q V_2$; when $x = y, \alpha < \beta$ (say), there exist fuzzy α -b-open sets U, V in X such that $x_\alpha \in U, y_\beta q V$ and $U \not q V$.

Let us now recall the following definition from [4] for ready references.

Definition 5.14 ([4]). An fts (X, τ) is said to be a *fuzzy regular space*, if for any fuzzy point x_t in X and any fuzzy closed set F in X with $x_t \notin F$, there exist fuzzy open sets U, V in X such that $x_t q U, F \leq V$ and $U \not q V$.

Remark 5.15. It is clear from Note 5.3 that fuzzy α - b -regular space is fuzzy regular and fuzzy T_2 -space is fuzzy α - b - T_2 -space. But the reverse implications are not necessarily true, follow from the next example.

Example 5.16. Consider Example 5.10. It is clear that (X, τ) is fuzzy regular and fuzzy α - b - T_2 -space (as every fuzzy set is fuzzy α - b -open set as well as fuzzy α - b -closed set). Now consider the fuzzy point $a_{0.4}$ and a fuzzy set A defined by $A(a) = 0.3$. Then $a_{0.4} \notin A \in F\alpha bC(X)$. But there do not exist any fuzzy open set U and a fuzzy α - b -open set V in X such that $a_{0.4} q U, A \leq V$ and $U \not q V$ (because 1_X is the only fuzzy open set in X with $a_{0.4} q 1_X$ and $1_X q V$ for all fuzzy set $V (\neq 0_X)$ in X). Thus X is not fuzzy α - b -regular space.

Consider two fuzzy points $a_{0.4}$ and $a_{0.5}$ in X . But there do not exist fuzzy open sets U, V in X such that $a_{0.4} \in U, a_{0.5} q V$ and $U \not q V$. So X is not fuzzy T_2 -space.

6. APPLICATIONS OF FUZZY α - b - r -CONTINUOUS, α - b -CONTINUOUS AND ALMOST α - b -CONTINUOUS FUNCTIONS

In this section, the applications of the functions introduced in this paper are established. First we recall the following definition from [20] for ready references.

Definition 6.1 ([20]). A function $f : X \rightarrow Y$ is said to be a *fuzzy open function*, if $f(U)$ is a fuzzy open set in Y for every fuzzy open set U in X .

Theorem 6.2. Let (X, τ) and (Y, τ_1) be two fts's and $f : X \rightarrow Y$ be a surjective, fuzzy α - b - r -continuous function. If X is a fuzzy α - b -compact space, then Y is a fuzzy s -closed space.

Proof. Let $\mathcal{U} = \{U_\alpha : \alpha \in \Lambda\}$ be a fuzzy covering of Y by fuzzy regular closed sets of Y . As f is a fuzzy α - b - r -continuous function, $\mathcal{V} = \{f^{-1}(U_\alpha) : \alpha \in \Lambda\}$ covers X by fuzzy α - b -open sets of X . As X is a fuzzy α - b -compact space, there exists a finite subset Λ_0 of Λ such that $1_X = \bigvee_{\alpha \in \Lambda_0} f^{-1}(U_\alpha)$. Then we have

$$1_Y = f\left(\bigvee_{\alpha \in \Lambda_0} f^{-1}(U_\alpha)\right) = \bigvee_{\alpha \in \Lambda_0} f(f^{-1}(U_\alpha)) \leq \bigvee_{\alpha \in \Lambda_0} U_\alpha.$$

Thus Y is a fuzzy s -closed space. □

Theorem 6.3. Let (X, τ) and (Y, τ_1) be two fts's and $f : X \rightarrow Y$ be a fuzzy α - b -continuous function. If A is fuzzy α - b -compact set relative to X , then the image $f(A)$ is fuzzy compact relative to Y .

Proof. Let A be fuzzy α - b -compact relative to X and $\mathcal{U} = \{U_\alpha : \alpha \in \Lambda\}$ be a fuzzy covering of $f(A)$ by fuzzy open sets of Y , i.e., $f(A) \leq \bigvee_{\alpha \in \Lambda} U_\alpha$. Then $A \leq$

$f^{-1}\left(\bigvee_{\alpha \in \Lambda} U_\alpha\right) = \bigvee_{\alpha \in \Lambda} f^{-1}(U_\alpha)$. Thus $\mathcal{V} = \{f^{-1}(U_\alpha) : \alpha \in \Lambda\}$ is a fuzzy covering of A by fuzzy α - b -open sets in X . As A is fuzzy α - b -compact set relative to X ,

there exists a finite subcollection $\mathcal{V}_0 = \{f^{-1}(U_{\alpha_i}) : 1 \leq i \leq n\}$ of \mathcal{V} such that $A \leq \bigvee_{i=1}^n f^{-1}(U_{\alpha_i})$. So $f(A) \leq f(\bigvee_{i=1}^n f^{-1}(U_{\alpha_i})) = \bigvee_{i=1}^n f(f^{-1}(U_{\alpha_i})) \leq \bigvee_{i=1}^n U_{\alpha_i}$. Hence $\mathcal{U}_0 = \{U_{\alpha_i} : 1 \leq i \leq n\}$ is a finite subcovering of $f(A)$. Therefore $f(A)$ is fuzzy compact relative to Y . \square

Theorem 6.4. *Let (X, τ) and (Y, τ_1) be two fts's and $f : X \rightarrow Y$ be a fuzzy almost α - b -continuous function. If A is fuzzy α - b -compact relative to X , then the image $f(A)$ is fuzzy nearly compact relative to Y .*

Proof. The proof is similar to that of Theorem 6.3. \square

Theorem 6.5. *Let (X, τ) and (Y, τ_1) be two fts's and $f : X \rightarrow Y$ be an injective, fuzzy α - b -continuous function and Y is a fuzzy T_2 -space. Then X is a fuzzy α - b - T_2 -space.*

Proof. Let x_α and y_β be two distinct fuzzy points in X , where $x \neq y$. As f is injective, $f(x_\alpha) \neq f(y_\beta)$. As Y is a fuzzy T_2 -space, there exist fuzzy open sets U_1, U_2, V_1, V_2 in Y such that $f(x_\alpha) \in U_1, f(y_\beta)qV_1$ and $U_1 \not/qV_1$ and $f(x_\alpha)qU_2, f(y_\beta) \in V_2$ and $U_2 \not/qV_2$. Then $x_\alpha \in f^{-1}(U_1), y_\beta qf^{-1}(V_1)$ and $f^{-1}(U_1) \not/qf^{-1}(V_1)$. Assume that $f^{-1}(U_1)qf^{-1}(V_1)$. Then there exists $z \in X$ such that

$$f^{-1}(U_1)(z) + f^{-1}(V_1)(z) > 1.$$

Thus $U_1(f(z)) + V_1(f(z)) > 1$. So U_1qV_1 . This is a contradiction. Also, we have

$$x_\alpha qf^{-1}(U_2), y_\beta \in f^{-1}(V_2) \text{ and } f^{-1}(U_2) \not/qf^{-1}(V_2),$$

where $f^{-1}(U_1), f^{-1}(V_1), f^{-1}(U_2), f^{-1}(V_2) \in F\alpha bO(X, \tau_1)$.

Similarly, when $x = y, \alpha < \beta$ (say), there exist $U_1, U_2 \in \tau_1$ such that

$$f(x_\alpha) \in U_1, f(y_\beta)qU_2 \text{ and } U_1 \not/qU_2.$$

Then $x_\alpha \in f^{-1}(U_1), y_\beta qf^{-1}(U_2)$ and $f^{-1}(U_1) \not/qf^{-1}(U_2)$, where $f^{-1}(U_1), f^{-1}(U_2) \in F\alpha bO(X, \tau_1)$. Thus X is a fuzzy α - b - T_2 -space. \square

Theorem 6.6. *If a bijective function $h : X \rightarrow Y$ is a fuzzy α - b -continuous, fuzzy open function from a fuzzy α - b -regular space X onto an fts Y , then Y is fuzzy regular space.*

Proof. Let y_α be a fuzzy point in Y and F , a fuzzy closed set in Y with $y_\alpha \notin F$. As h is bijective, there exists unique $x \in X$ such that $h(x) = y$. Then $h(x_\alpha) \notin F$. As h is fuzzy α - b -continuous function, $x_\alpha \notin h^{-1}(F) \in F\alpha bC(X)$. As X is fuzzy α - b -regular space, there exist a fuzzy open set U and a fuzzy α - b -open set V in X such that

$$x_\alpha qU, h^{-1}(F) \leq V \text{ and } U \not/qV.$$

Since X is fuzzy α - b -regular, by Note 5.3, V is also fuzzy open set in X . As h is fuzzy open function, we have $h(x_\alpha)qh(U), F \leq h(V)$ and $h(U) \not/qh(V)$, where $h(U), h(V)$ are fuzzy open sets in Y . Thus Y is a fuzzy regular space. \square

7. CONCLUSIONS

In this paper we have introduced and characterized only fuzzy α - b -regularity, fuzzy α - b -compactness and fuzzy α - b - T_2 property. A thorough discussion on these three spaces are done in a separate article which has already been communicated.

Acknowledgements. I express my sincere gratitude to the referees for their valuable remark on this article.

REFERENCES

- [1] A. S. Bin Shahna, On fuzzy strong semicontinuity and fuzzy precontinuity, *Fuzzy Sets and Systems* 44 (1991), 303–308.
- [2] C. L. Chang, Fuzzy topological spaces, *J. Math. Anal. Appl.* 24 (1968) 182–190.
- [3] K. K. Azad, On fuzzy semi-continuity, fuzzy almost continuity and fuzzy weakly continuity, *J. Math. Anal. Appl.* 82 (1981) 14–32.
- [4] B. Hutton and I. Reilly, Separation axioms in fuzzy topological spaces, *Fuzzy sets and Systems* 31 (1980) 93–104.
- [5] J. G. Lee, G. Şenel, P. K. Lim, J. Kim and K. Hur, Octahedron sets, *Ann. Fuzzy Math. Inform.* 19 (3) (2020) 211–238.
- [6] G. Şenel and N. Çağman, Soft closed sets on soft bitopological space, *Journal of New Results in Science* 3 (5) (2014) 57–66.
- [7] G. Şenel and N. Çağman, Soft topological subspaces, *Ann. Fuzzy Math. Inform.* 10 (4) (2015) 525–535.
- [8] G. Şenel, A new approach to Hausdorff space theory via the soft sets, *Mathematical Problems in Engineering* 1–6, Doi:10.1155/2016/2196743.
- [9] G. Şenel, Soft topology generated by L -Soft sets, *Journal of New Theory* 24 (4) (2018) 88–100.
- [10] G. Şenel, J. G. Lee and K. Hur, Distance and similarity measures for octahedron sets and their applications to MCGDM problems, *Mathematics* 8 (2020) 16–90.
- [11] L. A. Zadeh, Fuzzy Sets, *Information and Control* 8 (1965) 338–353.
- [12] Pao Ming Pu and Ying Ming Liu, Fuzzy topology I. Neighbourhood structure of a fuzzy point and Moore-Smith Convergence, *J. Math. Anal. Appl.* 76 (1980) 571–599.
- [13] S. Nanda, Strongly compact fuzzy topological spaces, *Fuzzy Sets and Systems* 42 (1991) 259–262.
- [14] M. A. Fath Alla, On fuzzy topological spaces, Ph.D. Thesis, Assiut univ., Sohag, Egypt (1984).
- [15] Anjana Bhattacharyya, Concerning almost quasi continuous fuzzy multifunctions, *Universitatea Din Bacău Studii Şi Cercetări, Ştiinţifice Seria : Matematică* 11 (2001) 35–48.
- [16] Anjana Bhattacharyya, Several concepts of continuity in fuzzy setting, *The 10th International Conference MSAST 2016, December 21-23 (2016), Kolkata, 282–293.*
- [17] M. N. Mukherjee and B. Ghosh, On nearly compact and θ -rigid fuzzy sets in fuzzy topological spaces, *Fuzzy Sets and Systems* 43 (1991) 57–68.
- [18] S. Ganguly and S. Saha, A note on compactness in fuzzy setting, *Fuzzy Sets and Systems* 34 (1990) 117–124.
- [19] S. P. Sinha and S. Malakar, On s -closed fuzzy topological spaces, *J. Fuzzy Math.* 2 (1) (1994) 95–103.
- [20] C. K. Wong, Fuzzy points and local properties of fuzzy topology, *J. Math. Anal. Appl.* 46 (1974) 316–328.

ANJANA BHATTACHARYYA (anjanabhattacharyya@hotmail.com)

Department of Mathematics, Victoria Institution (College), 78 B, A.P.C. Road, Kolkata - 700009, India