

Generalized fuzzy sets in UP-algebras

NOPPHARAT DOKKHAMDANG, AKEKARIN KESORN, AIYARED IAMPAN

Received 30 April 2018; Revised 23 May 2018; Accepted 3 June 2018

ABSTRACT. Based on the theory of fuzzy sets, the notion of fuzzy UP-subalgebras (resp., fuzzy UP-filters, fuzzy UP-ideals, fuzzy strongly UP-ideals) with thresholds of UP-algebras is introduced, some properties of them are discussed, and its generalizations are proved. Further, we discuss the relations between fuzzy UP-subalgebras (resp., fuzzy UP-filters, fuzzy UP-ideals, fuzzy strongly UP-ideals) with thresholds and their level subsets.

2010 AMS Classification: 03G25

Keywords: UP-algebra, Fuzzy UP-subalgebra with thresholds, Fuzzy UP-filter with thresholds, Fuzzy UP-ideal with thresholds, Fuzzy strongly UP-ideal with thresholds.

Corresponding Author: Aiyared Iampan (aiyared.ia@up.ac.th)

1. INTRODUCTION

A fuzzy set in a nonempty set X is an arbitrary function from the set X into $[0, 1]$ where $[0, 1]$ is the unit segment of the real line. The concept of a fuzzy set in a nonempty set was first considered by Zadeh [16] in 1965. The fuzzy set theories developed by Zadeh and others have found many applications in the domain of mathematics and elsewhere. There are several kinds of fuzzy set extensions in the fuzzy set theory, for example, intuitionistic fuzzy sets, interval-valued fuzzy sets, vague sets, bipolar-valued fuzzy sets etc.

After the introduction of the notion of fuzzy sets by Zadeh [16], several researches were conducted on the generalizations of the notion of fuzzy sets and application to many logical algebras such as: In 2005, Jun [5] introduced the notion of (α, β) -fuzzy subalgebras of BCK/BCI-algebras. In 2007, Jun [6] introduced the notion of fuzzy subalgebras with thresholds of BCK/BCI-algebras. In 2009, Saeid [11] introduced the notion of new fuzzy subalgebras with thresholds of BCK/BCI-algebras. Zhan et al. [17] introduced the notions of $(\in, \in \vee q)$ -fuzzy p -ideals, $(\in, \in \vee q)$ -fuzzy q -ideals

and $(\in, \in \vee q)$ -fuzzy a -ideals in BCI-algebras. In 2010, Akram and Dar [1] studied generalized fuzzy K-algebras. In 2015, Senapati [12] introduced the notion of T-fuzzy subalgebras of KU-algebras.

Iampan [3] introduced a new branch of the logical algebra, called a UP-algebra. Later Guntasow et al. [2] studied fuzzy translations of a fuzzy set in UP-algebras. Senapati et al. [13, 14] applied cubic set and interval-valued intuitionistic fuzzy structure in UP-algebras. In this paper, we introduce the notion of fuzzy UP-subalgebras (resp., fuzzy UP-filters, fuzzy UP-ideals, fuzzy strongly UP-ideals) with thresholds of UP-algebras, investigate their properties, and generalize the notions. Also, the characterizations of fuzzy UP-subalgebras (resp., fuzzy UP-filters, fuzzy UP-ideals, fuzzy strongly UP-ideals) with thresholds are obtained. Further, we discuss the relations between fuzzy UP-subalgebras (resp., fuzzy UP-filters, fuzzy UP-ideals, fuzzy strongly UP-ideals) with thresholds and their level subsets.

2. BASIC RESULTS ON UP-ALGEBRAS

Before we begin our study, we will introduce the definition of a UP-algebra.

Definition 2.1 ([3]). An algebra $A = (A, \cdot, 0)$ of type $(2, 0)$ is called a UP-algebra, where A is a nonempty set, \cdot is a binary operation on A and 0 is a fixed element of A (i.e., a nullary operation), if it satisfies the following axioms: for any $x, y, z \in A$,

$$(UP-1) (y \cdot z) \cdot ((x \cdot y) \cdot (x \cdot z)) = 0,$$

$$(UP-2) 0 \cdot x = x,$$

$$(UP-3) x \cdot 0 = 0,$$

$$(UP-4) x \cdot y = 0 \text{ and } y \cdot x = 0 \text{ imply } x = y.$$

From [3], we know that the notion of UP-algebras is a generalization of KU-algebras.

Example 2.2 ([3]). Let X be a universal set. Define two binary operations \cdot and $*$ on the power set of X by putting $A \cdot B = B \cap A'$ and $A * B = B \cup A'$, for all $A, B \in \mathcal{P}(X)$. Then $(\mathcal{P}(X), \cdot, \emptyset)$ and $(\mathcal{P}(X), *, X)$ are UP-algebras and we shall call it the power UP-algebra of type 1 and the power UP-algebra of type 2, respectively.

Example 2.3. Let \mathbb{N} be the set of all natural numbers with two binary operations \circ and \bullet defined by

$$x \circ y = \begin{cases} y & \text{if } x < y, \\ 0 & \text{otherwise} \end{cases}$$

and

$$x \bullet y = \begin{cases} y & \text{if } x > y \text{ or } x = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Then $(\mathbb{N}, \circ, 0)$ and $(\mathbb{N}, \bullet, 0)$ are UP-algebras.

For more examples of UP-algebras we refer to [13, 14].

In what follows, let A be a UP-algebra unless otherwise specified. The following proposition is very important for the study of UP-algebras.

Proposition 2.4 ([3]). *The following properties hold: for any $x, y, z \in A$,*

$$(1) x \cdot x = 0,$$

$$(2) x \cdot y = 0 \text{ and } y \cdot z = 0 \text{ imply } x \cdot z = 0,$$

- (3) $x \cdot y = 0$ implies $(z \cdot x) \cdot (z \cdot y) = 0$,
- (4) $x \cdot y = 0$ implies $(y \cdot z) \cdot (x \cdot z) = 0$,
- (5) $x \cdot (y \cdot x) = 0$,
- (6) $(y \cdot x) \cdot x = 0$ if and only if $x = y \cdot x$, and
- (7) $x \cdot (y \cdot y) = 0$.

Definition 2.5 ([3]). A subset S of A is called a *UP-subalgebra* of A , if the constant 0 of A is in S and $(S, \cdot, 0)$ itself forms a UP-algebra.

Iampan [3] proved the useful criteria that a nonempty subset S of a UP-algebra $A = (A, \cdot, 0)$ is a UP-subalgebra of A if and only if S is closed under the \cdot multiplication on A .

Definition 2.6 ([3, 15]). A subset S of A is called a

- (1) UP-ideal of A , if it satisfies the following properties:
 - (i) the constant 0 of A is in S ,
 - (ii) for any $x, y, z \in A, x \cdot (y \cdot z) \in S$ and $y \in S$ imply $x \cdot z \in S$.
- (2) UP-filter of A , if it satisfies the following properties:
 - (i) the constant 0 of A is in S ,
 - (ii) for any $x, y \in A, x \cdot y \in S$ and $x \in S$ imply $y \in S$.
- (3) strongly UP-ideal of A , if it satisfies the following properties:
 - (i) the constant 0 of A is in S ,
 - (ii) for any $x, y, z \in A, (z \cdot y) \cdot (z \cdot x) \in S$ and $y \in S$ imply $x \in S$.

Guntasow et al. [2] proved the generalization that the notion of UP-subalgebras is a generalization of UP-filters, the notion of UP-filters is a generalization of UP-ideals, and the notion of UP-ideals is a generalization of strongly UP-ideals. Moreover, they also proved that a UP-algebra A is the only one strongly UP-ideal of itself.

Definition 2.7 ([16]). A *fuzzy set* in a nonempty set X (or a fuzzy subset of X) is an arbitrary function from the set X into $[0, 1]$, where $[0, 1]$ is the unit segment of the real line.

Somjanta et al. [15] introduced the notion of fuzzy UP-subalgebras (resp., fuzzy UP-filters, fuzzy UP-ideals, fuzzy strongly UP-ideals) of UP-algebras as follows.

Definition 2.8. A fuzzy set f in A is called a

- (1) fuzzy UP-subalgebra of A , if for any $x, y \in A, f(x \cdot y) \geq \min\{f(x), f(y)\}$.
- (2) fuzzy UP-filter of A if for any $x, y \in A$,
 - (i) $f(0) \geq f(x)$,
 - (ii) $f(y) \geq \min\{f(x \cdot y), f(x)\}$.
- (3) fuzzy UP-ideal of A , if for any $x, y, z \in A$,
 - (i) $f(0) \geq f(x)$,
 - (ii) $f(x \cdot z) \geq \min\{f(x \cdot (y \cdot z)), f(y)\}$.
- (4) fuzzy strongly UP-ideal of A if for any $x, y, z \in A$,
 - (i) $f(0) \geq f(x)$,
 - (ii) $f(x) \geq \min\{f((z \cdot y) \cdot (z \cdot x)), f(y)\}$.

They also proved that the notion of fuzzy UP-subalgebras is a generalization of fuzzy UP-filters, the notion of fuzzy UP-filters is a generalization of fuzzy UP-ideals, and the notion of fuzzy UP-ideals is a generalization of fuzzy strongly UP-ideals.

Definition 2.9 ([8]). Let X and Y be any two nonempty sets and let $f: X \rightarrow Y$ be any function. A fuzzy set μ in X is called f -invariant, if $f(x) = f(y)$ implies $\mu(x) = \mu(y)$, for all $x, y \in X$.

Definition 2.10 ([3]). Let $(A, \cdot, 0_A)$ and $(B, *, 0_B)$ be UP-algebras. A mapping f from A to B is called a UP-homomorphism, if

$$f(x \cdot y) = f(x) * f(y), \text{ for all } x, y \in A.$$

Iampan [3] proved that $f(0_A) = 0_B$.

3. FUZZY UP-SUBALGEBRAS (RESP., FUZZY UP-FILTERS, FUZZY UP-IDEALS, FUZZY STRONGLY UP-IDEALS) WITH THRESHOLDS

In this section, we introduce the notion of fuzzy UP-subalgebras (resp., fuzzy UP-filters, fuzzy UP-ideals, fuzzy strongly UP-ideals) with thresholds of UP-algebras, investigate their properties, and generalize the notions.

Definition 3.1. A fuzzy set f in A is called a fuzzy UP-subalgebra with thresholds ε and δ of A , where $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$, if for any $x, y \in A$,

$$\max\{f(x \cdot y), \varepsilon\} \geq \min\{f(x), f(y), \delta\}.$$

Example 3.2. Let $A = \{0, 1, 2, 3\}$ be a set with a binary operation \cdot defined by the following Cayley table:

\cdot	0	1	2	3
0	0	1	2	3
1	0	0	3	0
2	0	1	0	0
3	0	1	3	0

Then $(A, \cdot, 0)$ is a UP-algebra. We defined a fuzzy set $f: A \rightarrow [0, 1]$ as follows:

$$f(0) = 0.6, f(1) = 0.7, f(2) = 0.3, \text{ and } f(3) = 0.2.$$

Then f is a fuzzy UP-subalgebra with thresholds $\varepsilon = 0.8$ and $\delta = 0.9$ of A .

Lemma 3.3. *If f is a fuzzy UP-subalgebra with thresholds ε and δ of A , then $\max\{f(0), \varepsilon\} \geq \min\{f(x), \delta\}$ for all $x \in A$. In particular, if there exists $y \in A$ such that $f(y) > \delta$, then $f(0) > \varepsilon$.*

Proof. For all $x \in A$,

$$\begin{aligned} \text{(Proposition 2.4 (1))} \quad \max\{f(0), \varepsilon\} &= \max\{f(x \cdot x), \varepsilon\} \\ &\geq \min\{f(x), f(x), \delta\} \\ &= \min\{f(x), \delta\}. \end{aligned}$$

If there exists $y \in A$ such that $f(y) > \delta$, then

$$\max\{f(0), \varepsilon\} \geq \min\{f(y), \delta\} = \delta > \varepsilon.$$

Thus $f(0) = \max\{f(0), \varepsilon\} > \varepsilon$. □

Definition 3.4. A fuzzy set f in A is called a fuzzy UP-filter with thresholds ε and δ of A , where $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$, if it satisfies the following properties: for any $x, y \in A$,

- (1) $\max\{f(0), \varepsilon\} \geq \min\{f(x), \delta\}$,
- (2) $\max\{f(y), \varepsilon\} \geq \min\{f(x \cdot y), f(x), \delta\}$.

Example 3.5. Let $A = \{0, 1, 2, 3, 4\}$ be a set with a binary operation \cdot defined by the following Cayley table:

\cdot	0	1	2	3	4
0	0	1	2	3	4
1	0	0	2	0	0
2	0	1	0	0	0
3	0	1	2	0	4
4	0	1	2	3	0

Then $(A, \cdot, 0)$ is a UP-algebra. We defined a fuzzy set $f : A \rightarrow [0, 1]$ as follows:

$$f(0) = 0.9, f(1) = 0.1, f(2) = 0.5, f(3) = 0.4, \text{ and } f(4) = 0.4.$$

Then f is a fuzzy UP-filter with thresholds $\varepsilon = 0.8$ and $\delta = 0.9$ of A .

Definition 3.6. A fuzzy set f in A is called a fuzzy UP-ideal with thresholds ε and δ of A , where $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$, if it satisfies the following properties: for any $x, y, z \in A$,

- (1) $\max\{f(0), \varepsilon\} \geq \min\{f(x), \delta\}$,
- (2) $\max\{f(x \cdot z), \varepsilon\} \geq \min\{f(x \cdot (y \cdot z)), f(y), \delta\}$.

Example 3.7. Let $A = \{0, 1, 2, 3\}$ be a set with a binary operation \cdot defined by the following Cayley table:

\cdot	0	1	2	3
0	0	1	2	3
1	0	0	3	3
2	0	1	0	0
3	0	1	3	0

Then $(A, \cdot, 0)$ is a UP-algebra. We defined a fuzzy set $f : A \rightarrow [0, 1]$ as follows:

$$f(0) = 0, f(1) = 0.2, f(2) = 0.1, \text{ and } f(3) = 0.3.$$

Then f is a fuzzy UP-ideal with thresholds $\varepsilon = 0.4$ and $\delta = 0.9$ of A .

Definition 3.8. A fuzzy set f in A is called a fuzzy strongly UP-ideal with thresholds ε and δ of A , where $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$, if it satisfies the following properties: for any $x, y, z \in A$,

- (1) $\max\{f(0), \varepsilon\} \geq \min\{f(x), \delta\}$,
- (2) $\max\{f(x), \varepsilon\} \geq \min\{f((z \cdot y) \cdot (z \cdot x)), f(y), \delta\}$.

Example 3.9. Let $A = \{0, 1, 2, 3, 4\}$ be a set with a binary operation \cdot defined by the following Cayley table:

\cdot	0	1	2	3	4
0	0	1	2	3	4
1	0	0	2	0	4
2	0	1	0	0	0
3	0	1	2	0	4
4	0	1	2	3	0

Then $(A, \cdot, 0)$ is a UP-algebra. We defined a fuzzy set $f : A \rightarrow [0, 1]$ as follow:

$$f(0) = 0.3, f(1) = 0.4, f(2) = 0, f(3) = 0.1, \text{ and } f(4) = 0.$$

Then f is a fuzzy strongly UP-ideal with thresholds $\varepsilon = 0.5$ and $\delta = 0.8$ of A .

Theorem 3.10. *Every fuzzy strongly UP-ideal with thresholds ε and δ of A is a fuzzy UP-ideal with thresholds ε and δ .*

Proof. Assume that f is a fuzzy strongly UP-ideal with thresholds ε and δ of A . Then $\max\{f(0), \varepsilon\} \geq \min\{f(x), \delta\}$, for all $x \in A$. Let $x, y, z \in A$.

Case 1: $f(0) \leq \varepsilon$. Then $f(x) \leq \varepsilon$ for all $x \in A$. Thus

$$\max\{f(x \cdot z), \varepsilon\} = \varepsilon \geq f(y) \geq \min\{f(x \cdot (y \cdot z)), f(y), \delta\}.$$

Case 2: $f(0) > \varepsilon$. Then

$$\begin{aligned} \max\{f(x \cdot z), \varepsilon\} &\geq \min\{f((z \cdot y) \cdot (z \cdot (x \cdot z))), f(y), \delta\} \\ \text{(Proposition 2.4 (5))} &= \min\{f((z \cdot y) \cdot 0), f(y), \delta\} \\ \text{((UP-3))} &= \min\{f(0), f(y), \delta\}. \end{aligned}$$

If $\min\{f(0), f(y), \delta\} = f(y)$ or δ , then we obtain immediately that

$$\max\{f(x \cdot z), \varepsilon\} \geq \min\{f(x \cdot (y \cdot z)), f(y), \delta\}.$$

Assume that $\min\{f(0), f(y), \delta\} = f(0)$. Then

$$\begin{aligned} \min\{f(0), f(y), \delta\} &= f(0) = \max\{f(0), \varepsilon\} \\ &\geq \min\{f(y), \delta\} \\ &\geq \min\{f(x \cdot (y \cdot z)), f(y), \delta\}. \end{aligned}$$

Thus, f is a fuzzy UP-ideal with thresholds ε and δ of A . □

Example 3.11. Let $A = \{0, 1, 2, 3, 4\}$ be a set with a binary operation \cdot defined by the following Cayley table:

\cdot	0	1	2	3	4
0	0	1	2	3	4
1	0	0	2	0	4
2	0	1	0	0	0
3	0	1	2	0	4
4	0	1	2	3	0

Then $(A, \cdot, 0)$ is a UP-algebra. We defined a fuzzy set $f: A \rightarrow [0, 1]$ as follow:

$$f(0) = 1, f(1) = 0.2, f(2) = 0.1, f(3) = 0.2, \text{ and } f(4) = 0.9.$$

Then f is a fuzzy UP-ideal with thresholds $\varepsilon = 0.5$ and $\delta = 0.9$ of A . Since $\max\{f(2), \varepsilon\} = \max\{0.1, 0.5\} = 0.5 \not\geq 0.9 = \min\{1, 1, 0.9\} = \min\{f((2 \cdot 0) \cdot (2 \cdot 2)), f(0), \delta\}$, we have f is not a fuzzy strongly UP-ideal with thresholds $\varepsilon = 0.5$ and $\delta = 0.9$ of A .

Theorem 3.12. *Every fuzzy UP-ideal with thresholds ε and δ of A is a fuzzy UP-filter with thresholds ε and δ .*

Proof. Assume that f is a fuzzy UP-ideal with thresholds ε and δ of A . Then $\max\{f(0), \varepsilon\} \geq \min\{f(x), \delta\}$, for all $x \in A$. Let $x, y \in A$. Thus

$$\begin{aligned} ((\text{UP-2})) \quad \max\{f(y), \varepsilon\} &= \max\{f(0 \cdot y), \varepsilon\} \\ &\geq \min\{f(0 \cdot (x \cdot y)), f(x), \delta\} \\ ((\text{UP-2})) \quad &= \min\{f(x \cdot y), f(x), \delta\}. \end{aligned}$$

So, f is a fuzzy UP-filter with thresholds ε and δ of A . □

Example 3.13. Let $A = \{0, 1, 2, 3, 4\}$ be a set with a binary operation \cdot defined by the following Cayley table:

\cdot	0	1	2	3	4
0	0	1	2	3	4
1	0	0	2	3	4
2	0	0	0	3	3
3	0	1	2	0	3
4	0	1	2	0	0

Then $(A, \cdot, 0)$ is a UP-algebra. We defined a fuzzy set $f: A \rightarrow [0, 1]$ as follow:

$$f(0) = 0.9, f(1) = 0.8, f(2) = 0.7, f(3) = 0.5, \text{ and } f(4) = 0.5.$$

Then f is a fuzzy UP-filter with thresholds $\varepsilon = 0.2$ and $\delta = 0.9$ of A . Since $\max\{f(3 \cdot 4), \varepsilon\} = \max\{0.5, 0.2\} = 0.5 \not\geq 0.7 = \min\{0.9, 0.7, 0.9\} = \min\{f(3 \cdot (2 \cdot 4)), f(2), \delta\}$, we have f is not a fuzzy UP-ideal with thresholds $\varepsilon = 0.2$ and $\delta = 0.9$ of A .

Theorem 3.14. *Every fuzzy UP-filter with thresholds ε and δ of A is a fuzzy UP-subalgebra with thresholds ε and δ .*

Proof. Assume that f is a fuzzy UP-filter with thresholds ε and δ of A . Then $\max\{f(0), \varepsilon\} \geq \min\{f(x), \delta\}$, for all $x \in A$. Let $x, y \in A$. Then

Case 1: $f(0) \leq \varepsilon$. Then $f(x) \leq \varepsilon$, for all $x \in A$. Thus

$$\max\{f(x \cdot y), \varepsilon\} = \varepsilon \geq f(x) \geq \min\{f(x), f(y), \delta\}.$$

Case 2: $f(0) > \varepsilon$. Then

$$\begin{aligned} \max\{f(x \cdot y), \varepsilon\} &\geq \min\{f(y \cdot (x \cdot y)), f(y), \delta\} \\ (\text{Proposition 2.4 (5)}) \quad &= \min\{f(0), f(y), \delta\}. \end{aligned}$$

If $\min\{f(0), f(y), \delta\} = f(y)$ or δ , then we obtain immediately that

$$\max\{f(x \cdot y), \varepsilon\} \geq \min\{f(x), f(y), \delta\}.$$

Assume that $\min\{f(0), f(y), \delta\} = f(0)$. Then

$$\min\{f(0), f(y), \delta\} = f(0) = \max\{f(0), \varepsilon\} \geq \min\{f(y), \delta\} \geq \min\{f(x), f(y), \delta\}.$$

Thus, f is a fuzzy UP-subalgebra with thresholds ε and δ of A . □

Example 3.15. Let $A = \{0, 1, 2, 3, 4\}$ be a set with a binary operation \cdot defined by the following Cayley table:

\cdot	0	1	2	3	4
0	0	1	2	3	4
1	0	0	2	3	4
2	0	0	0	3	3
3	0	1	2	0	3
4	0	1	2	0	0

Then $(A, \cdot, 0)$ is a UP-algebra. We defined a fuzzy set $f: A \rightarrow [0, 1]$ as follow:

$$f(0) = 0.8, f(1) = 0.7, f(2) = 0.4, f(3) = 0.3, \text{ and } f(4) = 0.2.$$

Then f is a fuzzy UP-subalgebra with thresholds $\varepsilon = 0.2$ and $\delta = 0.9$ of A . Since $\max\{f(4), \varepsilon\} = \max\{0.2, 0.2\} = 0.2 \not\geq 0.3 = \min\{0.3, 0.3, 0.9\} = \min\{f(3 \cdot 4), f(3), \delta\}$, we have f is not a fuzzy UP-filter with thresholds $\varepsilon = 0.2$ and $\delta = 0.9$ of A .

By Definition 2.8, we see that a fuzzy UP-subalgebra (resp., fuzzy UP-filter, fuzzy UP-ideal, fuzzy strongly UP-ideal) is a fuzzy UP-subalgebra (resp., fuzzy UP-filter, fuzzy UP-ideal, fuzzy strongly UP-ideal) with thresholds 0 and 1. By Theorems 3.10, 3.12, and 3.14 and Examples 3.11, 3.13, and 3.15, we have that the notion of fuzzy UP-subalgebras with thresholds ε and δ is a generalization of fuzzy UP-filters with thresholds ε and δ , the notion of fuzzy UP-filters with thresholds ε and δ is a generalization of fuzzy UP-ideals with thresholds ε and δ , and the notion of fuzzy UP-ideals with thresholds ε and δ is a generalization of fuzzy strongly UP-ideals with thresholds ε and δ .

Theorem 3.16. *Let $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$. If a fuzzy set f in A is constant, then it is a fuzzy strongly UP-ideal with thresholds ε and δ of A .*

Proof. Assume that f is a constant fuzzy set in A and let $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$. Then $f(x) = f(0)$, for all $x \in A$. Let $x \in A$.

$$\max\{f(0), \varepsilon\} \geq f(0) = f(x) \geq \min\{f(x), \delta\}.$$

Let $x, y, z \in A$. Then

$$\begin{aligned} \max\{f(x), \varepsilon\} &\geq f(x) = \min\{f((z \cdot y) \cdot (z \cdot x)), f(y)\} \\ &\geq \min\{f((z \cdot y) \cdot (z \cdot x)), f(y), \delta\}. \end{aligned}$$

Thus, f is a fuzzy strongly UP-ideal with thresholds ε and δ of A . □

Theorem 3.17. *Let $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$. If a fuzzy set f in A is such that $f(x) \leq \varepsilon$, for all $x \in A$, then it is a fuzzy strongly UP-ideal (resp. fuzzy UP-ideal, fuzzy UP-filter and fuzzy UP-subalgebra) with thresholds ε and δ of A .*

Proof. (1) Let $x \in A$. Then

$$\max\{f(0), \varepsilon\} = \varepsilon \geq f(x) = \min\{f(x), \delta\}.$$

Let $x, y, z \in A$. Then

$$\max\{f(x), \varepsilon\} = \varepsilon \geq \min\{f((z \cdot y) \cdot (z \cdot x)), f(y)\} = \min\{f((z \cdot y) \cdot (z \cdot x)), f(y), \delta\}.$$

Thus, f is a fuzzy strongly UP-ideal with thresholds ε and δ of A .

(2) Let $x, y, z \in A$. Then

$$\max\{f(x \cdot z), \varepsilon\} = \varepsilon \geq \min\{f(x \cdot (y \cdot z)), f(y)\} = \min\{f(x \cdot (y \cdot z)), f(y), \delta\}.$$

Thus, f is a fuzzy UP-ideal with thresholds ε and δ of A .

(3) Let $x, y \in A$. Then

$$\max\{f(y), \varepsilon\} = \varepsilon \geq \min\{f(x \cdot y), f(x)\} = \min\{f(x \cdot y), f(x), \delta\}.$$

Thus, f is a fuzzy UP-filter with thresholds ε and δ of A .

(4) Let $x, y \in A$. Then

$$\max\{f(x \cdot y), \varepsilon\} = \varepsilon \geq \min\{f(x), f(y)\} = \min\{f(x), f(y), \delta\}.$$

Thus, f is a fuzzy UP-subalgebra with thresholds ε and δ of A . □

Theorem 3.18. *Let $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$. If a fuzzy set f in A with $\varepsilon \leq f(x) \leq \delta$ is a fuzzy strongly UP-ideal with thresholds ε and δ of A , then it is constant.*

Proof. For all $x \in A$,

$$f(0) = \max\{f(0), \varepsilon\} \geq \min\{f(x), \delta\} = f(x)$$

and

$$\begin{aligned} f(x) &= \max\{f(x), \varepsilon\} \geq \min\{f((x \cdot 0) \cdot (x \cdot x)), f(0), \delta\} \\ \text{(Proposition 2.4 (1))} &= \min\{f((x \cdot 0) \cdot 0), f(0), \delta\} \\ \text{((UP-3))} &= \min\{f(0), f(0), \delta\} \\ &= \min\{f(0), \delta\} \\ &= f(0). \end{aligned}$$

Then, $f(x) = f(0)$, for all $x \in A$. Thus f is constant. □

Theorem 3.19. *Let $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$. If a fuzzy set f in A is such that $f(x) \geq \delta$ for all $x \in A$, then it is a fuzzy strongly UP-ideal (resp. fuzzy UP-ideal, fuzzy UP-filter and fuzzy UP-subalgebra) with thresholds ε and δ of A .*

Proof. (1) Let $x \in A$. Then

$$\max\{f(0), \varepsilon\} = f(0) \geq \delta = \min\{f(x), \delta\}.$$

Let $x, y, z \in A$. Then

$$\max\{f(x), \varepsilon\} = f(x) \geq \delta = \min\{f((z \cdot y) \cdot (z \cdot x)), f(y), \delta\}.$$

Thus, f is a fuzzy strongly UP-ideal with thresholds ε and δ of A .

(2) Let $x, y, z \in A$. Then

$$\max\{f(x \cdot z), \varepsilon\} = f(x \cdot z) \geq \delta = \min\{f(x \cdot (y \cdot z)), f(y), \delta\}.$$

Thus, f is a fuzzy UP-ideal with thresholds ε and δ of A .

(3) Let $x, y \in A$. Then

$$\max\{f(y), \varepsilon\} = f(y) \geq \delta = \min\{f(x \cdot y), f(x), \delta\}.$$

Thus, f is a fuzzy UP-filter with thresholds ε and δ of A .

(4) Let $x, y \in A$. Then

$$\max\{f(x \cdot y), \varepsilon\} = f(x \cdot y) \geq \delta = \min\{f(x), f(y), \delta\}.$$

Thus, f is a fuzzy UP-subalgebra with thresholds ε and δ of A . □

Corollary 3.20. *Let $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$. Then a fuzzy set f in A with $\varepsilon \leq f(x) \leq \delta$ is a fuzzy strongly UP-ideal with thresholds ε and δ of A if and only if it is constant.*

Proof. It is straightforward by Theorem 3.16 and 3.18. □

4. UPPER t -LEVEL SUBSETS OF A FUZZY SET

In this section, we discuss the relations between fuzzy UP-subalgebras (resp., fuzzy UP-filters, fuzzy UP-ideals, fuzzy strongly UP-ideals) with thresholds and their level subsets.

Let f be a fuzzy set in A . For any $t \in [0, 1]$, the set

$$U(f, t) = \{x \in A \mid f(x) \geq t\}$$

is called an upper t -level subset [15] of f .

Theorem 4.1. *Let f be a fuzzy set in A and $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$. Then f is a fuzzy UP-subalgebra with thresholds ε and δ of A if and only if for all $t \in (\varepsilon, \delta]$, $U(f, t)$ is a UP-subalgebra of A , if $U(f, t)$ is nonempty.*

Proof. Assume that f is a fuzzy UP-subalgebra with thresholds ε and δ of A . Let $t \in (\varepsilon, \delta]$ be such that $U(f, t) \neq \emptyset$ and let $x, y \in U(f, t)$. Then $f(x) \geq t, f(y) \geq t$, and $\delta \geq t$. Thus t is a lower bound of $\{f(x), f(y), \delta\}$. Since f is a fuzzy UP-subalgebra with thresholds ε and δ of A , we have

$$\max\{f(x \cdot y), \varepsilon\} \geq \min\{f(x), f(y), \delta\} \geq t > \varepsilon.$$

So $\max\{f(x \cdot y), \varepsilon\} = f(x \cdot y)$. Since $\max\{f(x \cdot y), \varepsilon\} \geq t$, we get $f(x \cdot y) \geq t$. Hence $x \cdot y \in U(f, t)$. Therefore, $U(f, t)$ is a UP-subalgebra of A .

Conversely, assume that for all $t \in (\varepsilon, \delta]$, $U(f, t)$ is a UP-subalgebra of A , if $U(f, t)$ is nonempty. Let $x, y \in A$. Then $f(x), f(y) \in [0, 1]$. Choose $t = \min\{f(x), f(y)\}$. Then $f(x) \geq t$ and $f(y) \geq t$. Thus $x, y \in U(f, t) \neq \emptyset$. By assumption, we have $U(f, t)$ is a UP-subalgebra of A . So $x \cdot y \in U(f, t)$, that is, $f(x \cdot y) \geq t = \min\{f(x), f(y)\}$. So

$$\max\{f(x \cdot y), \varepsilon\} \geq f(x \cdot y) \geq \min\{f(x), f(y)\} \geq \min\{f(x), f(y), \delta\}.$$

Hence, f is a UP-subalgebra with thresholds ε and δ of A . □

Theorem 4.2. *Let f be a fuzzy set in A and $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$. Then f is a fuzzy UP-filter with thresholds ε and δ of A if and only if for all $t \in (\varepsilon, \delta]$, $U(f, t)$ is a UP-filter of A , if $U(f, t)$ is nonempty.*

Proof. Assume that f is a fuzzy UP-filter with thresholds ε and δ of A . Let $t \in (\varepsilon, \delta]$ be such that $U(f, t) \neq \emptyset$ and let $a \in U(f, t)$. Then $f(a) \geq t$ and $\delta \geq t$. Thus t is a lower bound of $\{f(a), \delta\}$. Since f is a fuzzy UP-filter with thresholds ε and δ of A , we have

$$\max\{f(0), \varepsilon\} \geq \min\{f(a), \delta\} \geq t > \varepsilon.$$

So $\max\{f(0), \varepsilon\} = f(0)$. Since $\max\{f(0), \varepsilon\} \geq t$, we get $f(0) \geq t$. Hence $0 \in U(f, t)$.

Next, let $x, y \in A$ be such that $x \cdot y \in U(f, t)$ and $x \in U(f, t)$. Then $f(x \cdot y) \geq t$, $f(x) \geq t$, and $\delta \geq t$. Thus t is a lower bound of $\{f(x \cdot y), f(x), \delta\}$. Since f is a fuzzy UP-filter with thresholds ε and δ of A , we have

$$\max\{f(y), \varepsilon\} \geq \min\{f(x \cdot y), f(x), \delta\} \geq t > \varepsilon.$$

So $\max\{f(y), \varepsilon\} = f(y)$. Since $\max\{f(y), \varepsilon\} \geq t$, we get $f(y) \geq t$. Hence $y \in U(f, t)$. Therefore, $U(f, t)$ is a UP-filter of A .

Conversely, assume that for all $t \in (\varepsilon, \delta]$, $U(f, t)$ is a UP-filter of A , if $U(f, t)$ is nonempty. Let $x \in A$. Then $f(x) \in [0, 1]$. Choose $t = f(x)$. Then $f(x) \geq t$. Thus $x \in U(f, t) \neq \emptyset$. By assumption, we have $U(f, t)$ is a UP-filter of A . So $0 \in U(f, t)$, that is, $f(0) \geq t = f(x)$, Hence

$$\max\{f(0), \varepsilon\} \geq f(0) \geq f(x) \geq \min\{f(x), \delta\}.$$

Next, let $x, y \in A$. Then $f(x \cdot y), f(x) \in [0, 1]$. Choose $t = \min\{f(x \cdot y), f(x)\}$. Then $f(x \cdot y) \geq t$ and $f(x) \geq t$. Thus $x \cdot y, x \in U(f, t) \neq \emptyset$. By assumption, we have $U(f, t)$ is a UP-filter of A . So $y \in U(f, t)$, that is, $f(y) \geq t = \min\{f(x \cdot y), f(x)\}$. Hence

$$\max\{f(y), \varepsilon\} \geq f(y) \geq \min\{f(x \cdot y), f(x)\} \geq \min\{f(x \cdot y), f(x), \delta\}.$$

f is a fuzzy UP-filter with thresholds ε and δ of A . □

Theorem 4.3. *Let f be a fuzzy set in A and $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$. Then f is a fuzzy UP-ideal with thresholds ε and δ of A if and only if for all $t \in (\varepsilon, \delta]$, $U(f, t)$ is a UP-ideal of A , if $U(f, t)$ is nonempty.*

Proof. Assume that f is a fuzzy UP-ideal with thresholds ε and δ of A . Let $t \in (\varepsilon, \delta]$ be such that $U(f, t) \neq \emptyset$ and let $a \in U(f, t)$. Then $f(a) \geq t$ and $\delta \geq t$. Thus t is a lower bound of $\{f(a), \delta\}$. Since f is a fuzzy UP-ideal with thresholds ε and δ of A , we have

$$\max\{f(0), \varepsilon\} \geq \min\{f(a), \delta\} \geq t > \varepsilon.$$

So $\max\{f(0), \varepsilon\} = f(0)$. Since $\max\{f(0), \varepsilon\} \geq t$, we get $f(0) \geq t$. Hence $0 \in U(f, t)$.

Next, let $x, y, z \in A$ be such that $x \cdot (y \cdot z) \in U(f, t)$ and $y \in U(f, t)$. Then $f(x \cdot (y \cdot z)) \geq t$, $f(y) \geq t$, and $\delta \geq t$. Thus t is a lower bound of $\{f(x \cdot (y \cdot z)), f(y), \delta\}$. Since f is a fuzzy UP-ideal with thresholds ε and δ of A , we have

$$\max\{f(x \cdot z), \varepsilon\} \geq \min\{f(x \cdot (y \cdot z)), f(y), \delta\} \geq t > \varepsilon.$$

So $\max\{f(x \cdot z), \varepsilon\} = f(x \cdot z)$. Since $\max\{f(x \cdot z), \varepsilon\} \geq t$, we get $f(x \cdot z) \geq t$. Hence $x \cdot z \in U(f, t)$. Therefore, $U(f, t)$ is a UP-ideal of A .

Conversely, assume that for all $t \in (\varepsilon, \delta]$, $U(f, t)$ is a UP-ideal of A , if $U(f, t)$ is nonempty. Let $x \in A$. Then $f(x) \in [0, 1]$. Choose $t = f(x)$. Then $f(x) \geq t$. Thus $x \in U(f, t) \neq \emptyset$. By assumption, we have $U(f, t)$ is a UP-ideal of A . So $0 \in U(f, t)$, that is, $f(0) \geq t = f(x)$, Hence

$$\max\{f(0), \varepsilon\} \geq f(0) \geq f(x) \geq \min\{f(x), \delta\}.$$

Next, let $x, y, z \in A$. Then $f(x \cdot (y \cdot z)), f(y) \in [0, 1]$. Choose $t = \min\{f(x \cdot (y \cdot z)), f(y)\}$. Then $f(x \cdot (y \cdot z)) \geq t$ and $f(y) \geq t$. Thus $x \cdot (y \cdot z), y \in U(f, t) \neq \emptyset$.

By assumption, we have $U(f, t)$ is a UP-ideal of A . Thus $x \cdot z \in U(f, t)$, that is, $f(x \cdot z) \geq t = \min\{f(x \cdot (y \cdot z)), f(y)\}$. So

$$\max\{f(x \cdot z), \varepsilon\} \geq f(x \cdot z) \geq \min\{f(x \cdot (y \cdot z)), f(y)\} \geq \min\{f(x \cdot (y \cdot z)), f(y), \delta\}.$$

Hence, f is a fuzzy UP-ideal with thresholds ε and δ of A . \square

Theorem 4.4. *Let f be a fuzzy set in A and $\varepsilon, \delta \in [0, 1]$ with $\varepsilon < \delta$. Then f is a fuzzy strongly UP-ideal with thresholds ε and δ of A if and only if for all $t \in (\varepsilon, \delta]$, $U(f, t)$ is a strongly UP-ideal of A , if $U(f, t)$ is nonempty.*

Proof. Assume that f is a fuzzy strongly UP-ideal with thresholds ε and δ of A . Let $t \in (\varepsilon, \delta]$ be such that $U(f, t) \neq \emptyset$ and let $a \in U(f, t)$. Then $f(a) \geq t$ and $\delta \geq t$. Thus t is a lower bound of $\{f(a), \delta\}$. Since f is a fuzzy strongly UP-ideal with thresholds ε and δ of A , we have

$$\max\{f(0), \varepsilon\} \geq \min\{f(a), \delta\} \geq t > \varepsilon.$$

So $\max\{f(0), \varepsilon\} = f(0)$. Since $\max\{f(0), \varepsilon\} \geq t$, we get $f(0) \geq t$. Hence $0 \in U(f, t)$.

Next, let $x, y, z \in A$ be such that $(z \cdot y) \cdot (z \cdot x) \in U(f, t)$ and $y \in U(f, t)$. Then $f((z \cdot y) \cdot (z \cdot x)) \geq t, f(y) \geq t$, and $\delta \geq t$. Thus t is a lower bound of $\{f((z \cdot y) \cdot (z \cdot x)), f(y), \delta\}$. Since f is a fuzzy strongly UP-ideal with thresholds ε and δ of A , we have

$$\max\{f(x), \varepsilon\} \geq \min\{f((z \cdot y) \cdot (z \cdot x)), f(y), \delta\} \geq t > \varepsilon.$$

So $\max\{f(x), \varepsilon\} = f(x)$. Since $\max\{f(x), \varepsilon\} \geq t$, we get $f(x) \geq t$. Hence $x \in U(f, t)$. Therefore, $U(f, t)$ is a strongly UP-ideal of A .

Conversely, assume that for all $t \in (\varepsilon, \delta]$, $U(f, t)$ is a strongly UP-ideal of A , if $U(f, t)$ is nonempty. Let $x \in A$. Then $f(x) \in [0, 1]$. Choose $t = f(x)$. Then $f(x) \geq t$. Thus $x \in U(f, t) \neq \emptyset$. By assumption, we have $U(f, t)$ is a strongly UP-ideal of A . So $0 \in U(f, t)$, that is, $f(0) \geq t = f(x)$. Hence

$$\max\{f(0), \varepsilon\} \geq f(0) \geq f(x) \geq \min\{f(x), \delta\}.$$

Next, let $x, y, z \in A$. Then $f((z \cdot y) \cdot (z \cdot x)), f(y) \in [0, 1]$. Choose $t = \min\{f((z \cdot y) \cdot (z \cdot x)), f(y)\}$. Then $f((z \cdot y) \cdot (z \cdot x)) \geq t$ and $f(y) \geq t$. Thus $(z \cdot y) \cdot (z \cdot x), y \in U(f, t) \neq \emptyset$. By assumption, we have $U(f, t)$ is a strongly UP-ideal of A . So $x \in U(f, t)$, that is, $f(x) \geq t = \min\{f((z \cdot y) \cdot (z \cdot x)), f(y)\}$. Hence

$$\begin{aligned} \max\{f(x), \varepsilon\} &\geq f(x) \geq \min\{f((z \cdot y) \cdot (z \cdot x)), f(y)\} \\ &\geq \min\{f((z \cdot y) \cdot (z \cdot x)), f(y), \delta\}. \end{aligned}$$

Therefore, f is a fuzzy strongly UP-ideal with thresholds ε and δ of A . \square

5. IMAGE AND PREIMAGE OF A FUZZY SET

Definition 5.1 ([4]). Let f be a function from a nonempty set X to a nonempty set Y . If μ is a fuzzy set in X , then the fuzzy set β in Y defined by

$$\beta(y) = \begin{cases} \sup\{\mu(t)\}_{t \in f^{-1}(y)} & \text{if } f^{-1}(y) \neq \emptyset, \\ 0 & \text{otherwise} \end{cases}$$

is said to be the image of μ under f .

Similarly, if β is a fuzzy set in Y , then the fuzzy set $\mu = \beta \circ f$ in X (i.e., the fuzzy set defined by $\mu(x) = \beta(f(x))$ for all $x \in X$) is called the preimage of β under f .

Definition 5.2 ([10]). A fuzzy set f in A has sup property, if for any nonempty subset T of A , there exists $t_0 \in T$ such that $f(t_0) = \sup\{f(t)\}_{t \in T}$.

Lemma 5.3 ([9]). Let $(A, \cdot, 0_A)$ and $(B, *, 0_B)$ be UP-algebras and let $f: A \rightarrow B$ be a surjective UP-homomorphism. Let μ be an f -invariant fuzzy set in A with sup property. For any $a, b \in B$, there exist $a_0 \in f^{-1}(a)$ and $b_0 \in f^{-1}(b)$ such that $\beta(a) = \mu(a_0)$, $\beta(b) = \mu(b_0)$, and $\beta(a * b) = \mu(a_0 \cdot b_0)$.

Theorem 5.4. Let $(A, \cdot, 0_A)$ and $(B, *, 0_B)$ be UP-algebras and let $f: A \rightarrow B$ be a surjective UP-homomorphism. Then the following statements hold:

- (1) if μ is an f -invariant fuzzy UP-subalgebra with thresholds ε and δ of A with sup property, then β is a fuzzy UP-subalgebra with thresholds ε and δ of B ,
- (2) if μ is an f -invariant fuzzy UP-filter with thresholds ε and δ of A with sup property, then β is a fuzzy UP-filter with thresholds ε and δ of B ,
- (3) if μ is an f -invariant fuzzy UP-ideal with thresholds ε and δ of A with sup property, then β is a fuzzy UP-ideal with thresholds ε and δ of B , and
- (4) if μ is an f -invariant fuzzy strongly UP-ideal with thresholds ε and δ of A with sup property, then β is a fuzzy strongly UP-ideal with thresholds ε and δ of B .

Proof. (1) Assume that μ is an f -invariant fuzzy UP-subalgebra with thresholds ε and δ of A with sup property. Let $a, b \in B$. Then by Lemma 5.3, there exist $a_0 \in f^{-1}(a)$ and $b_0 \in f^{-1}(b)$ such that $\beta(a) = \mu(a_0)$, $\beta(b) = \mu(b_0)$, and $\beta(a * b) = \mu(a_0 \cdot b_0)$. Thus

$$\max\{\beta(a * b), \varepsilon\} = \max\{\mu(a_0 \cdot b_0), \varepsilon\} \geq \min\{\mu(a_0), \mu(b_0), \delta\} = \min\{\beta(a), \beta(b), \delta\}.$$

So, β is a fuzzy UP-subalgebra with thresholds ε and δ of B .

(2) Assume that μ is an f -invariant fuzzy UP-filter with thresholds ε and δ of A with sup property. Since $f(0_A) = 0_B$, we have $f^{-1}(0_B) \neq \emptyset$. Then there exists $x_1 \in f^{-1}(0_B)$ such that $\mu(x_1) = \beta(0_B)$. Thus $f(x_1) = 0_B = f(0_A)$. Since μ is f -invariant, $\mu(x_1) = \mu(0_A)$. So, $\mu(0_A) = \beta(0_B)$.

Let $y \in B$. Since f is surjective, we have $f^{-1}(y) \neq \emptyset$. Then there exists $x \in f^{-1}(y)$ such that $\mu(x) = \beta(y)$. Thus

$$\max\{\beta(0_B), \varepsilon\} = \max\{\mu(0_A), \varepsilon\} \geq \min\{\mu(x), \delta\} = \min\{\beta(y), \delta\}.$$

Next, let $a, b \in B$. Then by Lemma 5.3, there exist $a_0 \in f^{-1}(a)$ and $b_0 \in f^{-1}(b)$ such that $\mu(a_0) = \beta(a)$, $\mu(b_0) = \beta(b)$, and $\mu(a_0 \cdot b_0) = \beta(a * b)$. Thus

$$\max\{\beta(b), \varepsilon\} = \max\{\mu(b_0), \varepsilon\} \geq \min\{\mu(a_0 \cdot b_0), \mu(a_0), \delta\} = \min\{\beta(a * b), \beta(a), \delta\}.$$

So, β is a fuzzy UP-filter with thresholds ε and δ of B .

(3) Assume that μ is an f -invariant fuzzy UP-ideal with thresholds ε and δ of A with sup property. Since $f(0_A) = 0_B$, we have $f^{-1}(0_B) \neq \emptyset$. Then there exists $x_1 \in f^{-1}(0_B)$ such that $\mu(x_1) = \beta(0_B)$. Thus $f(x_1) = 0_B = f(0_A)$. So $\mu(x_1) = \mu(0_A)$, because μ is f -invariant. Hence, $\mu(0_A) = \beta(0_B)$.

Let $y \in B$. Since f is surjective, we have $f^{-1}(y) \neq \emptyset$. Then there exists $x \in f^{-1}(y)$ such that $\mu(x) = \beta(y)$. Thus

$$\max\{\beta(0_B), \varepsilon\} = \max\{\mu(0_A), \varepsilon\} \geq \min\{\mu(x), \delta\} = \min\{\beta(y), \delta\}.$$

Next, let $a, b, c \in B$. Then by Lemma 5.3, there exist $a_0 \in f^{-1}(a), b_0 \in f^{-1}(b)$ and $c_0 \in f^{-1}(c)$ such that $\beta(b) = \mu(b_0), \beta(a * c) = \mu(a_0 \cdot c_0)$ and $\beta(a * (b * c)) = \mu(a_0 \cdot (b_0 \cdot c_0))$. Thus

$$\begin{aligned} \max\{\beta(a * c), \varepsilon\} &= \max\{\mu(a_0 \cdot c_0), \varepsilon\} \geq \min\{\mu(a_0 \cdot (b_0 \cdot c_0)), \mu(b_0), \delta\} \\ &= \min\{\beta(a * (b * c)), \beta(b), \delta\}. \end{aligned}$$

So, β is a fuzzy UP-ideal with thresholds ε and δ of B .

(4) Assume that μ is an f -invariant fuzzy strongly UP-ideal with thresholds ε and δ of A with sup property. Since $f(0_A) = 0_B$, we have $f^{-1}(0_B) \neq \emptyset$. Then there exists $x_1 \in f^{-1}(0_B)$ such that $\mu(x_1) = \beta(0_B)$. Thus $f(x_1) = 0_B = f(0_A)$. So $\mu(x_1) = \mu(0_A)$, because μ is f -invariant. Hence, $\mu(0_A) = \beta(0_B)$.

Let $y \in B$. Since f is surjective, we have $f^{-1}(y) \neq \emptyset$. Then there exists $x \in f^{-1}(y)$ such that $\mu(x) = \beta(y)$. Thus

$$\max\{\beta(0_B), \varepsilon\} = \max\{\mu(0_A), \varepsilon\} \geq \min\{\mu(x), \delta\} = \min\{\beta(y), \delta\}.$$

Next, let $a, b, c \in B$. Then by Lemma 5.3, there exist $a_0 \in f^{-1}(a), b_0 \in f^{-1}(b)$ and $c_0 \in f^{-1}(c)$ such that $\mu(a_0) = \beta(a), \mu(b_0) = \beta(b)$ and $\mu((c_0 \cdot b_0) \cdot (c_0 \cdot a_0)) = \beta((c \cdot b) \cdot (c \cdot a))$. Thus

$$\begin{aligned} \max\{\beta(a), \varepsilon\} &= \max\{\mu(a_0), \varepsilon\} \geq \min\{\mu((c_0 \cdot b_0) \cdot (c_0 \cdot a_0)), \mu(b_0), \delta\} \\ &= \min\{\beta((c \cdot b) \cdot (c \cdot a)), \beta(b), \delta\}. \end{aligned}$$

So, β is a fuzzy strongly UP-ideal with thresholds ε and δ of B . □

Theorem 5.5. *Let $(A, \cdot, 0_A)$ and $(B, *, 0_B)$ be UP-algebras and let $f: A \rightarrow B$ be a UP-homomorphism. Then the following statements hold:*

- (1) *if β is a fuzzy UP-subalgebra with thresholds ε and δ of B , then μ is a fuzzy UP-subalgebra with thresholds ε and δ of A ,*
- (2) *if β is a fuzzy UP-filter with thresholds ε and δ of B , then μ is a fuzzy UP-filter with thresholds ε and δ of A ,*
- (3) *if β is a fuzzy UP-ideal with thresholds ε and δ of B , then μ is a fuzzy UP-ideal with thresholds ε and δ of A , and*
- (4) *if β is a fuzzy strongly UP-ideal with thresholds ε and δ of B , then μ is a fuzzy strongly UP-ideal with thresholds ε and δ of A .*

Proof. (1) Assume that β is a fuzzy UP-subalgebra with thresholds ε and δ of B . Let $x, y \in A$. Then

$$\begin{aligned} \max\{\mu(x \cdot y), \varepsilon\} &= \max\{(\beta \circ f)(x \cdot y), \varepsilon\} = \max\{\beta(f(x \cdot y)), \varepsilon\} \\ &= \max\{\beta(f(x) * f(y)), \varepsilon\} \\ &\geq \min\{\beta(f(x)), \beta(f(y)), \delta\} \\ &= \min\{(\beta \circ f)(x), (\beta \circ f)(y), \delta\} \\ &= \min\{\mu(x), \mu(y), \delta\}. \end{aligned}$$

Thus, μ is a fuzzy UP-subalgebra with thresholds ε and δ of A .

(2) Assume that β is a fuzzy UP-filter with thresholds ε and δ of B . Let $x \in A$. Then

$$\begin{aligned} \max\{\mu(0_A), \varepsilon\} &= \max\{(\beta \circ f)(0_A), \varepsilon\} = \max\{\beta(f(0_A)), \varepsilon\} \\ (f(0_A) = 0_B) \qquad &= \max\{\beta(0_B), \varepsilon\} \\ &\geq \min\{\beta(f(x)), \delta\} \\ &= \min\{(\beta \circ f)(x), \delta\} \\ &= \min\{\mu(x), \delta\}. \end{aligned}$$

Let $x, y \in A$. Then

$$\begin{aligned} \max\{\mu(y), \varepsilon\} &= \max\{(\beta \circ f)(y), \varepsilon\} = \max\{\beta(f(y)), \varepsilon\} \\ &\geq \min\{\beta(f(x) * f(y)), \beta(f(x)), \delta\} \\ &= \min\{\beta(f(x \cdot y)), \beta(f(x)), \delta\} \\ &= \min\{(\beta \circ f)(x \cdot y), (\beta \circ f)(x), \delta\} \\ &= \min\{\mu(x \cdot y), \mu(x), \delta\}. \end{aligned}$$

Thus, μ is a fuzzy UP-filter with thresholds ε and δ of A .

(3) Assume that β is a fuzzy UP-ideal with thresholds ε and δ of B . Let $x \in A$. Then

$$\begin{aligned} \max\{\mu(0_A), \varepsilon\} &= \max\{(\beta \circ f)(0_A), \varepsilon\} = \max\{\beta(f(0_A)), \varepsilon\} \\ (f(0_A) = 0_B) \qquad &= \max\{\beta(0_B), \varepsilon\} \\ &\geq \min\{\beta(f(x)), \delta\} \\ &= \min\{(\beta \circ f)(x), \delta\} \\ &= \min\{\mu(x), \delta\}. \end{aligned}$$

Let $x, y, z \in A$. Then

$$\begin{aligned} \max\{\mu(x \cdot z), \varepsilon\} &= \max\{(\beta \circ f)(x \cdot z), \varepsilon\} = \max\{\beta(f(x \cdot z)), \varepsilon\} \\ &= \min\{\beta(f(x) * f(z)), \delta\} \\ &\geq \min\{\beta(f(x) * (f(y) * f(z))), \beta(f(y)), \delta\} \\ &= \min\{\beta(f(x) * f(y \cdot z)), \beta(f(y)), \delta\} \\ &= \min\{\beta(f(x \cdot (y \cdot z))), \beta(f(y)), \delta\} \\ &= \min\{(\beta \circ f)(x \cdot (y \cdot z)), (\beta \circ f)(y), \delta\} \\ &= \min\{\mu(x \cdot (y \cdot z)), \mu(y), \delta\}. \end{aligned}$$

Thus, μ is a fuzzy UP-ideal with thresholds ε and δ of A .

(4) Assume that β is a fuzzy strongly UP-ideal with thresholds ε and δ of B . Let $x \in A$. Then

$$\begin{aligned} \max\{\mu(0_A), \varepsilon\} &= \max\{(\beta \circ f)(0_A), \varepsilon\} = \max\{\beta(f(0_A)), \varepsilon\} \\ (f(0_A) = 0_B) &= \max\{\beta(0_B), \varepsilon\} \\ &\geq \min\{\beta(f(x)), \delta\} \\ &= \min\{(\beta \circ f)(x), \delta\} \\ &= \min\{\mu(x), \delta\}. \end{aligned}$$

Let $x, y, z \in A$. Then

$$\begin{aligned} \max\{\mu(x), \varepsilon\} &= \max\{(\beta \circ f)(x), \varepsilon\} \\ &= \max\{\beta(f(x)), \varepsilon\} \\ &\geq \min\{\beta((f(z) * f(y)) * (f(z) * f(x))), \beta(f(y)), \delta\} \\ &= \min\{\beta(f(z \cdot y) * f(z \cdot x)), \beta(f(y)), \delta\} \\ &= \min\{\beta(f((z \cdot y) \cdot (z \cdot x))), \beta(f(y)), \delta\} \\ &= \min\{(\beta \circ f)((z \cdot y) \cdot (z \cdot x)), (\beta \circ f)(y), \delta\} \\ &= \min\{\mu((z \cdot y) \cdot (z \cdot x)), \mu(y), \delta\}. \end{aligned}$$

Thus, μ is a fuzzy strongly UP-ideal with thresholds ε and δ of A . □

Definition 5.6. Let f be a function from a nonempty set X to a nonempty set Y . If μ is a fuzzy set in X , then the fuzzy set β in Y defined by

$$\beta(y) = \begin{cases} \inf\{\mu(t)\}_{t \in f^{-1}(y)} & \text{if } f^{-1}(y) \neq \emptyset, \\ 1 & \text{otherwise} \end{cases}$$

is said to be the image of μ under f .

Similarly, if β is a fuzzy set in Y , then the fuzzy set $\mu = \beta \circ f$ in X (i.e., the fuzzy set defined by $\mu(x) = \beta(f(x))$ for all $x \in X$) is called the preimage of β under f .

Definition 5.7 ([10]). A fuzzy set f in A has inf property, if for any nonempty subset T of A , there exists $t_0 \in T$ such that $f(t_0) = \inf\{f(t)\}_{t \in T}$.

Lemma 5.8 ([7]). Let $(A, \cdot, 0_A)$ and $(B, *, 0_B)$ be UP-algebras and let $f: A \rightarrow B$ be a surjective UP-homomorphism. Let μ be an f -invariant fuzzy set in A with inf property. For any $a, b \in B$, there exist $a_0 \in f^{-1}(a)$ and $b_0 \in f^{-1}(b)$ such that $\beta(a) = \mu(a_0)$, $\beta(b) = \mu(b_0)$, and $\beta(a * b) = \mu(a_0 \cdot b_0)$.

Theorem 5.9. Let $(A, \cdot, 0_A)$ and $(B, *, 0_B)$ be UP-algebras and let $f: A \rightarrow B$ be a surjective UP-homomorphism. Then the following statements hold:

- (1) if μ is an f -invariant fuzzy UP-subalgebra with thresholds ε and δ of A with inf property, then β is a fuzzy UP-subalgebra with thresholds ε and δ of B ,
- (2) if μ is an f -invariant fuzzy UP-filter with thresholds ε and δ of A with inf property, then β is a fuzzy UP-filter with thresholds ε and δ of B ,
- (3) if μ is an f -invariant fuzzy UP-ideal with thresholds ε and δ of A with inf property, then β is a fuzzy UP-ideal with thresholds ε and δ of B , and

(4) if μ is an f -invariant fuzzy strongly UP-ideal with thresholds ε and δ of A with inf property, then β is a fuzzy strongly UP-ideal with thresholds ε and δ of B .

Proof. (1) Assume that μ is an f -invariant fuzzy UP-subalgebra with thresholds ε and δ of A with inf property. Let $a, b \in B$. Then by Lemma 5.8, there exist $a_0 \in f^{-1}(a)$ and $b_0 \in f^{-1}(b)$ such that $\beta(a) = \mu(a_0), \beta(b) = \mu(b_0)$, and $\beta(a * b) = \mu(a_0 \cdot b_0)$. Thus

$$\max\{\beta(a * b), \varepsilon\} = \max\{\mu(a_0 \cdot b_0), \varepsilon\} \geq \min\{\mu(a_0), \mu(b_0), \delta\} = \min\{\beta(a), \beta(b), \delta\}.$$

So, β is a fuzzy UP-subalgebra with thresholds ε and δ of B .

(2) Assume that μ is an f -invariant fuzzy UP-filter with thresholds ε and δ of A with inf property. Since $f(0_A) = 0_B$, we have $f^{-1}(0_B) \neq \emptyset$. Then there exists $x_1 \in f^{-1}(0_B)$ such that $\mu(x_1) = \beta(0_B)$. Thus $f(x_1) = 0_B = f(0_A)$. So $\mu(x_1) = \mu(0_A)$, because μ is f -invariant. Hence, $\mu(0_A) = \beta(0_B)$.

Let $y \in B$. Since f is surjective, we have $f^{-1}(y) \neq \emptyset$. Then there exists $x \in f^{-1}(y)$ such that $\mu(x) = \beta(y)$. Thus

$$\max\{\beta(0_B), \varepsilon\} = \max\{\mu(0_A), \varepsilon\} \geq \min\{\mu(x), \delta\} = \min\{\beta(y), \delta\}.$$

Next, let $a, b \in B$. Then by Lemma 5.8, there exist $a_0 \in f^{-1}(a)$ and $b_0 \in f^{-1}(b)$ such that $\beta(a) = \mu(a_0), \beta(b) = \mu(b_0)$, and $\beta(a * b) = \mu(a_0 \cdot b_0)$. Thus

$$\max\{\beta(b), \varepsilon\} = \max\{\mu(b_0), \varepsilon\} \geq \min\{\mu(a_0 \cdot b_0), \mu(a_0), \delta\} = \min\{\beta(a * b), \beta(a), \delta\}.$$

So, β is a fuzzy UP-filter with thresholds ε and δ of B .

(3) Assume that μ is an f -invariant fuzzy UP-ideal with thresholds ε and δ of A with inf property. Since $f(0_A) = 0_B$, we have $f^{-1}(0_B) \neq \emptyset$. Then there exists $x_1 \in f^{-1}(0_B)$ such that $\mu(x_1) = \beta(0_B)$. Thus $f(x_1) = 0_B = f(0_A)$. So $\mu(x_1) = \mu(0_A)$, because μ is f -invariant. Hence, $\mu(0_A) = \beta(0_B)$.

Let $y \in B$. Since f is surjective, we have $f^{-1}(y) \neq \emptyset$. Then there exists $x \in f^{-1}(y)$ such that $\mu(x) = \beta(y)$. Thus

$$\max\{\beta(0_B), \varepsilon\} = \max\{\mu(0_A), \varepsilon\} \geq \min\{\mu(x), \delta\} = \min\{\beta(y), \delta\}.$$

Next, let $a, b, c \in B$. Then by Lemma 5.8, there exist $a_0 \in f^{-1}(a), b_0 \in f^{-1}(b)$ and $c_0 \in f^{-1}(c)$ such that $\beta(b) = \mu(b_0), \beta(a * c) = \mu(a_0 \cdot c_0)$ and $\beta(a * (b * c)) = \mu(a_0 \cdot (b_0 \cdot c_0))$. Thus

$$\begin{aligned} \max\{\beta(a * c), \varepsilon\} &= \max\{\mu(a_0 \cdot c_0), \varepsilon\} \geq \min\{\mu(a_0 \cdot (b_0 \cdot c_0)), \mu(b_0), \delta\} \\ &= \min\{\beta(a * (b * c)), \beta(b), \delta\}. \end{aligned}$$

So, β is a fuzzy UP-ideal with thresholds ε and δ of B .

(4) Assume that μ is an f -invariant fuzzy strongly UP-ideal with thresholds ε and δ of A with inf property. Since $f(0_A) = 0_B$, we have $f^{-1}(0_B) \neq \emptyset$. Then there exists $x_1 \in f^{-1}(0_B)$ such that $\mu(x_1) = \beta(0_B)$. Thus $f(x_1) = 0_B = f(0_A)$. So $\mu(x_1) = \mu(0_A)$, because μ is f -invariant. Hence, $\mu(0_A) = \beta(0_B)$.

Let $y \in B$. Since f is surjective, we have $f^{-1}(y) \neq \emptyset$. Then there exists $x \in f^{-1}(y)$ such that $\mu(x) = \beta(y)$. Thus

$$\max\{\beta(0_B), \varepsilon\} = \max\{\mu(0_A), \varepsilon\} \geq \min\{\mu(x), \delta\} = \min\{\beta(y), \delta\}.$$

Next, let $a, b, c \in B$. Then by Lemma 5.8, there exist $a_0 \in f^{-1}(a), b_0 \in f^{-1}(b)$ and $c_0 \in f^{-1}(c)$ such that $\beta(a) = \mu(a_0), \beta(b) = \mu(b_0)$ and $\beta((c * b) * (c * a)) = \mu((c_0 \cdot b_0) \cdot (c_0 \cdot a_0))$. Thus

$$\begin{aligned} \max\{\beta(a), \varepsilon\} &= \max\{\mu(a_0), \varepsilon\} \geq \min\{\mu((c_0 \cdot b_0) \cdot (c_0 \cdot a_0)), \mu(b_0), \delta\} \\ &= \min\{\beta((c * b) * (c * a)), \beta(b), \delta\}. \end{aligned}$$

So, β is a fuzzy strongly UP-ideal with thresholds ε and δ of B . □

Theorem 5.10. *Let $(A, \cdot, 0_A)$ and $(B, *, 0_B)$ be UP-algebras and let $f: A \rightarrow B$ be a UP-homomorphism. Then the following statements hold:*

- (1) *if β is a fuzzy UP-subalgebra with thresholds ε and δ of B , then μ is a fuzzy UP-subalgebra with thresholds ε and δ of A ,*
- (2) *if β is a fuzzy UP-filter with thresholds ε and δ of B , then μ is a fuzzy UP-filter with thresholds ε and δ of A ,*
- (3) *if β is a fuzzy UP-ideal with thresholds ε and δ of B , then μ is a fuzzy UP-ideal with thresholds ε and δ of A , and*
- (4) *if β is a fuzzy strongly UP-ideal with thresholds ε and δ of B , then μ is a fuzzy strongly UP-ideal with thresholds ε and δ of A .*

Proof. (1) Assume that β is a fuzzy UP-subalgebra with thresholds ε and δ of B . Let $x, y \in A$. Then

$$\begin{aligned} \max\{\mu(x \cdot y), \varepsilon\} &= \max\{(\beta \circ f)(x \cdot y), \varepsilon\} = \max\{\beta(f(x \cdot y)), \varepsilon\} \\ &= \max\{\beta(f(x) * f(y)), \varepsilon\} \\ &\geq \min\{\beta(f(x)), \beta(f(y)), \delta\} \\ &= \min\{(\beta \circ f)(x), (\beta \circ f)(y), \delta\} \\ &= \min\{\mu(x), \mu(y), \delta\}. \end{aligned}$$

Thus, μ is a fuzzy UP-subalgebra with thresholds ε and δ of A .

(2) Assume that β is a fuzzy UP-filer with thresholds ε and δ of B . Let $x \in A$. Then

$$\begin{aligned} \max\{\mu(0_A), \varepsilon\} &= \max\{(\beta \circ f)(0_A), \varepsilon\} = \max\{\beta(f(0_A)), \varepsilon\} \\ (f(0_A) = 0_B) \quad &= \max\{\beta(0_B), \varepsilon\} \\ &\geq \min\{\beta(f(x)), \delta\} \\ &= \min\{(\beta \circ f)(x), \delta\} \\ &= \min\{\mu(x), \delta\}. \end{aligned}$$

Let $x, y \in A$. Then

$$\begin{aligned} \max\{\mu(y), \varepsilon\} &= \max\{(\beta \circ f)(y), \varepsilon\} = \max\{\beta(f(y)), \varepsilon\} \\ &\geq \min\{\beta(f(x) * f(y)), \beta(f(x)), \delta\} \\ &= \min\{\beta(f(x \cdot y)), \beta(f(x)), \delta\} \\ &= \min\{(\beta \circ f)(x \cdot y), (\beta \circ f)(x), \delta\} \\ &= \min\{\mu(x \cdot y), \mu(x), \delta\}. \end{aligned}$$

Thus, μ is a fuzzy UP-filter with thresholds ε and δ of A .

(3) Assume that β is a fuzzy UP-ideal with thresholds ε and δ of B . Let $x \in A$. Then

$$\begin{aligned} \max\{\mu(0_A), \varepsilon\} &= \max\{(\beta \circ f)(0_A), \varepsilon\} = \max\{\beta(f(0_A)), \varepsilon\} \\ (f(0_A) = 0_B) \qquad &= \max\{\beta(0_B), \varepsilon\} \\ &\geq \min\{\beta(f(x)), \delta\} \\ &= \min\{(\beta \circ f)(x), \delta\} \\ &= \min\{\mu(x), \delta\}. \end{aligned}$$

Let $x, y, z \in A$. Then

$$\begin{aligned} \max\{\mu(x \cdot z), \varepsilon\} &= \max\{(\beta \circ f)(x \cdot z), \varepsilon\} = \max\{\beta(f(x \cdot z)), \varepsilon\} \\ &= \max\{\beta(f(x) * f(z)), \varepsilon\} \\ &\geq \min\{\beta(f(x) * (f(y) * f(z))), \beta(f(y)), \delta\} \\ &= \min\{\beta(f(x) * f(y \cdot z)), \beta(f(y)), \delta\} \\ &= \min\{\beta(f(x \cdot (y \cdot z))), \beta(f(y)), \delta\} \\ &= \min\{(\beta \circ f)(x \cdot (y \cdot z)), (\beta \circ f)(y), \delta\} \\ &= \min\{\mu(x \cdot (y \cdot z)), \mu(y), \delta\}. \end{aligned}$$

Thus, μ is a fuzzy UP-ideal with thresholds ε and δ of A .

(4) Assume that β is a fuzzy strongly UP-ideal with thresholds ε and δ of B . Let $x \in A$. Then

$$\begin{aligned} \max\{\mu(0_A), \varepsilon\} &= \max\{(\beta \circ f)(0_A), \varepsilon\} = \max\{\beta(f(0_A)), \varepsilon\} \\ (f(0_A) = 0_B) \qquad &= \max\{\beta(0_B), \varepsilon\} \\ &\geq \min\{\beta(f(x)), \delta\} \\ &= \min\{(\beta \circ f)(x), \delta\} \\ &= \min\{\mu(x), \delta\}. \end{aligned}$$

Let $x, y, z \in A$. Then

$$\begin{aligned} \max\{\mu(x), \varepsilon\} &= \max\{(\beta \circ f)(x), \varepsilon\} \\ &= \max\{\beta(f(x)), \varepsilon\} \\ &\geq \min\{\beta((f(z) * f(y)) * (f(z) * f(x))), \beta(f(y)), \delta\} \\ &= \min\{\beta(f(z \cdot y) * f(z \cdot x)), \beta(f(y)), \delta\} \\ &= \min\{\beta(f((z \cdot y) \cdot (z \cdot x))), \beta(f(y)), \delta\} \\ &= \min\{(\beta \circ f)((z \cdot y) \cdot (z \cdot x)), (\beta \circ f)(y), \delta\} \\ &= \min\{\mu((z \cdot y) \cdot (z \cdot x)), \mu(y), \delta\}. \end{aligned}$$

Thus, μ is a fuzzy strongly UP-ideal with thresholds ε and δ of A . □

6. CONCLUSIONS

In the present paper, we have introduced the notion of fuzzy UP-subalgebras (resp., fuzzy UP-filters, fuzzy UP-ideals, fuzzy strongly UP-ideals) with thresholds of UP-algebras and proved its generalizations. We think this work would enhance

the scope for further study in UP-algebras and related algebraic systems. It is our hope that this work would serve as a foundation for the further study in UP-algebras and related algebraic systems.

Acknowledgements. This work was financially supported by the University of Phayao.

REFERENCES

- [1] A. Akram and K. H. Dar, Generalized fuzzy K-algebras, VDM Verlag 2010.
- [2] T. Guntasow, S. Sajak, A. Jomkham, and A. Iampan, Fuzzy translations of a fuzzy set in UP-algebras, *J. Indones. Math. Soc.* 23 (2) (2017) 1–19.
- [3] A. Iampan, A new branch of the logical algebra: UP-algebras, *J. Algebra Relat. Top.* 5 (1) (2017) 35–54.
- [4] Y. B. Jun, Closed fuzzy ideals in BCI-algebras, *Math. Japon.* 38 (1) (1993) 199–202.
- [5] Y. B. Jun, On (α, β) -fuzzy subalgebras of BCK/BCI-algebras, *Bull. Korean Math. Soc.* 42 (4) (2005) 703–711.
- [6] Y. B. Jun, Fuzzy subalgebras with thresholds in BCK/BCI-algebras, *Commun. Korean Math. Soc.* 22 (2) (2007) 173–181.
- [7] W. Kaijiae, P. Pongsumpao, S. Arayarangsi, and A. Iampan, UP-algebras characterized by their anti-fuzzy UP-ideals and anti-fuzzy UP-subalgebras, *Ital. J. Pure Appl. Math.* 36 (2016) 667–692.
- [8] J. N. Mordeson and D. S. Malik, Fuzzy commutative algebra, World Scientific, Singapore 1998.
- [9] P. Pongsumpao, W. Kaijiae, S. Arayarangsi, and A. Iampan, Fuzzy UP-ideals and fuzzy UP-subalgebras of UP-algebras in term of level subsets, Manuscript submitted for publication, February 2018.
- [10] A. Rosenfeld, Fuzzy groups, *J. Math. Anal. Appl.* 35 (1971) 512–517.
- [11] A. B. Saeid, Redefined fuzzy subalgebra (with thresholds) of BCK/BCI-algebras, *Iran. J. Math. Sci. Inform.* 4 (2) (2009) 19–24.
- [12] T. Senapati, T-fuzzy KU-subalgebras of KU-algebras, *Ann. Fuzzy Math. Inform.* 10 (2) (2015) 261–270.
- [13] T. Senapati, Y. B. Jun, and K. P. Shum, Cubic set structure applied in UP-algebras, *Discrete Math. Algorithms Appl.* 10 (4) (2018) 1850049.
- [14] T. Senapati, G. Muhiuddin, and K. P. Shum, Representation of UP-algebras in interval-valued intuitionistic fuzzy environment, *Ital. J. Pure Appl. Math.* 38 (2017) 497–517.
- [15] J. Somjanta, N. Thuekaew, P. Kumpeangkeaw, and A. Iampan, Fuzzy sets in UP-algebras, *Ann. Fuzzy Math. Inform.* 12 (6) (2016) 739–756.
- [16] L. A. Zadeh, Fuzzy sets, *Inf. Cont.* 8 (1965) 338–353.
- [17] J. Zhan, Y. B. Jun, and B. Davvaz, On $(\in, \in \vee q)$ -fuzzy ideals of BCI-algebras, *Iran. J. Fuzzy Syst.* (1) 6 (2009) 81–94.

NOPPHARAT DOKKHAMDANG (noppharattak@gmail.com)

Department of Mathematics, School of Science, University of Phayao, Phayao, postal code 56000, Thailand

AKEKARIN KESORN (the.akekarinza@gmail.com)

Department of Mathematics, School of Science, University of Phayao, Phayao, postal code 56000, Thailand

AIYARED IAMPAN (aiyared.ia@up.ac.th)

Department of Mathematics, School of Science, University of Phayao, Phayao, postal code 56000, Thailand