


Closure fuzzy ideals of MS-algebras

BERHANU ASSAYE ALABA, TEFERI GETACHEW ALEMAYEHU



Reprinted from the
Annals of Fuzzy Mathematics and Informatics
Vol. 16, No. 2, October 2018

Closure fuzzy ideals of MS-algebras

BERHANU ASSAYE ALABA, TEFERI GETACHEW ALEMAYEHU

Received 20 June 2018; Revised 13 July 2018; Accepted 30 July 2018

ABSTRACT. In this paper, we introduce concepts of dominator fuzzy ideals and closure fuzzy ideals in MS-algebras and study some properties of these fuzzy ideals. It is proved that the lattice of all closure fuzzy ideals is isomorphic to the fuzzy ideal lattice of the lattice of all principal dominator ideals. A set of equivalent conditions is obtained to characterize closure fuzzy ideals of MS-algebras. Finally some properties of closure fuzzy ideals are studied with respect to homomorphisms.

2010 AMS Classification: 06D99, 06D05, 06D30

Keywords: MS-algebra, Fuzzy ideals, Dominator fuzzy ideals, Closure fuzzy ideals, Homomorphisms.

Corresponding Author: Teferi Getachew Alemayehu (teferigetachew3@gmail.com)

1. INTRODUCTION

An Ockham algebra is a bounded distributive lattice with a dual endomorphism. The class of all Ockham algebras contains the well-known classes for examples Boolean algebras, de Morgan algebras, Kleene algebras and Stone algebras [5]. Blyth and Varlet [6] defined a subclass of Ockham algebras so called MS-algebras which generalize both de Morgan algebras and Stone algebras. These algebras belong to the class of Ockham algebras introduced by Berman [4]. The class of all MS-algebras forms an equational class. Blyth and Varlet [7] characterized the subvarieties of MS-algebras. Recently, Rao [13] introduced the notion of e-filters of MS-algebras, Rao [12] introduced the concepts of boosters and β -filters of MS-algebras and also Badawy and Rao [3] introduced the closure ideals of MS-algebras.

On the other hand, fuzzy set theory introduced by Zadeh [17] is generalization of classical set theory. Next, Rosenfeld [11] applied it to group theory and developed the theory of fuzzy subgroups. Also, many authors have worked on fuzzy lattice theory. They introduced the concepts of fuzzy sublattice, fuzzy ideal, fuzzy prime ideal, in a lattice and gave some interesting results (see [1, 2, 9, 10, 14, 16]). In this paper,

we introduce dominator fuzzy subsets and dominator fuzzy ideals and study some basic properties of dominator fuzzy subsets and dominator fuzzy ideals. Moreover the concept of closure fuzzy ideals is introduced in MS-algebras. Some properties of closure fuzzy ideals of an MS-algebra are observed. It is proved that the class $FI_C(L)$ of all closure fuzzy ideals of an MS-algebra L is a bounded distributive lattice with set inclusion. Also it is proved that $FI_C(L)$ is isomorphic to the ideal lattice of $FI(M_{\circ\circ}(L))$. Finally, some properties of closure fuzzy ideals are studied with respect to homomorphisms. The concept of dominator fuzzy ideal preserving homomorphism from an MS-algebra L into another MS-algebra M is introduced as a homomorphism f satisfying the condition $f(\mu_{\circ\circ}) = \{f(\mu)\}_{\circ\circ}$, for any fuzzy ideal μ of L . It is proved that the images and the inverse images, under this homomorphism, of closure fuzzy ideals are again closure fuzzy ideals.

2. PRELIMINARIES

In this section, we recall some definitions and results which will be used in this paper. For in details in ordinary crisp theory of closure ideals of MS-algebras, we refer to [3].

Definition 2.1 ([6]). An MS-algebra is an algebra $(L, \vee, \wedge, \circ, 0, 1)$ of type $(2, 2, 1, 0, 0)$, such that $(L, \vee, \wedge, 0, 1)$ is a bounded distributive lattice and $a \rightarrow a^\circ$ is a unary operation satisfies: $a \leq a^{\circ\circ}$, $(a \wedge b)^\circ = a^\circ \vee b^\circ$ and $1^\circ = 0$.

A Stone algebra $S = (S, \vee, \wedge, *, 0, 1)$ is also a bounded distributive lattice, endowed with a unary operation $x \rightarrow x^*$ satisfying $(x \wedge y)^* = x^* \vee y^*$, $x \wedge x^* = 0$ and $0^* = 1$.

A de Morgan algebra is an algebra $(L, \vee, \wedge, \bar{}, 0, 1)$ of type $(2, 2, 1, 0, 0)$ such that $(L, \vee, \wedge, 0, 1)$ is a bounded distributive lattice and $a \rightarrow \bar{a}$ is a unary operation satisfies: $\bar{\bar{a}} = a$, $\overline{(a \wedge b)} = \bar{a} \vee \bar{b}$ and $\bar{1} = 0$.

Lemma 2.2 ([6]). For any two elements a, b of an MS-algebra, we have the following:

- (1) $0^\circ = 1$,
- (2) $a \leq b \Rightarrow b^\circ \leq a^\circ$,
- (3) $a^{\circ\circ} = a^\circ$,
- (4) $(a \vee b)^\circ = a^\circ \wedge b^\circ$,
- (5) $(a \vee b)^{\circ\circ} = a^{\circ\circ} \vee b^{\circ\circ}$,
- (6) $(a \wedge b)^{\circ\circ} = a^{\circ\circ} \wedge b^{\circ\circ}$.

Definition 2.3 ([3]). For any nonempty subset A of an MS-algebra L , define the dominator of A as follows: $A_{\circ\circ} = \{x \in L : x \leq a^{\circ\circ} \text{ for some } a \in A\}$.

Note that $\{0\}_{\circ\circ} = \{0\}$ and $L_{\circ\circ} = L$.

Definition 2.4 ([3]). An ideal I of an MS-algebra L is called a dominator ideal, if $I = I_{\circ\circ}$.

The set of all dominator ideals of L denoted by $I_{\circ\circ}(L)$ and $I_{\circ\circ}(L)$ form a bounded distributive lattice with set inclusion.

Definition 2.5 ([8]). For any elements a in a lattice L , the principal ideal of L generated by a , $[a]$, defined by $[a] = \{x \in L : x \leq a\}$.

Definition 2.6 ([3]). For any element a of an MS-algebra L , the dominator $\{a\}_{\circ\circ}$ is called a principal dominator ideal.

Note that $\{0\}_{\circ\circ} = \{0\}$ and $\{1\}_{\circ\circ} = L$ and for any $a \in L$, $\{a\}_{\circ\circ} = [a]_{\circ\circ} = (a^{\circ\circ})$. The set of all principal dominator ideals of L denoted by $M_{\circ\circ}(L)$, i.e., $M_{\circ\circ}(L) = \{[a]_{\circ\circ} : a \in L\}$. $M_{\circ\circ}(L)$ is a bounded sublattice of $I_{\circ\circ}(L)$ and it is a de Morgan algebra.

Definition 2.7 ([3]). Let L be an MS-algebra. For any ideal I of L , define an operator $\sigma : I(L) \rightarrow I(M_{\circ\circ}(L))$ as follows: $\sigma(I) = \{[x]_{\circ\circ} : x \in I\}$.

Definition 2.8 ([3]). Let L be an MS-algebra. For any ideal \bar{I} of $M_{\circ\circ}(L)$, define an operator $\overleftarrow{\sigma} : I(M_{\circ\circ}(L)) \rightarrow I(L)$ as follows: $\overleftarrow{\sigma}(\bar{I}) = \{x \in L : [x]_{\circ\circ} \in \bar{I}\}$.

We recall that for any nonempty set L , the characteristic function of L ,

$$\chi_L(x) = \begin{cases} 1 & \text{if } x \in L, \\ 0 & \text{if } x \notin L. \end{cases}$$

Definition 2.9 ([3]). An ideal I of an MS-algebra L is a closure ideal, if $\overleftarrow{\sigma}\sigma(I) = I$.

Definition 2.10 ([2]). Let μ be a fuzzy subset of $(L, \wedge, \vee, 0, 1)$. For any $\alpha \in [0, 1]$, we shall denote the level subset $\mu^{-1}([\alpha, 1])$ by simply μ_α , i.e.,

$$\mu_\alpha = \{x \in L : \alpha \leq \mu(x)\}.$$

A fuzzy subset μ of L is proper, if it is a non constant function. A fuzzy subset μ such that $\mu(x) = 0$, for all $x \in L$ is an improper fuzzy subset.

Definition 2.11 ([11]). Let μ and θ be fuzzy subsets of a set L . Define the fuzzy subsets $\mu \cup \theta$ and $\mu \cap \theta$ of L as follows: for each $x \in L$,

$$(\mu \cup \theta)(x) = \mu(x) \vee \theta(x) \text{ and } (\mu \cap \theta)(x) = \mu(x) \wedge \theta(x).$$

Then $\mu \cup \theta$ and $\mu \cap \theta$ are called the union and intersection of μ and θ respectively.

We define the binary operations " \vee " and " \wedge " on all fuzzy subsets of L as:

$$(\mu \vee \theta)(x) = \sup\{\mu(a) \wedge \theta(b) : a, b \in L, a \vee b = x\}$$

and

$$(\mu \wedge \theta)(x) = \sup\{\mu(a) \wedge \theta(b) : a, b \in L, a \wedge b = x\}.$$

If μ and θ are fuzzy ideals of L , then $\mu \wedge \theta = \mu \cap \theta$ and $\mu \vee \theta$ is a fuzzy ideal generated by $\mu \cup \theta$.

Theorem 2.12 ([14]). Let μ be a fuzzy subset of L . Then μ is a fuzzy ideal of L if and only if any one of the following conditions are satisfied:

- (1) $\mu(0) = 1$ and $\mu(x \vee y) = \mu(x) \wedge \mu(y)$, for all $x, y \in L$,
- (2) $\mu(0) = 1$ and $\mu(x \vee y) \geq \mu(x) \wedge \mu(y)$ and $\mu(x \wedge y) \geq \mu(x) \vee \mu(y)$, for all $x, y \in L$.

Proposition 2.13 ([2]). Let μ is a fuzzy ideal of L . If $x \leq y$, then $\mu(x) \geq \mu(y)$, for all $x, y \in L$.

Theorem 2.14 ([14]). Let μ be a fuzzy subset of L . Then μ is a fuzzy ideal of L if and only if for any $\alpha \in [0, 1]$ such that $\mu_\alpha \neq \emptyset$, μ_α is an ideal of L .

Definition 2.15 ([2]). A proper fuzzy ideal μ of L is called prime fuzzy ideal, if for any two fuzzy ideals η, ν of L , $\eta \cap \nu \subseteq \mu$ implies $\eta \subseteq \mu$ or $\nu \subseteq \mu$.

Theorem 2.16 ([15]). For any prime ideal P of L and $\alpha \in [0, 1)$ if and only if there exists a prime fuzzy ideal P_α^1 of L is given by: for each $x \in L$,

$$P_\alpha^1(x) = \begin{cases} 1 & \text{if } x \in P, \\ \alpha & \text{if } x \notin P. \end{cases}$$

Theorem 2.17 ([16]). Let $f : L \rightarrow L'$ be an onto homomorphism. Then $f(\mu)$ is a fuzzy ideal of L' if μ is a fuzzy ideal of L .

Theorem 2.18 ([1]). Let $f : L \rightarrow L'$ be homomorphism. Then $f^{-1}(\mu)$ is a fuzzy ideal of L , if μ is a fuzzy ideal of L' .

Throughout in the next sections L stands for an MS-algebra unless otherwise mentioned.

3. DOMINATOR FUZZY SETS, DOMINATOR FUZZY IDEALS OF MS-ALGEBRAS

In this section, the concepts of dominator fuzzy subsets and dominator fuzzy ideals are introduced in MS-algebras. Some properties of these fuzzy ideals are investigated.

Definition 3.1. For any non-empty fuzzy subset μ of an MS-algebra L , define the dominator of μ as follows: $\mu_{\circ\circ}(x) = \sup\{\mu(a) : x \leq a^{\circ\circ}, a \in L\}$, for all $x \in L$

Obviously, $\{\chi_{\{0\}}\}_{\circ\circ} = \chi_{\{0\}}$ and $\{\chi_L\}_{\circ\circ} = \chi_L$.

Lemma 3.2. For any two fuzzy subsets μ, θ of MS-algebra L , we have the following:

- (1) $\mu \subseteq \mu_{\circ\circ}$,
- (2) $\mu \subseteq \theta$ implies $\mu_{\circ\circ} \subseteq \theta_{\circ\circ}$,
- (3) $\{\mu_{\circ\circ}\}_{\circ\circ} = \mu_{\circ\circ}$.

Proof. (1) $\mu_{\circ\circ}(x) = \sup\{\mu(a) : x \leq a^{\circ\circ}, a \in L\} \geq \mu(x)$, since $x \leq x^{\circ\circ}$. Then $\mu \subseteq \mu_{\circ\circ}$.

(2) Suppose $\mu \subseteq \theta$. Then for all $x \in L$,

$$\mu_{\circ\circ}(x) = \sup\{\mu(a) : x \leq a^{\circ\circ}, a \in L\} \leq \sup\{\theta(a) : x \leq a^{\circ\circ}, a \in L\} = \theta_{\circ\circ}(x).$$

Thus $\mu_{\circ\circ} \subseteq \theta_{\circ\circ}$.

- (3) Since $a^{\circ\circ} \leq y^{\circ\circ\circ\circ} = y^{\circ\circ}$,

$$\begin{aligned} \{\mu_{\circ\circ}\}_{\circ\circ}(x) &= \sup\{\mu_{\circ\circ}(a) : x \leq a^{\circ\circ}\} \\ &= \sup\{\sup\{\mu(y) : a \leq y^{\circ\circ}\} : x \leq a^{\circ\circ}\} \\ &= \sup\{\mu(y) : x \leq y^{\circ\circ}\} = \mu_{\circ\circ}(x). \end{aligned}$$

Then $\{\mu_{\circ\circ}\}_{\circ\circ} = \mu_{\circ\circ}$. □

Lemma 3.3. For any two fuzzy ideals μ and θ of MS-algebra L , the following hold:

- (1) $\mu_{\circ\circ}$ is a fuzzy ideal of L ,
- (2) $(\mu \cap \theta)_{\circ\circ} = \mu_{\circ\circ} \cap \theta_{\circ\circ}$,
- (3) $(\mu \vee \theta)_{\circ\circ} = \mu_{\circ\circ} \vee \theta_{\circ\circ}$.

Proof. (1) Clearly, $\mu_{\circ\circ}(0) = 1$. Let $x, y \in L$. Then

$$\begin{aligned} \mu_{\circ\circ}(x) \wedge \mu_{\circ\circ}(y) &= \sup\{\mu(a) : x \leq a^{\circ\circ}, a \in L\} \wedge \sup\{\mu(b) : y \leq b^{\circ\circ}, b \in L\} \\ &= \sup\{\mu(a) \wedge \mu(b) : x \leq a^{\circ\circ}, y \leq b^{\circ\circ}\} \\ &\leq \sup\{\mu(a \vee b) : x \vee y \leq a^{\circ\circ} \vee b^{\circ\circ} = (a \vee b)^{\circ\circ}, a, b \in L\} \\ &= \mu_{\circ\circ}(x \vee y) \end{aligned}$$

and

$$\begin{aligned} \mu_{\circ\circ}(x) \vee \mu_{\circ\circ}(y) &= \sup\{\mu(a) : x \leq a^{\circ\circ}, a \in L\} \vee \sup\{\mu(b) : y \leq b^{\circ\circ}, b \in L\} \\ &= \sup\{\mu(a) \vee \mu(b) : x \leq a^{\circ\circ}, y \leq b^{\circ\circ}, a, b \in L\} \\ &\leq \sup\{\mu(a \wedge b) : x \wedge y \leq a^{\circ\circ} \wedge b^{\circ\circ} = (a \wedge b)^{\circ\circ}, a, b \in L\} \\ &= \mu_{\circ\circ}(x \wedge y). \end{aligned}$$

Thus $\mu_{\circ\circ}$ is fuzzy ideal of L .

(2) Clearly, $(\mu \cap \theta)_{\circ\circ} \subseteq \mu_{\circ\circ} \cap \theta_{\circ\circ}$. Let $x \in L$. Then

$$\begin{aligned} (\mu_{\circ\circ} \cap \theta_{\circ\circ})(x) &= \mu_{\circ\circ}(x) \wedge \theta_{\circ\circ}(x) \\ &= \sup\{\mu(a) : x \leq a^{\circ\circ}, a \in L\} \wedge \sup\{\theta(b) : x \leq b^{\circ\circ}, b \in L\} \\ &\leq \sup\{\mu(a \wedge b) : x \leq (a \wedge b)^{\circ\circ}, a, b \in L\} \wedge \sup\{\theta(a \wedge b) : x \leq (a \wedge b)^{\circ\circ}, a, b \in L\}. \end{aligned}$$

Since $x \leq a^{\circ\circ}$ and $x \leq b^{\circ\circ}$, $x \leq (a \wedge b)^{\circ\circ}$.

$$\begin{aligned} &= \sup\{\mu(a \wedge b) \wedge \theta(a \wedge b) : x \leq (a \wedge b)^{\circ\circ}\} \\ &= \sup\{(\mu \cap \theta)(a \wedge b) : x \leq (a \wedge b)^{\circ\circ}\} \\ &= (\mu \cap \theta)_{\circ\circ}(x). \end{aligned}$$

Thus $(\mu \cap \theta)_{\circ\circ} \subseteq \mu_{\circ\circ} \cap \theta_{\circ\circ}$.

(3) Let $x \in L$. Then

$$\begin{aligned} (\mu_{\circ\circ} \vee \theta_{\circ\circ})(x) &= \sup\{\mu_{\circ\circ}(x_1) \wedge \theta_{\circ\circ}(x_2) : x = x_1 \vee x_2\} \\ &= \sup\{\mu(a_1) : x_1 \leq a_1^{\circ\circ}\} \wedge \sup\{\theta(a_2) : x_2 \leq a_2^{\circ\circ}\} : x_1 \vee x_2 = x \\ &= \sup\{\sup\{\mu(a_1) \wedge \theta(a_2) : x_1 \leq a_1^{\circ\circ}, x_2 \leq a_2^{\circ\circ}\} : x = x_1 \vee x_2\} \\ &= \sup\{\mu(a_1) \wedge \theta(a_2) : x_1 \leq a_1^{\circ\circ}, x_2 \leq a_2^{\circ\circ}, \text{where } x = x_1 \vee x_2\}. \end{aligned}$$

Put $A = \{\alpha, \text{ where } \alpha = \mu(a_1) \wedge \theta(a_2) : x_1 \leq a_1^{\circ\circ}, x_2 \leq a_2^{\circ\circ}, \text{where } x = x_1 \vee x_2\}$.

Then

$$\begin{aligned} (\mu \vee \theta)_{\circ\circ}(x) &= \sup\{(\mu \vee \theta)(a) : x \leq a^{\circ\circ}\} \\ &= \sup\{\sup\{\mu(a_1) \wedge \theta(a_2) : a_1 \vee a_2 = a\} : x \leq a^{\circ\circ}\} \\ &= \sup\{\mu(a_1) \wedge \theta(a_2) : a_1 \vee a_2 = a, x \leq a^{\circ\circ}\}. \end{aligned}$$

Put $B = \{\lambda, \text{ where } \lambda = \mu(a_1) \wedge \theta(a_2) : a_1 \vee a_2 = a, x \leq a^{\circ\circ}\}$ and we prove that $A = B$. Let $\alpha \in A$. Then $\alpha = \mu(a_1) \wedge \theta(a_2)$ such that $x_1 \leq a_1^{\circ\circ}, x_2 \leq a_2^{\circ\circ}$, where $x = x_1 \vee x_2$. Thus $x = x_1 \vee x_2 \leq a_1^{\circ\circ} \vee a_2^{\circ\circ} = (a_1 \vee a_2)^{\circ\circ} = (a)^{\circ\circ}, a = a_1 \vee a_2$. So $\alpha \in B$, i.e., $A \subseteq B$.

Let $\lambda \in B$. Then $\lambda = \mu(a_1) \wedge \theta(a_2)$ such that $a_1 \vee a_2 = a, x \leq a^{\circ\circ}$. Thus $x \leq (a_1 \wedge a_2)^{\circ\circ} = (a_1)^{\circ\circ} \vee (a_2)^{\circ\circ}$. So $x = x \wedge (a_1^{\circ\circ} \vee a_2^{\circ\circ}) = (x \wedge a_1^{\circ\circ}) \vee (x \wedge a_2^{\circ\circ})$. Let $x_1 = x \wedge a_1^{\circ\circ}, x_2 = x \wedge a_2^{\circ\circ}$. Since $x = x_1 \vee x_2, x_1 \leq a_1^{\circ\circ}, x_2 \leq a_2^{\circ\circ}$, where $a = a_1 \vee a_2$. This implies $\lambda \in A$ and $B \subseteq A$. Hence $A = B$, i.e., $\sup A = \sup B$. Therefore we get $(\mu \vee \theta)_{\circ\circ} = \mu_{\circ\circ} \vee \theta_{\circ\circ}$. \square

Corollary 3.4. If $\{\mu_i : i \in \Omega\}$ is a family of fuzzy ideals of L , then

$$\{\bigcap_{i \in \Omega} \mu_i\}_{\circ\circ} = \bigcap_{i \in \Omega} (\mu_i)_{\circ\circ}.$$

Definition 3.5. A fuzzy ideal μ of an MS-algebra L is called a dominator fuzzy ideal, if $\mu = \mu_{\circ\circ}$.

Theorem 3.6. μ is dominator fuzzy ideal in MS-algebra L if and only if $\mu_\alpha, \alpha \in Im(\mu)$, is a dominator ideal of L .

Proof. Suppose μ is dominator fuzzy ideal of L . Then $\mu_\alpha = (\mu_{\circ\circ})_\alpha, \forall \alpha \in Im(\mu)$. Clearly, $\mu_\alpha \subseteq (\mu_\alpha)_{\circ\circ}$. Let $x \in (\mu_\alpha)_{\circ\circ}$. Thus there exists $y \in \mu_\alpha$ such that $x \leq y^{\circ\circ}$. So $\mu_{\circ\circ}(x) = \sup\{\mu(y) : x \leq y^{\circ\circ}\} \geq \mu(y) \geq \alpha$. Hence $x \in (\mu_{\circ\circ})_\alpha = \mu_\alpha$. Therefore $(\mu_\alpha)_{\circ\circ} = \mu_\alpha$.

Conversely, suppose $(\mu_\alpha)_{\circ\circ} = \mu_\alpha$. Then, clearly, $\mu \subseteq \mu_{\circ\circ}$. Now let $\alpha = (\mu_{\circ\circ})(x) = \sup\{\mu(y) : x \leq y^{\circ\circ}\}$. Then for each $\epsilon > 0$, there exists $a \in L$ such that $\alpha - \epsilon \leq \mu(a)$ for $x \leq a^{\circ\circ}$. Since ϵ is arbitrary, $\alpha \leq \mu(a)$ for $x \leq a^{\circ\circ}$. Thus $a \in \mu_\alpha$, for $x \leq a^{\circ\circ}$. So $x \in (\mu_\alpha)_{\circ\circ} = \mu_\alpha$. Hence $\mu(x) \geq \alpha = (\mu_{\circ\circ})(x)$. Therefore $\mu = \mu_{\circ\circ}$. \square

Corollary 3.7. I is dominator ideal of MS-algebra L if and only if χ_I is dominator fuzzy ideal of L .

We denote the set of all dominator fuzzy ideals of L by $\mu_{\circ\circ}(L)$. Now we have the following theorem.

Theorem 3.8. The set of all dominator fuzzy ideals of L , $\mu_{\circ\circ}(L)$, is a bounded distributive lattice with set inclusion.

Proof. It is obvious by Lemma 3.3. \square

4. CLOSURE FUZZY IDEALS OF MS-ALGEBRA

In this section, the notion of closure fuzzy ideals is introduced in MS-algebras and some properties are observed. We denote the set of all fuzzy ideals of L by $FI(L)$ and the set of all fuzzy ideals of $M_{\circ\circ}(L)$ by $FI(M_{\circ\circ}(L))$.

Definition 4.1. Let L be an MS-algebra. For any fuzzy ideal μ of L , define an operator $\sigma : FI(L) \rightarrow FI(M_{\circ\circ}(L))$ as follows: for all $a \in L$,

$$\sigma(\mu)((a]_{\circ\circ}) = \sup\{\mu(b) : (a]_{\circ\circ} = (b]_{\circ\circ}, b \in L\}.$$

Definition 4.2. Let L be an MS-algebra. For any fuzzy ideal θ of $M_{\circ\circ}(L)$, define an operator $\overleftarrow{\sigma} : FI(M_{\circ\circ}(L)) \rightarrow FI(L)$ as follows: $\overleftarrow{\sigma}(\theta)(a) = \theta((a]_{\circ\circ})$, for all $a \in L$.

Lemma 4.3. Let L be an MS-algebra. Then we have the following:

- (1) for any fuzzy ideal μ of L , $\sigma(\mu)$ is fuzzy ideal of $M_{\circ\circ}(L)$,
- (2) for any fuzzy ideal θ of $M_{\circ\circ}(L)$, $\overleftarrow{\sigma}(\theta)$ is fuzzy ideal of L ,
- (3) $\overleftarrow{\sigma}$ and σ are isotones,
- (4) $\sigma(\overleftarrow{\sigma})(\mu) = \mu$, for all fuzzy ideal μ of $M_{\circ\circ}(L)$,
- (5) σ is a homomorphism from fuzzy ideal of L into fuzzy ideal of $M_{\circ\circ}(L)$.

Proof. Let $a, b \in L$. Then $(a]_{\circ\circ}, (b]_{\circ\circ} \in M_{\circ\circ}(L)$.

(1) Clearly, $\sigma(\mu)((0]_{\circ\circ}) = 1$. Then

$$\begin{aligned} \sigma(\mu)((a]_{\circ\circ}) \wedge \sigma(\mu)((b]_{\circ\circ}) &= \sup\{\mu(x) : (x]_{\circ\circ} = (a]_{\circ\circ}\} \wedge \sup\{\mu(y) : (y]_{\circ\circ} = (a]_{\circ\circ}\} \\ &= \sup\{\mu(x) \wedge \mu(y) : (x]_{\circ\circ} = (a]_{\circ\circ}, (y]_{\circ\circ} = (a]_{\circ\circ}\} \\ &\leq \sup\{\mu(x \vee y) : (x]_{\circ\circ} = (a]_{\circ\circ}, (y]_{\circ\circ} = (a]_{\circ\circ}\} \\ &\leq \sup\{\mu(x \vee y) : (x \vee y]_{\circ\circ} = (a \vee b]_{\circ\circ}\} \\ &= \sigma(\mu)((a \vee b]_{\circ\circ}) \end{aligned}$$

$$= \sigma(\mu)((a]_{\infty} \vee (b]_{\infty}))$$

and

$$\begin{aligned} & \sigma(\mu)((a]_{\infty}) \vee \sigma(\mu)((b]_{\infty}) \\ &= \sup\{\mu(x) : (x]_{\infty} = (a]_{\infty}\} \vee \sup\{\mu(y) : (y]_{\infty} = (b]_{\infty}\} \\ &\leq \sup\{\mu(x \wedge y) : (x \wedge y]_{\infty} = (a \wedge b]_{\infty}\} \vee \sup\{\mu(x \wedge y) : (x \wedge y]_{\infty} = (a \wedge b]_{\infty}\} \\ &= \sup\{\mu(x \wedge y) : (x \wedge y]_{\infty} = (a \wedge b]_{\infty}\} \\ &= \sigma(\mu)((a \wedge b]_{\infty}) \\ &= \sigma(\mu)((a]_{\infty} \wedge (b]_{\infty}). \end{aligned}$$

Thus $\sigma(\mu)$ is fuzzy ideal of $M_{\infty}(L)$.

(2) Clearly, $\overleftarrow{\sigma}(\theta)(0) = \theta((0]_{\infty}) = 1$

and

$$\overleftarrow{\sigma}(\theta)(a \vee b) = \theta((a \vee b]_{\infty}) = \theta((a]_{\infty}) \wedge \theta((b]_{\infty}) = \overleftarrow{\sigma}(\theta)(a) \wedge \overleftarrow{\sigma}(\theta)(b).$$

(3) Let μ, θ be fuzzy ideals of $M_{\infty}(L)$ such that $\mu \subseteq \theta$. Then

$$\overleftarrow{\sigma}(\mu)(a) = \mu((a]_{\infty}) \subseteq \theta((a]_{\infty}) = \overleftarrow{\sigma}(\theta)(a).$$

Similarly, σ is an isotone.

(4) Let $(a]_{\infty} \in M_{\infty}(L)$ and let μ fuzzy ideal of $M_{\infty}(L)$. Then

$$\begin{aligned} \sigma \overleftarrow{\sigma}(\mu)((a]_{\infty}) &= \sup\{\overleftarrow{\sigma}(\mu)(b) : (a]_{\infty} = (a]_{\infty}\} \\ &= \sup\{\sigma(\mu)(b]_{\infty}) : (a]_{\infty} = (b]_{\infty}\} \\ &= \mu((a]_{\infty}). \end{aligned}$$

Thus $\sigma(\overleftarrow{\sigma})(\mu) = \mu$, for all fuzzy ideals μ of $M_{\infty}(L)$.

(5) Let $\mu, \theta \in FI(L)$. Since σ is isotone, $\sigma(\mu \cap \theta) \subseteq \sigma(\mu) \cap \sigma(\theta)$. On the other hand,

$$\begin{aligned} & (\sigma(\mu) \cap \sigma(\theta))((a]_{\infty}) \\ &= \sigma(\mu)((a]_{\infty}) \wedge \sigma(\theta)((a]_{\infty}) \\ &= \sup\{\mu(x) : (x]_{\infty} = (a]_{\infty}\} \wedge \sup\{\theta(y) : (y]_{\infty} = (a]_{\infty}\} \\ &\leq \sup\{\mu(x \wedge y) : (x \wedge y]_{\infty} = (a]_{\infty}\} \wedge \sup\{\theta(x \wedge y) : (x \wedge y]_{\infty} = (a]_{\infty}\} \\ &= \sup\{\mu(x \wedge y) \wedge \theta(x \wedge y) : (x \wedge y]_{\infty} = (a]_{\infty}\} \\ &= \sup\{(\mu \cap \theta)(x \wedge y) : (x \wedge y]_{\infty} = (a]_{\infty}\} \\ &= \sigma(\mu \cap \theta)((a]_{\infty}). \end{aligned}$$

Then $\sigma(\mu) \cap \sigma(\theta) \subseteq \sigma(\mu \cap \theta)$. Also clearly, $\sigma(\mu) \vee \sigma(\theta) \subseteq \sigma(\mu \vee \theta)$. Again

$$\begin{aligned} & \sigma(\mu \vee \theta)((x]_{\infty}) \\ &= \sup\{(\mu \vee \theta)(y) : (y]_{\infty} = (x]_{\infty}\} \\ &= \sup\{\sup\{\mu(a) \wedge \theta(b) : y = a \vee b\} : (y]_{\infty} = (x]_{\infty}\} \\ &\leq \sup\{\sup\{\mu(a_1) \wedge \theta(b_1) : (a_1]_{\infty} = (a]_{\infty}, (b_1]_{\infty} = (b]_{\infty}\} : (y]_{\infty} = (x]_{\infty}\} \\ &= \sup\{\sup\{\mu(a_1) : (a_1]_{\infty} = (a]_{\infty}\} \wedge \sup\{\theta(b_1) : (b_1]_{\infty} = (b]_{\infty}\} : \\ & \hspace{15em} (a]_{\infty} \vee (b]_{\infty} = (x]_{\infty}\} \\ &= \sup\{\sigma(\mu)((a]_{\infty}) \wedge \sigma(\theta)((b]_{\infty}) : (a]_{\infty} \vee (b]_{\infty} = (x]_{\infty}\} \\ &= (\sigma(\mu) \vee \sigma(\theta))((x]_{\infty}). \end{aligned}$$

Thus $\sigma(\mu \vee \theta) \subseteq \sigma(\mu) \vee \sigma(\theta)$. So $\sigma(\mu \vee \theta) = \sigma(\mu) \vee \sigma(\theta)$. Hence σ is a homomorphism from the lattice of fuzzy ideals of L into the lattice of fuzzy ideals of $M_{\infty}(L)$. \square

Theorem 4.4. *The map $\overleftarrow{\sigma} \sigma : FI(L) \rightarrow FI(L)$ is a closure operator, that is, for any $\mu, \theta \in FI(L)$,*

- (1) $\mu \subseteq \overleftarrow{\sigma} \sigma(\mu)$,
- (2) $\mu \subseteq \theta$ implies $\overleftarrow{\sigma} \sigma(\mu) \subseteq \overleftarrow{\sigma} \sigma(\theta)$,
- (3) $\overleftarrow{\sigma} \sigma\{\overleftarrow{\sigma} \sigma(\mu)\} = \overleftarrow{\sigma} \sigma(\mu)$.

Proof. The prove of (1) and (2) is straight forward.

(3) For any $x \in L$,

$$\begin{aligned} \overleftarrow{\sigma}\sigma\{\overleftarrow{\sigma}\sigma(\mu)\}(x) &= \sigma\{\overleftarrow{\sigma}\sigma(\mu)\}((x]_{\circ\circ}) \\ &= \sup\{\overleftarrow{\sigma}\sigma(\mu)(y) : (x]_{\circ\circ} = (y]_{\circ\circ}\} \\ &= \sup\{\sigma(\mu)((y]_{\circ\circ}) : ((x]_{\circ\circ} = ((y]_{\circ\circ})\} \\ &= \sigma(\mu)((x]_{\circ\circ}) \\ &= \overleftarrow{\sigma}\sigma(\mu)(x). \end{aligned}$$

□

Corollary 4.5. For any two fuzzy ideals μ and θ of an MS-algebra L , we have

$$\overleftarrow{\sigma}\sigma(\mu \cap \theta) = \overleftarrow{\sigma}\sigma(\mu) \cap \overleftarrow{\sigma}\sigma(\theta).$$


Proof. By Lemma 4.3 (5), $\sigma(\mu \cap \theta) = \sigma(\mu) \cap \sigma(\theta)$. Now for any $y \in L$,

$$\begin{aligned} \overleftarrow{\sigma}\sigma(\mu \cap \theta)(y) &= \sigma(\mu \cap \theta)((y]_{\circ\circ}) \\ &= \sigma(\mu)((y]_{\circ\circ}) \wedge \sigma(\theta)((y]_{\circ\circ}) \\ &= \overleftarrow{\sigma}\sigma(\mu)(y) \wedge \overleftarrow{\sigma}\sigma(\theta)(y). \end{aligned}$$

Then $\overleftarrow{\sigma}\sigma(\mu \cap \theta) = \overleftarrow{\sigma}\sigma(\mu) \cap \overleftarrow{\sigma}\sigma(\theta)$. □


Definition 4.6. A fuzzy ideal μ of L is called a closure fuzzy ideal, if $\overleftarrow{\sigma}\sigma(\mu) = \mu$.

Example 4.7. Let us consider the following figure



Then clearly $(L, \vee, \wedge, \circ, 0, 1)$ is an MS-algebra. Define a fuzzy subset μ of L as $\mu(x) = \mu(0) = 1$ and $\mu(1) = \mu(y) = 0.5$. We can easily verify that μ is closure fuzzy ideal of L .

Example 4.8. Let us consider the following figure with given table:



x	0	a	b	c	d	e	1
x°	1	e	e	e	e	e	0

Then clearly, $(L, \vee, \wedge, \circ, 0, 1)$ is an MS-algebra. Let $\mu : L \rightarrow [0, 1]$ defined as $\mu(a) = \mu(b) = 0.8, \mu(c) = \mu(d) = 0.6, \mu(e) = \mu(1) = 0$ and $\mu(0) = 1$. We can easily verify that μ is fuzzy ideal of L but not closure fuzzy ideal of L .

In the following Theorems we characterize closure fuzzy ideal in terms of level subsets and characteristic functions.

Theorem 4.9. For proper fuzzy subsets μ of L , μ is closure fuzzy ideal if and only if $\mu_\alpha, \forall \alpha \in Im(\mu)$, is closure ideal of L .

Proof. Suppose μ is closure fuzzy ideal of L . Then $(\overleftarrow{\sigma}\sigma(\mu))_\alpha = \mu_\alpha$. To prove each level subsets of μ is closure fuzzy ideal of L , it is enough to show $\overleftarrow{\sigma}\sigma(\mu_\alpha) = \mu_\alpha$. Clearly, $\mu_\alpha \subseteq \overleftarrow{\sigma}\sigma(\mu_\alpha)$. Let $x \in \overleftarrow{\sigma}\sigma(\mu_\alpha)$. Then $(x]_{\circ\circ} \in \sigma(\mu_\alpha)$. Thus there exists $y \in \mu_\alpha$ such that $(x]_{\circ\circ} = (y]_{\circ\circ}$, i.e., $\mu(y) \geq \alpha$ such that $(x]_{\circ\circ} = (y]_{\circ\circ}$. So $\sigma(\mu)((x]_{\circ\circ}) = \sup\{\mu(x) : (x]_{\circ\circ} = (y]_{\circ\circ}\} \geq \alpha$ and thus $\overleftarrow{\sigma}\sigma(\mu)(x) \geq \alpha$. Hence $x \in (\overleftarrow{\sigma}\sigma(\mu))_\alpha = \mu_\alpha$, i.e., $\overleftarrow{\sigma}\sigma(\mu_\alpha) \subseteq \mu_\alpha$. Therefore $\overleftarrow{\sigma}\sigma(\mu_\alpha) = \mu_\alpha$.

Conversely, clearly $\mu \subseteq \overleftarrow{\sigma}\sigma(\mu)$. Let $\alpha = \overleftarrow{\sigma}\sigma(\mu)(x) = \sup\{\mu(y) : (y]_{\circ\circ} = (x]_{\circ\circ}\}$. Then for each $\epsilon > 0$, there is $a \in L, (a]_{\circ\circ} = (x]_{\circ\circ}$ such that $\mu(a) \geq \alpha - \epsilon$. Since ϵ is arbitrary, $\mu(a) \geq \alpha$ such that $(a]_{\circ\circ} = (x]_{\circ\circ}$. Thus $a \in \mu_\alpha$. So $x \in \overleftarrow{\sigma}\sigma(\mu_\alpha) = \mu_\alpha$. Hence $\mu(x) \geq \alpha = \overleftarrow{\sigma}\sigma(\mu)(x)$. Therefore $\mu = \overleftarrow{\sigma}\sigma(\mu)$. \square

Theorem 4.10. For a nonempty subset I of L , I is a closure ideal if and only if χ_I is closure fuzzy ideal of L .

Theorem 4.11. Let μ be fuzzy ideal of an MS-algebra L . Then the following conditions are equivalent:

- (1) μ is a closure fuzzy ideal,
- (2) For all $x, y \in L, (x]_{\circ\circ} = (y]_{\circ\circ}$ implies $\mu(x) = \mu(y)$,
- (3) $\mu = \mu_{\circ\circ}$.

Proof. (1) \Rightarrow (2): Suppose μ is a closure fuzzy ideal. Let $x, y \in L$ such that $(x]_{\circ\circ} = (y]_{\circ\circ}$. Then

$$\mu(x) = \overleftarrow{\sigma}\sigma(\mu)(x) = \sigma(\mu)((x]_{\circ\circ}) = \sigma(\mu)((y]_{\circ\circ}) = \overleftarrow{\sigma}\sigma(\mu)(y) = \mu(y).$$

(2) \Rightarrow (1): Suppose for all $x, y \in L, (x]_{\circ\circ} = (y]_{\circ\circ}$ implies $\mu(x) = \mu(y)$. Then

$$\overleftarrow{\sigma}\sigma(\mu)(x) = \sigma(\mu)((x]_{\circ\circ}) = \sup\{\mu(y) : (x]_{\circ\circ} = (y]_{\circ\circ}\} = \mu(x).$$

(1) \Rightarrow (3): Suppose μ is closure fuzzy ideal of L . By Lemma 3.2, $\mu \subseteq \mu_{\circ\circ}$. Let $\forall x \in L$. Then

$$\begin{aligned} \mu_{\circ\circ}(x) &= \sup\{\mu(y) : x \leq y^{\circ\circ}\} \\ &= \sup\{\mu(y) : (x]_{\circ\circ} \subseteq (y]_{\circ\circ}\} \\ &\leq \sigma(\mu)((x]_{\circ\circ}) = \overleftarrow{\sigma}\sigma(\mu)(x) \\ &= \mu(x). \end{aligned}$$

Thus $\mu_{\circ\circ} \subseteq \mu$. So $\mu_{\circ\circ} = \mu$.

(3) \Rightarrow (1): Suppose $\mu = \mu_{\circ\circ}$, for all $x \in L$. Then

$$\begin{aligned} \overleftarrow{\sigma}\sigma(\mu)(x) &= \sigma(\mu)((x]_{\circ\circ}) \\ &= \sup\{\mu(y) : (x]_{\circ\circ} = (y]_{\circ\circ}\} \\ &\leq \sup\{\mu(y) : x \leq y^{\circ\circ}\} \\ &= \mu_{\circ\circ}(x) = \mu(x). \end{aligned}$$

It is clear that $\mu \subseteq \overleftarrow{\sigma}\sigma(\mu)$. Thus $\mu = \overleftarrow{\sigma}\sigma(\mu)$. By transitivity, (2) \Rightarrow (3) \Rightarrow (2). \square

Lemma 4.12. The following conditions are held in any MS-algebra L :

- (1) for any fuzzy ideal μ of $L, \overleftarrow{\sigma}\sigma(\mu) = \mu_{\circ\circ}$,
- (2) for any fuzzy ideal μ of $L, \mu_{\circ\circ}$ is a closure fuzzy ideal,
- (3) the map $\overleftarrow{\sigma}\sigma : FI(L) \rightarrow FI(L)$ is a $(0,1)$ -homomorphism.

Proof. (1) $\overleftarrow{\sigma}\sigma(\mu)(x) = \sup\{\mu(y) : (y]_{\circ\circ} = (x]_{\circ\circ}\} \leq \sup\{\mu(y) : x \leq y^{\circ\circ}\} = \mu_{\circ\circ}(x)$, for all $x \in L$. On the other hand, for all $x \in L$,

$$\begin{aligned} \mu_{\circ\circ}(x) &= \sup\{\mu(y) : x \leq y^{\circ\circ}\} \\ &= \sup\{\mu(y) : (x]_{\circ\circ} \subseteq (y^{\circ\circ}] = (y]_{\circ\circ}\} \\ &= \sigma((x]_{\circ\circ}) \\ &= \overleftarrow{\sigma}\sigma(\mu)(x). \end{aligned}$$

Then $\overleftarrow{\sigma}\sigma(\mu) = \mu_{\circ\circ}$.

(2) From (1) $\overleftarrow{\sigma}\sigma(\mu_{\circ\circ}) = \{\mu_{\circ\circ}\}_{\circ\circ}$. By Lemma 3.2 (3), $\{\mu_{\circ\circ}\}_{\circ\circ} = \mu_{\circ\circ}$. Then $\overleftarrow{\sigma}\sigma(\mu_{\circ\circ}) = \mu_{\circ\circ}$. Thus $\mu_{\circ\circ}$ is a closure fuzzy ideal of L .

(3) Clearly, $\overleftarrow{\sigma}\sigma(\chi_{\{0\}}) = \chi_{\{0\}}$ and $\overleftarrow{\sigma}\sigma(\chi_{\{L\}}) = \chi_{\{L\}}$. Let $\mu, \theta \in FI(L)$. Then by (2) and Lemma 3.3 (2),(3), we get

$$\overleftarrow{\sigma}\sigma(\mu \cap \theta) = (\mu \cap \theta)_{\circ\circ} = \mu_{\circ\circ} \cap \theta_{\circ\circ} = \overleftarrow{\sigma}\sigma(\mu) \cap \overleftarrow{\sigma}\sigma(\theta)$$

and

$$\overleftarrow{\sigma}\sigma(\mu \vee \theta) = (\mu \vee \theta)_{\circ\circ} = \mu_{\circ\circ} \vee \theta_{\circ\circ} = \overleftarrow{\sigma}\sigma(\mu) \vee \overleftarrow{\sigma}\sigma(\theta).$$

Thus $\overleftarrow{\sigma}\sigma$ is a (0,1)-homomorphism. \square

Now, we denote the set of all closure fuzzy ideals of L by $FI_C(L)$. Then we have the following theorem

Theorem 4.13. *Let L be an MS-algebra, the set $FI_C(L)$ forms a bounded distributive lattice.*

Definition 4.14. A proper closure fuzzy ideal μ of L is called prime closure fuzzy ideal, if for any two fuzzy ideals η, ν of L , $\eta \cap \nu \subseteq \mu$ implies $\eta \subseteq \mu$ or $\nu \subseteq \mu$.

Example 4.15. In Example 4.8, clearly $A = \{0\}$, $B = \{0, a\}$, $C = \{0, a, b\}$, $D = \{0, a, c\}$, $E = \{0, a, b, c, d\}$, $F = \{0, a, b, c, d, e\}$ are ideals of L and A, E and F are prime ideals of L . By the Theorem 2.16, A_α^1, E_α^1 and F_α^1 are prime fuzzy ideals of L , for any $\alpha \in [0, 1)$. Now, we prove that A_α^1, E_α^1 and F_α^1 are prime closure fuzzy ideals or not. We have:

$$\overleftarrow{\sigma}\sigma(A_\alpha^1)(0) = \sup\{A_\alpha^1(x) : (x]_{\circ\circ} = (0]_{\circ\circ}, x \in L\} = 1 = A_\alpha^1(0)$$

and for any $x \in L, x \neq 0$,

$$\overleftarrow{\sigma}\sigma(A_\alpha^1)(x) = \sup\{A_\alpha^1(y) : (x]_{\circ\circ} = (y]_{\circ\circ}, y \in L\} = \alpha = A_\alpha^1(x).$$

Then $\overleftarrow{\sigma}\sigma(A_\alpha^1) = A_\alpha^1$. Since $(e]_{\circ\circ} = (a]_{\circ\circ} = (b]_{\circ\circ} = (c]_{\circ\circ} = (d]_{\circ\circ}$ and $e \notin E$,

$$\begin{aligned} \overleftarrow{\sigma}\sigma(E_\alpha^1)(e) &= \sup\{E_\alpha^1(y) : (e]_{\circ\circ} = (y]_{\circ\circ}, y \in L\} \\ &= \sup\{E_\alpha^1(a), E_\alpha^1(b), E_\alpha^1(c), E_\alpha^1(d), E_\alpha^1(e)\} \\ &= \sup\{1, 1, 1, 1, \alpha\} \\ &= 1 \\ &\neq \alpha \\ &= E_\alpha^1(e). \end{aligned}$$

Thus $\overleftarrow{\sigma}\sigma(E_\alpha^1) \neq E_\alpha^1$. Similarly, $\overleftarrow{\sigma}\sigma(F_\alpha^1) = F_\alpha^1$. So A_α^1 and F_α^1 are prime closure fuzzy ideals. Hence E_α^1 is prime fuzzy ideal of L but not prime closure fuzzy ideal of L .

Theorem 4.16. *Let L be an MS-algebra. Then there is an isomorphism of the lattice of closure fuzzy ideals of L onto the lattice of fuzzy ideals on $M_{\circ\circ}(L)$. Under*

this isomorphism the prime closure fuzzy ideals of L corresponding to prime fuzzy ideals of $M_{oo}(L)$.

Proof. Define the mapping $h : FI_C(L) \rightarrow FI(M_{oo}(L))$ by $h(\mu) = \sigma(\mu), \forall \mu \in FI_C(L)$. Obviously, $h(\chi_{\{0\}}) = \chi_{\{0\}}$ and $h(\chi_L) = \sigma(\chi_L) = \chi_{M_{oo}(L)}$. Let μ and θ be any two closure fuzzy ideals of L . Since σ is a homomorphism, we have get

$$h(\mu \cap \theta) = \sigma(\mu \cap \theta) = \sigma(\mu) \cap \sigma(\theta) = h(\mu) \cap h(\theta)$$

and

$$h(\mu \vee \theta) = \sigma(\mu \vee \theta) = \sigma(\mu) \vee \sigma(\theta) = h(\mu) \vee h(\theta).$$

Then h is a $(0, 1)$ -lattice homomorphism. Let $\mu, \theta \in FI_C(L)$ such that $h(\mu) = h(\theta)$. Then $\sigma(\mu) = \sigma(\theta)$. Thus $\mu = \overleftarrow{\sigma} \sigma(\mu) = \overleftarrow{\sigma} \sigma(\theta) = \theta$. So h is one to one.

Now we prove that h is an onto. Let $\mu \in FI(M_{oo}(L))$. Then $\overleftarrow{\sigma}(\mu)$ is fuzzy ideal of L , by Lemma 4.3 (1) and by Lemma 4.3 (3), $\sigma \overleftarrow{\sigma}(\mu) = \mu$. Thus $\overleftarrow{\sigma} \{ \sigma \overleftarrow{\sigma}(\mu) \} = \overleftarrow{\sigma}(\mu)$. So $\overleftarrow{\sigma}(\mu) \in FI_C(L)$. Hence h is onto. Therefore h is isomorphism.

Also we have obtained one-to-one correspondence between prime closure fuzzy ideals of L and the prime fuzzy ideals of $M_{oo}(L)$. Let μ be any prime closure fuzzy ideal of L . We need to prove that $\sigma(\mu)$ is prime ideal of $M_{oo}(L)$. Let θ, ν are any ideals of $M_{oo}(L)$ such that $\theta \cap \nu \subseteq \sigma(\mu)$. Since h is onto, there exist a closure fuzzy ideals of L , θ_1, ν_1 such that $\sigma(\theta_1) = \theta$ and $\sigma(\nu_1) = \nu$. Now $\sigma(\theta_1) \cap \sigma(\nu_1) \subseteq \sigma(\mu)$. Then $\sigma(\theta_1 \cap \nu_1) \subseteq \sigma(\mu)$. Thus $\theta_1 \cap \nu_1 \subseteq \mu$. So $\theta_1 \subseteq \mu$ or $\nu_1 \subseteq \mu$. Hence $\sigma(\theta_1) \subseteq \sigma(\mu)$ or $\sigma(\nu_1) \subseteq \sigma(\mu)$. Therefore $\sigma(\mu)$ is prime fuzzy ideal of L .

Conversely, let μ be any prime ideal of $M_{oo}(L)$. Then there exists any closure fuzzy ideal μ_1 of L such that $\sigma(\mu_1) = \mu$. Let ϕ, ξ any fuzzy ideals of L such that $\phi \cap \xi \subseteq \mu_1$. Then $\sigma(\phi \cap \xi) = \sigma(\phi) \cap \sigma(\xi) \subseteq \sigma(\mu_1)$. Since $\sigma(\mu_1)$ is prime ideal of L , $\sigma(\phi) \subseteq \sigma(\mu_1)$ or $\sigma(\xi) \subseteq \sigma(\mu_1)$. Thus $\phi \subseteq \overleftarrow{\sigma} \sigma(\phi) \subseteq \overleftarrow{\sigma} \sigma(\mu_1) = \mu_1$ or $\xi \subseteq \overleftarrow{\sigma} \sigma(\xi) \subseteq \overleftarrow{\sigma} \sigma(\mu_1) = \mu_1$. So μ_1 is prime closure fuzzy ideal of L . Hence prime closure fuzzy ideals of L one to one correspondence to prime ideals of $M_{oo}(L)$. \square

5. CLOSURE FUZZY IDEALS AND HOMOMORPHISMS OF MS-ALGEBRAS

In this section, some properties of the homomorphic images and the inverse images of closure fuzzy ideals are observed. Homomorphism on an MS-algebra L , that means a lattice homomorphism f which preserves "°" such that $(f(x))^\circ = f(x^\circ)$, for all $x \in L$.

Lemma 5.1. *Let L and M be two MS-algebras and $f : L \rightarrow M$ be a homomorphism. Then we have the following:*

- (1) for any non empty fuzzy subset μ of L , $f(\mu_{oo}) \subseteq \{f(\mu)\}_{oo}$,
- (2) for any non empty fuzzy subset θ of M , $\{f^{-1}(\theta)\}_{oo} \subseteq f^{-1}(\theta_{oo})$.

Proof. (1) Let $y \in M$.

Case 1: Suppose $f^{-1}(y) = \emptyset$. Then $f(\mu_{oo})(y) = 0 \leq \{f(\mu)\}_{oo}(y)$.

Case 2: Suppose $f^{-1}(y) \neq \emptyset$. Then

$$\begin{aligned} f(\mu_{oo})(y) &= \sup\{\mu_{oo}(b) : f(b) = y, b \in L\} \\ &= \sup\{\sup\{\mu(a) : b \leq a^\circ, a \in L\} : f(b) = y, b \in L\} \\ &= \sup\{\mu(a) : b \leq a^\circ, f(b) = y, a, b \in L\} \end{aligned}$$

and

$$\begin{aligned} \{f(\mu)\}_{\circ\circ}(y) &= \sup\{f(\mu)(x) : y \leq x^{\circ\circ}, x \in M\} \\ &= \sup\{\sup\{\mu(a) : f(a) = x, a \in L\} : y \leq x^{\circ\circ}, x \in M\} \\ &= \sup\{\mu(a) : f(a) = x, y \leq x^{\circ\circ}, a \in L, x \in M\}. \end{aligned}$$

Let $A = \{\alpha, \text{ where } \mu(a) = \alpha, b \leq a^{\circ\circ}, f(b) = y, a, b \in L\}$ and let $B = \{\lambda, \text{ where } \lambda = \mu(a) : f(a) = x, y \leq x^{\circ\circ}, a \in L, x, y \in M\}$. We prove that $A \subseteq B$. Let $\alpha \in A$. Then $b \leq a^{\circ\circ}, f(b) = y, a, b \in L$. Thus $f(b) \leq f(a^{\circ\circ}) = (f(a))^{\circ\circ}, f(b) = y, a, b \in L$. So $y \leq (x)^{\circ\circ}, f(a) = x, a \in L, x \in M$. Hence $\alpha \in B$ and thus $A \subseteq B$. This implies $\sup A \leq \sup B$. Therefore $f(\mu_{\circ\circ}) \subseteq \{f(\mu)\}_{\circ\circ}$.

(2) Let $x \in L$. Then

$$\begin{aligned} \{f^{-1}(\theta)\}_{\circ\circ}(x) &= \sup\{f^{-1}(\theta)(a) : x \leq a^{\circ\circ}\} \\ &= \sup\{\theta(f(a)) : x \leq a^{\circ\circ}\} \\ &\leq \sup\{\theta(f(a)) : f(x) \leq (f(a))^{\circ\circ}\} \\ &= \theta_{\circ\circ}(f(x)) \\ &= f^{-1}(\theta_{\circ\circ})(x). \end{aligned}$$

Thus $\{f^{-1}(\theta)\}_{\circ\circ} \subseteq f^{-1}(\theta_{\circ\circ})$. □

Definition 5.2. Let $f : L \rightarrow M$ be a homomorphism of an MS-algebra L into an MS-algebra M . Then f is called dominator fuzzy ideal preserving, if $f(\mu_{\circ\circ}) = \{f(\mu)\}_{\circ\circ}$.

Theorem 5.3. Let $f : L \rightarrow M$ be a homomorphism of an MS-algebra L onto an MS-algebra M . Then f is a dominator fuzzy ideal preserving.

Proof. Let $x, y \in M$. Then

$$\begin{aligned} \{f(\mu)\}_{\circ\circ}(y) &= \sup\{f(\mu)(x) : y \leq x^{\circ\circ}\} \\ &= \sup\{\sup\{\mu(a) : f(a) = x, a \in L\} : y \leq x^{\circ\circ}\} \\ &= \sup\{\mu(a) : y \leq f(a^{\circ\circ})\} \\ &= f(\mu_{\circ\circ}). \end{aligned}$$

Thus f is dominator fuzzy ideal preserving. □

Theorem 5.4. Let $f : L \rightarrow M$ be a homomorphism of an MS-algebra L onto an MS-algebra M . Then for any closure fuzzy ideal μ of L , $f(\mu)$ is closure fuzzy ideal of M .

Proof. By Lemma 4.12 (2), Theorem 5.3 and Theorem 4.11 (3),

$$\overleftarrow{\sigma} \sigma(f(\mu)) = (f(\mu))_{\circ\circ} = f(\mu_{\circ\circ}) = f(\mu).$$

□

Theorem 5.5. Let L and M be two MS-algebras and $f : L \rightarrow M$ be a homomorphism. Then we have the following:

- (1) for any closure fuzzy ideal θ of M , $f^{-1}(\theta)$ is a closure fuzzy ideal of L ,
- (2) for any closure fuzzy ideal θ of M , $f^{-1}(\theta_{\circ\circ}) = \{f^{-1}(\theta)\}_{\circ\circ}$.

Proof. (1) Clearly, $f^{-1}(\theta) \subseteq \overleftarrow{\sigma} \sigma(f^{-1}(\theta))$. By Lemma 4.12 (2), Lemma 5.1 (2) and Theorem 4.11 (3), $\overleftarrow{\sigma} \sigma(f^{-1}(\theta)) = \{f^{-1}(\theta)\}_{\circ\circ} \subseteq f^{-1}(\theta_{\circ\circ}) = f^{-1}(\theta)$. Then we have $\overleftarrow{\sigma} \sigma(f^{-1}(\theta)) = f^{-1}(\theta)$.

(2) Since θ is closure fuzzy ideal of M . Then by (1), $f^{-1}(\theta)$ is closure fuzzy ideal of L . By Theorem 4.11 (3), $\{f^{-1}(\theta)\}_{\circ\circ} = f^{-1}(\theta) = f^{-1}(\theta_{\circ\circ})$. □

Theorem 5.6. *Let $f : L \rightarrow M$ be an onto homomorphism between MS-algebras L and M . Then $FI_C(L)$ is homomorphic of $FI_C(M)$.*

Proof. Define the map $h : FI_C(L) \rightarrow FI_C(M)$ by $h(\mu) = f(\mu)$, for all $\mu \in FI_C(L)$. It is clear that $h(\chi_{\{0_L\}}) = \chi_{\{0_L\}}$ and $h(\chi_L) = \chi_M$. Let $\mu, \theta \in FI_C(L)$. Then for all $y \in M$,

$$\begin{aligned} h(\mu \cap \theta)(y) &= f(\mu \cap \theta)(y) = \sup\{(\mu \cap \theta)(x) : f(x) = y\} \\ &= \sup\{\mu(x) \wedge \theta(x) : f(x) = y\} \\ &= \sup\{\mu(x) : f(x) = y\} \wedge \sup\{\theta(x) : f(x) = y\} \\ &= f(\mu)(y) \wedge f(\theta)(y) = h(\mu)(y) \wedge h(\theta)(y). \end{aligned}$$

Thus $h(\mu \cap \theta) = h(\mu) \cap h(\theta)$ and for any $y \in M$,

$$\begin{aligned} (h(\mu) \vee h(\theta))(y) &= (f(\mu) \vee f(\theta))(y) \\ &= \sup\{f(\mu)(a) \vee f(\theta)(b) : a \vee b = y, a, b \in M\} \\ &= \sup\{\sup\{\mu(a_1) : f(a_1) = a\} \vee \sup\{\theta(b_1) : f(b_1) = b\}, \\ &\quad a \vee b = y, a_1, b_1 \in L, a, b \in M\} \\ &= \sup\{\mu(a_1) \vee \theta(b_1) : f(a_1) \vee f(b_1) = f(a_1 \vee b_1) = y\} \\ &= \sup\{(\mu \vee \theta)(x) : x = a_1 \vee b_1 : f(x) = y\} \\ &= f(\mu \vee \theta)(y) = h(\mu \vee \theta)(y). \end{aligned}$$

So h is a homomorphism. □

Acknowledgements. The authors would like to thank the referees for their valuable comments and constructive suggestions.

REFERENCES

- [1] N. Ajmal, Fuzzy lattice, Inform. Sci. 79 (1994) 271–291.
- [2] M. Attallah, Completely fuzzy prime ideals of distributive lattices, The Journal of Fuzzy Mathematics 8 (1) (2000) 151–156.
- [3] E. Badawy and M. Sambasiva Rao, Closure ideals of MS-algebras, Chamchuri J. Math. 6 (2014) 31–46.
- [4] J. Berman, Distributive lattices with an additional unary operation, Aequationes Math. 16 (1977) 165–171.
- [5] T. S. Blyth and J. C. Varlet, Ockham Algebras, Oxford University Press 1994.
- [6] T. S. Blyth and J. C. Varlet, On a common abstraction of de Morgan and Stone algebras, Proc.Roy.Soc.Edinburgh Sect. A (1983) 94A: 301–308.
- [7] T. S. Blyth and J. C. Varlet, Subvarieties of the class of MS-algebras, Proc.Roy.Soc.Edinburgh Sect. A (1983) 95A: 157–169.
- [8] G. Gratzer, General lattice theory, New York, San Francisco, U S A: Academic Press 1978.
- [9] A. Kadji, C. Lele and M. Tonga, Fuzzy prime and maximal filters of residual lattice, Soft Comput, Springer-Verlag Berlin Heidelberg (2016) DOI 10.1007/s00500-016-2113-2.
- [10] B. B. N. Kogup, C. Nkuimi and C. Lele, On Fuzzy prime ideals of lattice, SJPAM 3 (2008) 1–11.
- [11] A. Rosenfeld, Fuzzy groups, J. Math. Anal. Appl. 35 (1971) 512–517.
- [12] M. Sambasiva Rao, β -filters of Ms-Algebras, Asian-Eur. J. Math. 5 (2012) 1–8.
- [13] M. Sambasiva Rao, e -filters of MS-algebras, Acta Mathematica Scientia 33 (2013) 738–746.
- [14] U. M. Swamy and D. V. Raju, Fuzzy ideals and congruences of lattices, Fuzzy sets and systems 95 (1998) 249–253.
- [15] U. M. Swamy and D. V. Raju, Irreducibility in algebraic fuzzy systems, Fuzzy Sets and Systems 41 (1991) 233–241.
- [16] Bo. Yuan and W. Wu, Fuzzy ideals on a distributive lattice, Fuzzy Sets and Systems 35 (1990) 231–240.

- [17] L. A. Zadeh, Fuzzy sets, Information and Control 8 (1965) 338–353.

BERHANU ASSAYE ALABA (berhanu_assaye@yahoo.com)

Department of Mathematics, Bahir Dar University, Bahir Dar, Ethiopia

TEFERI GETACHEW ALEMAYEHU (teferigetachew3@gmail.com)

Department of Mathematics, Debre Berhan University , Debre Berhan, Ethiopia