


β -fuzzy set and some relations on it

F. SALAMA



Reprinted from the
Annals of Fuzzy Mathematics and Informatics
Vol. 20, No. 3, December 2020

β -fuzzy set and some relations on it

F. SALAMA

Received 21 July 2020; Revised 2 August 2020; Accepted 21 August 2020

ABSTRACT. In this paper, we introduced a new extension of the concept of fuzzy set that will be given the name “ β -fuzzy set”. β -fuzzy t -norm, β -fuzzy t -conorm and β -fuzzy relations are defined. Some characteristics of β -fuzzy relations like symmetry, transitive and reflexive are intensively studied. Finally, the results on these characteristics are deduced.

2010 AMS Classification: 94D05, 08A72, 03E72, 20N25

Keywords: β -fuzzy Set, β -fuzzy t -norm, β -fuzzy t -conorm, β -fuzzy Relations.

Corresponding Author: F. Salama (fatma.salama@science.tanta.edu.eg)

1. INTRODUCTION

Zadeh [25], in 1965, introduced the notion of fuzzy set as an extension of a crisp set. In a crisp set, the characteristic function takes value from $\{0,1\}$, but every element in fuzzy set has grade of membership lies on the interval $[0,1]$. Fuzziness appears in many areas of daily life, for example in engineering, medicine, meteorology, manufacturing, control systems, modeling, signal processing and expert systems, for more details see [4, 5, 6, 8, 7, 9, 14, 15, 16].

Goguen [10], in 1967, generalized the work of Zadeh where initiated L -fuzzy set to develop a new point of view toward optimization problems. L might be a semigroup, a posset, a lattice, a Boolean σ -ring, or subintervals of the real numbers. Goguen in L -fuzzy set did not discuss the value that satisfied the counter-relation.

Atanassov [1], in 1986, instituted the concept of intuitionistic fuzzy set as a generalization of fuzzy set. An intuitionistic fuzzy set is a pair of mappings, one of them connect the element of intuitionistic fuzzy set with degree of membership in $[0,1]$ and any other element in intuitionistic fuzzy set has degree of non-membership in $[0,1]$ such that the sum of the degree of membership and the degree of non-membership for the same element equal 1. Recently, the intuitionistic fuzzy set has been discussed in [19, 24].

Zhang [28], in 1994, introduced an extension of fuzzy sets, the new concept called bipolar fuzzy sets. A bipolar fuzzy set is also a pair of mappings, a positive membership degree means the element satisfy the property corresponding the bipolar fuzzy set while the negative membership degree indicate that the element satisfy some implicit counter property corresponding the bipolar fuzzy set. In recent years, many researchers have studied bipolar fuzzy sets (See [29, 30, 31]).

In the present paper, we introduced a new extension of the concept of fuzzy set that will be given the name β -fuzzy set. A β -fuzzy set K of an universal set H is a map which connect every element in the set H with a membership degree belong to the interval $[-1, 1]$. An element x with a positive value of $\beta_K(x)$ indicates that it satisfies the property of β -fuzzy set. On the other hand, an element x with a negative value of $\beta_K(x)$ satisfies a counter-property and value means the element is not relevant to the corresponding property. β -fuzzy t -norm, a β -fuzzy t -conorm and β -fuzzy relations are defined. Some characteristics of β -fuzzy relations like symmetry, transitive and reflexive are intensively studied. Finally, the results on these characteristics are deduced.

2. PRELIMINARIES

In this section, we briefly recall some concepts on fuzzy set, fuzzy relation, intuitionistic fuzzy set and bipolar fuzzy set that are necessary for this paper.

Definition 2.1 ([27]). Let H be a universal set. Then a fuzzy set K in H (i.e., a fuzzy subset K of H) is characterized by a function of the form $\beta_K:H \rightarrow [0, 1]$ such a function β_K is called the membership function and for each $x \in H$, $\beta_K(x)$ is the degree of membership of x (membership grade of x) in the fuzzy set K .

Definition 2.2 ([17]). An L-fuzzy set A on a set X is a function $A:X \rightarrow L$, where L might be a semigroup, a poset, a lattice, a Boolean σ -ring, or subintervals of the real numbers.

Definition 2.3 ([1]). An intuitionistic fuzzy set K in a set H is defined as an object of the form $K = \{(x, \beta_K(x), \gamma_K(x)) : x \in H\}$, where $\beta_K:H \rightarrow [0, 1]$ and $\gamma_K:H \rightarrow [0, 1]$ define the degree of membership and the degree of non-membership of the element $x \in H$, respectively and for every $x \in H$, $0 \leq \beta_K(x) + \gamma_K(x) \leq 1$.

Definition 2.4 ([28]). Let H be a nonempty set. A bipolar fuzzy set K in H is an object having the form

$$K = \{(x, \beta_K^P(x), \beta_K^N(x)) : x \in H\},$$

where $\beta_K^P:H \rightarrow [0, 1]$ and $\beta_K^N:H \rightarrow [-1, 0]$ are two mappings. We use the positive membership degree $\beta_K^P(x)$ to denote the satisfaction degree of an element x to the property corresponding to a bipolar fuzzy set K , and the negative membership degree $\beta_K^N(x)$ to denote the satisfaction degree of an element x to some implicit counter-property corresponding to a bipolar fuzzy set K . If $\beta_K^P(x) \neq 0$ and $\beta_K^N(x) = 0$, it is the situation that x is regarded as having only positive satisfaction for K . If $\beta_K^P(x) = 0$ and $\beta_K^N(x) \neq 0$, it is the situation that x does not satisfy the property of K but somewhat satisfies the counter property of K . It is possible for an element K to be such that $\beta_K^P(x) \neq 0$ and $\beta_K^N(x) \neq 0$ when the membership function of the property overlaps that of its counter property over some portion of H .

Definition 2.5 ([2]). A map $\tau : H \times H \rightarrow [0, 1]$ is called a fuzzy relation on H , if

$$\tau(x, y) \leq K(x) \wedge K(y), \forall x, y \in H.$$

- (i) τ is symmetric, if $\tau(x, y) = \tau(y, x)$ for all $x, y \in H$.
- (ii) τ is transitive, if $\tau^2 \subseteq \tau$.
- (iii) τ is said to be reflexive on K , if $\tau(x, x) = K(x)$ for all $x \in H$.

For more details and background see [3, 11, 12, 13, 18, 20, 21, 22, 23, 26].

3. MAIN RESULTS

Firstly, we introduce a new extension of fuzzy set as follows:

Definition 3.1. β -fuzzy set K on the universal set H is a set of ordered pairs $K = \{(x, \beta_K(x)) : x \in H\}$ in which $\beta_K(x)$ is the grade of membership function of the element x in H , $\beta_K(x) : H \rightarrow [-1, 1]$. We can say, a β -fuzzy set K (shortly, β -FS) is a map that connect every element in the set H with a membership degree in $[-1, 1]$. Therefore, when $0 < \beta_K(x)$ means x satisfies the fuzziness property of a β -fuzzy set and if $\beta_K(x) < 0$ this means x satisfies a counter of fuzziness property. On the other hand, 0 value means the element x is not relevant to the corresponding property.

Remark 3.2. If $\beta_K(x) = 0$ and there exist an element y such that $x < y$ and $0 < \beta_K(y)$, then we write $\beta_K(x) = 0^+$. In the same way, $\beta_K(x) = 0^-$ means $\beta_K(x) = 0$ and there exist an element z , such that $z < x$ and $\beta_K(z) < 0$.

Definition 3.3. If K_1 and K_2 are two β -FSs of H , then the intersection of K_1 and K_2 is defined as

$$(K_1 \cap K_2)(x) = K_1(x) \wedge K_2(x) = \min\{K_1(x), K_2(x)\}.$$

Definition 3.4. The union of any two β -FSs K_1 and K_2 is β -fuzzy set given by

$$(K_1 \cup K_2)(x) = K_1(x) \vee K_2(x) = \max\{K_1(x), K_2(x)\}.$$

Example 3.5. Consider K_1 and K_2 are two β -FSs of a set $H = \{1, 2, 3, 4, 5\}$ defined as follows:

$$K_1 = \{(1, 0.2), (2, -0.5), (3, 0), (4, -0.1), (5, 0.3)\},$$

$$K_2 = \{(1, -0.3), (2, 0.4), (3, 1), (4, 0), (5, -0.2)\}.$$

Then we can easily have

$$K_1 \cap K_2 = \{(1, -0.3), (2, -0.5), (3, 0), (4, -0.1), (5, -0.2)\},$$

$$K_1 \cup K_2 = \{(1, 0.2), (2, 0.4), (3, 1), (4, 0), (5, 0.3)\}.$$

Next, we define β -fuzzy t -norm and β -fuzzy t -conorm as follows:

Definition 3.6. β -fuzzy t -norm is a function $T : [-1, 1] \times [-1, 1] \rightarrow [-1, 1]$ satisfies the following conditions:

- (i) $T(1, x) = x, \forall x \in [-1, 1]$,
- (ii) $T(x, y) = T(y, x), \forall x, y \in [-1, 1]$,
- (iii) $T(x, T(y, z)) = T(T(x, y), z), \forall x, y, z \in [-1, 1]$,
- (iv) $\forall x, y, z, w \in [-1, 1]$, if $x \leq y$ and $z \leq w$, then $T(x, z) \leq T(y, w)$.

Example 3.7. Consider the function $T: [-1, 1] \times [-1, 1] \rightarrow [-1, 1]$ defined as:

$$T(x, y) = x \wedge y \text{ for all } x, y \in [-1, 1].$$

Then we prove that T will satisfies the conditions of Definition 3.6.

(i) Let $x \in [-1, 1]$. Then we have easily

$$T(1, x) = 1 \wedge x = \min\{1, x\} = x.$$

(ii) Let $x, y \in [-1, 1]$. Then we have easily

$$T(x, y) = x \wedge y = \min\{x, y\} = \min\{y, x\} = y \wedge x = T(y, x).$$

(iii) Let $x, y, z \in [-1, 1]$. Then we have

$$\begin{aligned} T(x, T(y, z)) &= x \wedge T(y, z) = x \wedge (y \wedge z) \\ &= (x \wedge y) \wedge z = T(x, y) \wedge z \\ &= T(T(x, y), z). \end{aligned}$$

(iv) Let $x, y, z, w \in [-1, 1]$ such that $x \leq y, z \leq w$. Then we have

$$T(x, z) = x \wedge z = \min\{x, z\} \leq \min\{y, w\} = y \wedge w = T(y, w).$$

Thus $T(x, z) \leq T(y, w)$. So from (i)–(iv), T is a β -fuzzy t -norm.

Definition 3.8. A function $T^*: [-1, 1] \times [-1, 1] \rightarrow [-1, 1]$ is β -fuzzy t -conorm, if satisfies the following conditions:

(i) $T^*(-1, x) = x, \forall x \in [-1, 1]$,

(ii) $T^*(x, y) = T^*(y, x), \forall x, y \in [-1, 1]$,

(iii) $T^*(x, T^*(y, z)) = T^*(T^*(x, y), z), \forall x, y, z \in [-1, 1]$,

(iv) $\forall x, y, z, w \in [-1, 1]$, if $x \leq y$ and $z \leq w$, then $T^*(x, z) \leq T^*(y, w)$.

Example 3.9. Consider the function $T^*: [-1, 1] \times [-1, 1] \rightarrow [-1, 1]$ given by:

$$T^*(x, y) = x \vee y.$$

Then we check that T will satisfies the conditions of Definition 3.8.

(i) Let $x \in [-1, 1]$. Then we have easily

$$T^*(-1, x) = -1 \vee x = \max\{-1, x\} = x.$$

(ii) Let $x, y \in [-1, 1]$. Then we have easily

$$T^*(x, y) = x \vee y = \max\{x, y\} = \max\{y, x\} = y \vee x = T^*(y, x).$$

(iii) Let $x, y, z \in [-1, 1]$. Then we have

$$\begin{aligned} T^*(x, T^*(y, z)) &= x \vee T^*(y, z) = x \vee (y \vee z) \\ &= (x \vee y) \vee z = T^*(x, y) \vee z \\ &= T^*(T^*(x, y), z). \end{aligned}$$

(iv) Let $x, y, z, w \in [-1, 1]$ such that $x \leq y, z \leq w$. Then we have

$$T^*(x, z) = x \vee z = \max\{x, z\} \leq \max\{y, w\} = y \vee w = T^*(y, w).$$

Thus $T(x, z) \leq T(y, w)$. So So from (i)–(iv), T^* is a β -fuzzy t -conorm.

Definition 3.10. β -fuzzy complement is a function $C: [-1, 1] \rightarrow [-1, 1]$ satisfies the following conditions:

(i) $C(-1) = 1, C(1) = -1$,

(ii) if $x \leq y$, then $C(x) \geq C(y) \forall x, y \in [-1, 1]$.

Example 3.11. Consider the function $C: [-1, 1] \rightarrow [-1, 1]$ defined by:

$$C(x) = -x, \forall x \in [-1, 1].$$

Then we show that C will satisfies the conditions of Definition 3.10.

(i) Clearly, $C(-1) = -(-1) = 1, C(1) = -1.$

(ii) Let $x, y \in [-1, 1]$ such that $x \leq y.$ Then $C(x) = -x \geq -y = C(y).$ Thus $C(x) \geq C(y).$ So from (i) and (ii), C is β -fuzzy complement.

Definition 3.12. Let K be a β -FS on a set H and let τ be the relation defined by:

$$\tau : H \times H \rightarrow [-1, 1].$$

Then τ is a β -fuzzy relation on $K,$ if $\tau(x, y) \leq K(x) \wedge K(y), \forall x, y \in H.$

Example 3.13. Given a universal set $H = \{x, y, z\},$ a β -PFS is given by

$$K(x) = -0.1, K(y) = 0.7, K(z) = 0,$$

Define a relation $\tau : H \times H \rightarrow [-1, 1]$ as follows:

$\tau(x, x) = -0.2$	$\tau(y, x) = -0.3,$	$\tau(z, x) = -0.5,$
$\tau(x, y) = -0.1$	$\tau(y, y) = 0.5,$	$\tau(z, y) = -0.2,$
$\tau(x, z) = -0.5$	$\tau(y, z) = 0.2,$	$\tau(z, z) = -0.9.$

Since the given relation τ satisfy the condition $\tau(x, y) \leq K(x) \wedge K(y), \forall x, y \in H.$ Then τ is a β -fuzzy relation (shortly, β -FR) on $K.$

Definition 3.14. Let K_1 and K_2 be β -FSs on sets H_1 and $H_2,$ respectively. Then a β -FR from the β -FS K_1 into the β -FS K_2 is a β -FS τ of $H_1 \times H_2$ defined by $\tau : H_1 \times H_2 \rightarrow [-1, 1]$ such that $\tau(x, y) \leq K_1(x) \wedge K_2(y), \forall x \in H_1, y \in H_2.$

Definition 3.15. A strong β -FR τ_K on a β -FS K of a set H is a β -FR defined by $\tau(x, y) = K(x) \wedge K(y), \forall x, y \in H.$

Example 3.16. Let H be a universal set given by $H = \{x, y, z\},$ a β -FS K is defined as:

$$K(x) = -0.2, K(y) = 0.6, K(z) = 0.$$

A relation $\tau : H \times H \rightarrow [-1, 1]$ is given by:

$\tau(x, x) = -0.2,$	$\tau(y, x) = -0.2,$	$\tau(z, x) = -0.2,$
$\tau(x, y) = -0.2,$	$\tau(y, y) = 0.6,$	$\tau(z, y) = 0,$
$\tau(x, z) = -0.2,$	$\tau(y, z) = 0,$	$\tau(z, z) = 0,$

Then it is clear that $\tau(x, y) = K(x) \wedge K(y), \forall x, y \in H.$ Thus τ is a strong β -FR on $K,$ i.e., $\tau_K = \tau.$

Definition 3.17. Let $\tau_1 : H_1 \times H_2 \rightarrow [-1, 1]$ be a β -FR from a β -fuzzy set K_1 of H_1 into a β -FS K_2 of H_2 and $\tau_2 : H_2 \times H_3 \rightarrow [-1, 1]$ is a β -FR from a β -FS K_2 of H_2 into a β -FS K_3 of $H_3.$ Then, a composition of τ_1 and $\tau_2, \tau_1 \circ \tau_2 : H_1 \times H_3 \rightarrow [-1, 1],$ is defined by $\tau_1 \circ \tau_2(x, z) = \vee \{\tau_1(x, y) \wedge \tau_2(y, z) : y \in H_2\}$ for all $x \in H_1$ and $z \in H_3.$

In the following we denote by $\beta\text{-FR}(H)$ the family of all β -FRs on a β -FS K of a universal set H through our study of operations on β -FRs.

Proposition 3.18. *Let $\tau_1, \tau_2 \in \beta - FR(H)$. Then a composition $\tau_1 \circ \tau_2$ is also belong to $\beta - FR(H)$.*

Proof. Let $x, y, z \in H$. Since $\tau_1, \tau_2 \in \beta - FR(H)$, we have

$$\tau_1(x, y) \leq K(x) \wedge K(y) \text{ and } \tau_2(y, z) \leq K(y) \wedge K(z).$$

Thus $\tau_1(x, y) \wedge \tau_2(y, z) \leq K(x) \wedge K(y) \wedge K(z) \leq K(x) \wedge K(z)$. So we have

$$\tau_1 \circ \tau_2(x, z) = \vee \{ \tau_1(x, y) \wedge \tau_2(y, z) : y \in H \} \leq K(x) \wedge K(z).$$

Hence the composition of τ_1 and τ_2 is a $\beta - FR$ on K . □

Proposition 3.19. *If $\tau_1, \tau_2, \tau_3 \in \beta - FR(H)$, then $(\tau_1 \circ \tau_2) \circ \tau_3$ and $\tau_1 \circ (\tau_2 \circ \tau_3)$ are also $\beta - FR$ s on K .*

Proof. Since $\tau_1, \tau_2, \tau_3 \in \beta - FR(H)$, from Proposition 3.18, $\tau_2 \circ \tau_3 \in \beta - FR(H)$. Since τ_1 and $\tau_2 \circ \tau_3$ are two $\beta - FR$ s on K , from Proposition 3.18, the relation $\tau_1 \circ (\tau_2 \circ \tau_3)$ is also $\beta - FR$ on K . By the same manner, $(\tau_1 \circ \tau_2) \circ \tau_3$ is a $\beta - FR$ on K . □

Proposition 3.20. *If $\tau_1, \tau_2, \tau_3 \in \beta - FR(H)$, then the following relation holds:*

$$(\tau_1 \circ \tau_2) \circ \tau_3 = \tau_1 \circ (\tau_2 \circ \tau_3).$$

Proof. Consider $\tau_1, \tau_2, \tau_3 \in \beta - FR(H)$. Then the $\beta - FR$ s $\tau_1 \circ \tau_2$ and $\tau_2 \circ \tau_3$ are defined by: for all $x, z \in H$ and $y, w \in H$,

$$\tau_1 \circ \tau_2(x, z) = \vee \{ \tau_1(x, y) \wedge \tau_2(y, z) : y \in H \}$$

and

$$\tau_2 \circ \tau_3(y, w) = \vee \{ \tau_2(y, z) \wedge \tau_3(z, w) : z \in H \}.$$

Thus, a $\beta - FR$ $(\tau_1 \circ \tau_2) \circ \tau_3$ is defined as: for all $x, w \in H$,

$$\begin{aligned} [(\tau_1 \circ \tau_2) \circ \tau_3](x, w) &= \vee \{ \tau_1 \circ \tau_2(x, z) \wedge \tau_3(z, w) : z \in H \} \\ &= \vee \{ \vee \{ \tau_1(x, y) \wedge \tau_2(y, z) : y \in H \} \wedge \tau_3(z, w) : z \in H \} \\ (3.1) \qquad \qquad \qquad &= \vee \{ \tau_1(x, y) \wedge \tau_2(y, z) \wedge \tau_3(z, w) : y, z \in H \}. \end{aligned}$$

Also $\tau_1 \circ (\tau_2 \circ \tau_3)$ is defined by: for all $x, w \in H$,

$$\begin{aligned} [\tau_1 \circ (\tau_2 \circ \tau_3)](x, w) &= \vee \{ \tau_1(x, y) \wedge \tau_2 \circ \tau_3(y, w) : y \in H \} \\ &= \vee \{ \vee \{ \tau_2(y, z) \wedge \tau_3(z, w) : z \in H \} \wedge \tau_1(x, y) : y \in H \} \\ (3.2) \qquad \qquad \qquad &= \vee \{ \tau_1(x, y) \wedge \tau_2(y, z) \wedge \tau_3(z, w) : y, z \in H \}. \end{aligned}$$

From (3.1) and (3.2), we find $(\tau_1 \circ \tau_2) \circ \tau_3 = \tau_1 \circ (\tau_2 \circ \tau_3)$. □

Definition 3.21. A $\beta - FR$ τ on a $\beta - FS$ K of a set H is said to be a reflexive, if $\tau(x, x) = K(x)$ for all $x \in H$.

Theorem 3.22. *For any two $\beta - FR$ s τ_1 and τ_2 on $\beta - FR(H)$, if τ_1 is reflexive on K , then $\tau_2 \subseteq \tau_1 \circ \tau_2$ and $\tau_2 \subseteq \tau_2 \circ \tau_1$.*

Proof. Since $\tau_1, \tau_2 \in \beta\text{-FR}(H)$ and τ_1 is reflexive, $\tau_1(x, x) = K(x)$ for all $x \in H$ and $\tau_2(x, z) \leq K(x) \wedge K(z)$. Then from Definition 3.17, the composition of τ_1 and τ_2 is given by: for all $x, z \in H$,

$$\begin{aligned} \tau_1 \circ \tau_2(x, z) &= \bigvee \{ \tau_1(x, y) \wedge \tau_2(y, z) : y \in H \} \\ &\geq \tau_1(x, x) \wedge \tau_2(x, z) \\ &= K(x) \wedge \tau_2(x, z) = \tau_2(x, z). \end{aligned}$$

Thus $\tau_2 \subseteq \tau_1 \circ \tau_2$. By the same way, we can prove that $\tau_2 \subseteq \tau_2 \circ \tau_1$. □

Corollary 3.23. *If τ is a reflexive β -FR, then $\tau \subseteq \tau^2$.*

Proof. Since τ is a reflexive β -FR, from Theorem 3.22, $\tau \subseteq \tau \circ \tau = \tau^2$. □

Theorem 3.24. *Let τ be a reflexive β -FR on K . Then*

$$\tau(x, x) = \tau^2(x, x) = \dots = \tau^\infty(x, x) = k(x)$$

for all $x \in H$.

Proof. To prove this theorem, we will use mathematical induction as follows.

Let τ be a reflexive β -FR on a β -FS K of a set H . Then $\tau(x, x) = K(x)$ for all $x \in H$. Assume that the relation $\tau^n(x, x) = K(x)$ is true at $m = n$ for all $x \in H$. Now, we will prove that the relation $\tau^{n+1}(x, x) = K(x)$ is true at $m = n + 1$ for all $x \in H$ as follows:

$$\begin{aligned} \tau^{n+1}(x, x) &= \tau \circ \tau^n(x, x) \\ &= \bigvee \{ \tau(x, y) \wedge \tau^n(y, x) : y \in H \} \\ &\leq \bigvee \{ K(x) \wedge K(x) : y \in H \}. \end{aligned}$$

Then we have

$$(3.3) \quad \tau^{n+1}(x, x) \leq K(x).$$

Also,

$$\begin{aligned} \tau^{n+1}(x, x) &= \tau \circ \tau^n(x, x) \\ &= \bigvee \{ \tau(x, y) \wedge \tau^n(y, x) : y \in H \} \\ &\geq \tau(x, x) \wedge \tau^n(x, x) \\ &= K(x) \wedge K(x) = K(x). \end{aligned}$$

Thus we have

$$(3.4) \quad \tau^{n+1}(x, x) \geq K(x).$$

So from (3.3) and (3.4), we have $\tau^{n+1}(x, x) = K(x)$ for all $x \in H$. □

Theorem 3.25. *Let τ_1 and τ_2 be reflexive β -FRs on a β -FS K . Then $\tau_1 \circ \tau_2$ and $\tau_2 \circ \tau_1$ are reflexive.*

Proof. Suppose τ_1 and τ_2 are reflexive β -FRs. Then for all $x \in H$,

$$\begin{aligned} \tau_1 \circ \tau_2(x, x) &= \bigvee \{ \tau_1(x, y) \wedge \tau_2(y, x) : y \in H \} \\ &\leq \bigvee \{ K(x) \wedge K(x) : y \in H \} = K(x). \end{aligned}$$

Thus we have

$$(3.5) \quad \tau_1 \circ \tau_2(x, x) \leq K(x).$$

Also, for all $x \in H$,

$$\begin{aligned} \tau_1 \circ \tau_2(x, x) &= \bigvee \{ \tau_1(x, y) \wedge \tau_2(y, x) : y \in H \} \\ &\geq \tau_1(x, x) \wedge \tau_2(x, x) \\ &= K(x) \wedge K(x) = K(x). \end{aligned}$$

So we have

$$(3.6) \quad \tau_1 \circ \tau_2(x, x) \geq K(x).$$

Hence, from (3.5) and (3.6), we have $\tau_1 \circ \tau_2(x, x) = K(x)$ for all $x \in H$. This means that $\tau_1 \circ \tau_2$ is reflexive.

Similarly, we can prove $\tau_2 \circ \tau_1$ is also reflexive. □

Definition 3.26. A β -FR τ on a β -fuzzy set K of a set H is symmetric, if $\tau(x, y) = \tau(y, x)$ for all $x, y \in H$.

Theorem 3.27. Let τ_1 and τ_2 be a symmetric β -FRs on K , then $\tau_1 \circ \tau_2$ is symmetric if and only if $\tau_1 \circ \tau_2 = \tau_2 \circ \tau_1$.

Proof. Suppose $\tau_1 \circ \tau_2$ is symmetric and let $x, z \in H$. Then clearly, $\tau_1 \circ \tau_2(x, z) = \tau_1 \circ \tau_2(z, x)$. Thus we have

$$\begin{aligned} \tau_1 \circ \tau_2(x, z) &= \bigvee \{ \tau_1(x, y) \wedge \tau_2(y, z) : y \in H \} \\ &= \bigvee \{ \tau_1(y, x) \wedge \tau_2(z, y) : y \in H \} \\ &= \bigvee \{ \tau_2(z, y) \wedge \tau_1(y, x) : y \in H \} \\ &= \tau_2 \circ \tau_1(z, x). \end{aligned}$$

Since $\tau_1 \circ \tau_2(x, z) = \tau_1 \circ \tau_2(z, x)$, $\tau_1 \circ \tau_2(z, x) = \tau_2 \circ \tau_1(z, x)$. So $\tau_1 \circ \tau_2 = \tau_2 \circ \tau_1$.

Conversely, suppose $\tau_1 \circ \tau_2 = \tau_2 \circ \tau_1$ and let $x, z \in H$. Then $\tau_1 \circ \tau_2(z, x) = \tau_2 \circ \tau_1(z, x)$. Thus we have

$$\begin{aligned} \tau_1 \circ \tau_2(x, z) &= \bigvee \{ \tau_1(x, y) \wedge \tau_2(y, z) : y \in H \} \\ &= \bigvee \{ \tau_1(y, x) \wedge \tau_2(z, y) : y \in H \} \\ &= \bigvee \{ \tau_2(z, y) \wedge \tau_1(y, x) : y \in H \} \\ &= \tau_2 \circ \tau_1(z, x) \\ &= \tau_1 \circ \tau_2(z, x). \end{aligned}$$

So $\tau_1 \circ \tau_2(x, z)$ is symmetric. □

Theorem 3.28. Let τ be a symmetric β -FR on a β -fuzzy set K of a set H . Then every power of τ is also symmetric.

Proof. If τ is a symmetric β -FR, then from Theorem 3.27, $\tau^2 = \tau \circ \tau$ is symmetric. Assume that τ^n is symmetric at $m = n$ and let $x, y \in H$. Then we have

$$\begin{aligned} \tau^{n+1}(x, z) &= \tau \circ \tau^n(x, z) \\ &= \bigvee \{ \tau(x, y) \wedge \tau^n(y, z) : y \in H \} \\ &= \bigvee \{ \tau^n(z, y) \wedge \tau(y, x) : y \in H \} \\ &= \tau^n \circ \tau(z, x) \\ &= \tau^{n+1}(z, x). \end{aligned}$$

Thus τ^{n+1} is symmetric. □

Definition 3.29. A β -FR τ on a β -FS K of a set H is transitive, if $\tau^2 \subseteq \tau$.

Theorem 3.30. If $\tau_1, \tau_2, \tau_3, \tau_4 \in \beta - FR(H)$, $\tau_1 \subseteq \tau_2$ and $\tau_3 \subseteq \tau_4$. Then $\tau_1 \circ \tau_3 \subseteq \tau_2 \circ \tau_4$.

Proof. Let $\tau_1 \subseteq \tau_2$ and $\tau_3 \subseteq \tau_4$, and let $x, y, z \in H$. Then $\tau_1(x, y) \leq \tau_2(x, y)$ and $\tau_3(y, z) \leq \tau_4(y, z)$. Thus we have

$$\begin{aligned} \tau_1 \circ \tau_3(x, z) &= \bigvee \{ \tau_1(x, y) \wedge \tau_3(y, z) : y \in H \} \\ &\leq \bigvee \{ \tau_2(x, y) \wedge \tau_4(y, z) : y \in H \} \\ &= \tau_2 \circ \tau_4(x, z). \end{aligned}$$

So $\tau_1 \circ \tau_3 \subseteq \tau_2 \circ \tau_4$. □

Theorem 3.31. Let τ_1 and τ_2 be transitive β -FRs on a β -fuzzy set K of a set H and $\tau_1 \circ \tau_2 = \tau_2 \circ \tau_1$. Then $\tau_1 \circ \tau_2$ is transitive.

Proof. Since τ_1 and τ_2 are transitive, $\tau_1^2 \subseteq \tau_1$ and $\tau_2^2 \subseteq \tau_2$. Then from Theorem 3.30, we have $\tau_1^2 \circ \tau_2^2 \subseteq \tau_1 \circ \tau_2$. Thus

$$\begin{aligned} (\tau_1 \circ \tau_2)^2 &= (\tau_1 \circ \tau_2) \circ (\tau_1 \circ \tau_2). \\ &= \tau_1 \circ (\tau_2 \circ \tau_1) \circ \tau_2 \\ &= \tau_1 \circ (\tau_1 \circ \tau_2) \circ \tau_2 \\ &= (\tau_1 \circ \tau_1) \circ (\tau_2 \circ \tau_2) \\ &= \tau_1^2 \circ \tau_2^2 \subseteq \tau_1 \circ \tau_2. \end{aligned}$$

So $\tau_1 \circ \tau_2$ is transitive. □

Theorem 3.32. Let τ be a symmetric and a transitive β -FR on K , where K is a β -FS of a set H . Then $\tau(x, y) \leq \tau(x, x)$ and $\tau(y, x) \leq \tau(x, x)$ for all $x, y \in H$.

Proof. Consider τ is a symmetric and transitive β -FR on K . Then $\tau \circ \tau \subseteq \tau$, i.e., $\tau \circ \tau(x, x) \leq \tau(x, x)$ for all $x \in H$. Thus

$$\begin{aligned} \tau \circ \tau(x, x) &= \bigvee \{ \tau(x, y) \wedge \tau(y, x) : y \in H \} \\ &= \bigvee \{ \tau(x, y) \wedge \tau(x, y) : y \in H \} \\ &= \tau(x, y). \end{aligned}$$

So $\tau \circ \tau(x, x) = \tau(x, y) \leq \tau(x, x)$. Hence $\tau(x, y) \leq \tau(x, x)$ for all $x, y \in H$. Since τ is symmetric, $\tau(y, x) \leq \tau(x, x)$ for all $x, y \in H$. □

Theorem 3.33. Let $\tau_1, \tau_2, \tau_3 \in \beta - FR(H)$. If τ_1 is transitive and $\tau_2 \subseteq \tau_1, \tau_3 \subseteq \tau_1$, then $\tau_2 \circ \tau_3 \subseteq \tau_1$.

Proof. Since $\tau_2 \subseteq \tau_1$ and $\tau_3 \subseteq \tau_1$, from Theorem 3.30, $\tau_2 \circ \tau_3 \subseteq \tau_1^2$. Since τ_1 is transitive, $\tau_1^2 \subseteq \tau_1$. Then $\tau_2 \circ \tau_3 \subseteq \tau_1$. \square

Theorem 3.34. If τ is transitive β -FR on a β -FS K of a set H , then every power of τ is also transitive.

Theorem 3.35. Let τ_1 be a transitive β -FR on a β -FS of a set H , let τ_2 be a reflexive β -fuzzy relations on K and let $\tau_2 \subseteq \tau_1$. Then $\tau_1 \circ \tau_2 = \tau_2 \circ \tau_1 = \tau_1$.

Proof. Since τ_1 is a transitive and $\tau_2 \subseteq \tau_1$, from Theorem 3.33, $\tau_1 \circ \tau_2 \subseteq \tau_1$. Then for all $x, z \in H$,

$$\begin{aligned} \tau_1 \circ \tau_2(x, z) &= \bigvee \{ \tau_1(x, y) \wedge \tau_2(y, z) : y \in H \} \\ &\geq \tau_1(x, z) \wedge \tau_2(z, z) \\ &= \tau_1(x, z) \wedge K(z) = \tau_1(x, z) \end{aligned}$$

Thus $\tau_1 \subseteq \tau_1 \circ \tau_2$. So $\tau_1 \circ \tau_2 = \tau_1$.

By the same way, we can prove that $\tau_2 \circ \tau_1 = \tau_1$. \square

In the following, we introduce special type of intersection and of β -fuzzy set.

Definition 3.36. For any two β -FSs K_1 and K_2 of a set H , an F-intersection of K_1 and K_2 , denoted by $K_1 \cap^F K_2$, is defined as: for each $x \in H$,

$$\begin{aligned} (K_1 \cap^F K_2)(x) &= K_2(x) \wedge^F K_1(x) \\ &= \begin{cases} \min \{K_1(x), K_2(x)\} & \text{if } K_1(x) \text{ and } K_2(x) \text{ have the same sign} \\ 0 & \text{if } K_1(x) \text{ and } K_2(x) \text{ have different sign} \\ 0 & \text{if } K_1(x) = 0, \text{ or } K_2(x) = 0. \end{cases} \end{aligned}$$

Definition 3.37. Let K_1 and K_2 are two β -FSs of H , then an F- union of K_1 and K_2 , denoted by $K_1 \cup^F K_2$, is defined as: for each $x \in H$,

$$\begin{aligned} (K_1 \cup^F K_2)(x) &= K_2(x) \vee^F K_1(x) \\ &= \begin{cases} \max \{K_1(x), K_2(x)\} & \text{if } K_1(x) \text{ and } K_2(x) \text{ have the same sign} \\ 0 & \text{if } K_1(x) \text{ and } K_2(x) \text{ have different sign} \\ 0 & \text{if } K_1(x) = 0, \text{ or } K_2(x) = 0. \end{cases} \end{aligned}$$

Example 3.38. Let $H = \{x_1, x_2, x_3, x_4, x_5\}$ be a universal set and let K_1 and K_2 be the β -FSs given by:

$$K_1 = \{(x_1, 0.7), (x_2, 0.5), (x_3, 0), (x_4, -0.1), (x_5, 0.3)\},$$

$$K_2 = \{(x_1, -0.3), (x_2, 0.4), (x_3, 1), (x_4, -0.2), (x_5, -0.2)\}.$$

Then clearly,

$$K_1 \cap^F K_2 = \{(x_1, 0), (x_2, 0.4), (x_3, 0), (x_4, -0.2), (x_5, 0)\},$$

$$K_1 \cup^F K_2 = \{(x_1, 0), (x_2, 0.5), (x_3, 0), (x_4, -0.1), (x_5, 0)\}.$$

Now, a new concept of composition of β -FRs is introduced as follows.

Definition 3.39. Let τ_1 be a β -FR from a β -FS K_1 of H_1 into a β -FS K_2 of H_2 , i.e., $\tau_1 : H_1 \times H_2 \rightarrow [-1, 1]$ and τ_2 is a β -FR from a β -FS K_2 of H_2 into a β -FS K_3 of H_3 , $\tau_2 : H_2 \times H_3 \rightarrow [-1, 1]$. Then an F -composition of τ_1 and τ_2 , $\tau_1 \circ^F \tau_2 : H_1 \times H_3 \rightarrow [-1, 1]$ is defined by: for all $x \in H_1$ and $z \in H_3$,

$$\tau_1 \circ^F \tau_2 (x, z) = \vee \{ |\tau_1(x, y) \wedge^F \tau_2(y, z)| : y \in H_2 \}.$$

Example 3.40. Let P be a set of patients, let $S = \{s_1, s_2, s_3, s_4, s_5\}$ be a set of symptoms and let N be a set of illnesses. If τ_1 is a β -FR from P to S defined as:

- $\tau_1(p, s) > 0$ is the degree that a symptoms appear on a patient p ,
- $\tau_1(p, s) = 0$ is the degree that a symptoms is not appear on a patient p ,
- $\tau_1(p, s) < 0$ is the degree that a counter- symptoms appear on a patient p .

Let τ_2 be a β -FR from S to N defined by:

- $\tau_2(s, n) > 0$ is the degree that s is a symptom of illness n ,
- $\tau_2(s, n) = 0$ is the degree that s is not a symptom of illness n ,
- $\tau_2(s, n) < 0$ is the degree that s is a counter-symptom of illness n .

Let us consider p is a patient in P , n is illness in N and τ_1, τ_2 are given by:

τ_1	s_1	s_2	s_3	s_4	s_5
p	0	0.4	0.7	-0.6	-0.3

Table 1.

τ_2	n
s_1	0.7
s_2	0.1
s_3	-0.5
s_4	1
s_5	-0.2

Table 2.

Then we have

$$\tau_1 \circ^F \tau_2 (x, z) = \vee \{0, 0.1, 0, 0, 0.3\} = 0.3.$$

It is clear from the formula of the F -composition of τ_1 and τ_2 that two symptoms of illness n shown by patient p The proportion of his suffering from this disease is 8%.

4. CONCLUSION

We introduced a new extension of the concept of fuzzy set that was given the name β -fuzzy set. A β -fuzzy set K of H is a map which connect every element in the set H with a membership degree belong to the interval $[-1, 1]$. An element x with a positive value of $\beta_K(x)$ indicates that it satisfies the property of a perfect fuzzy set. On the other hand, an element x with a negative value of $\beta_K(x)$ satisfies a counter-property and 0 value means the element is not relevant to the corresponding property. A β -fuzzy t -norm, a β -fuzzy t -conorm and β -fuzzy relations are defined. Some characteristics of β -fuzzy relations like symmetry, transitive and reflexive are intensively studied. A β -fuzzy set is an extremely useful tool for solving many problems in different areas of human life.

Declarations:

Availability of data and materials: All the data in the manuscript are public.

Conflicts of Interest: The author declares no conflict of interest.

Acknowledgements: The author is deeply grateful to Chief Editor and referees for valuable comments and suggestions which made the paper more readable.

REFERENCES

- [1] K. T. Atanassov, Intuitionistic fuzzy sets, *Fuzzy Sets and Systems* 20 (1986) 87–96.
- [2] B. De Baets and E. E. Kerre, Fuzzy relations and applications, *Adv. Electron. Electron Phys.* 89 (1994) 255–324.
- [3] B. De Baets and E. E. Kerre, The cutting of compositions, *Fuzzy Sets and Systems* 62 (1994) 295–309.
- [4] R. Belohlave, T. Kuhr and V. Vychodil, Confluence and termination of fuzzy relations, *Inform. Sci.* 180 (2010) 3288–3303.
- [5] C. Bo and X. Zhang, New operations of picture fuzzy relations and fuzzy comprehensive evaluation, *Symmetry* 2017, 9, 268; doi:10.3390/sym9110268.
- [6] G. Deschrijver and E. E. Kerre, On the relationship between some extensions of fuzzy set theory, *Fuzzy Sets and Systems* 133 (2003) 227–235.
- [7] S. Díaz, B. De Baets and S. Montes, On the Ferrers property of valued interval orders, 19 (2011) 421–427.
- [8] S. Díaz, B. De Baets and S. Montes, Transitivity and negative transitivity in the fuzzy setting, *Adv. Intell. Soft Comput.* 117 (2012) 91–100.
- [9] D. Dubois and H. Prade, A class of fuzzy measures based on triangular norms, *Int. J. General Systems* 8 (1982) 43–61.
- [10] J. Goguen, L-fuzzy sets, *J. Math. Anal. Appl.* 18 (1967) 145–174.
- [11] S. Güzide, A Comparative research on the definition of soft point, *International Journal of Computer Applications* 163 (2) (2017) 1–4. Doi: 10.5120/ijca2017913471.
- [12] S. Güzide, Analyzing the locus of soft spheres: Illustrative cases and drawings, *European Journal of Pure and Applied Mathematics* 11 (4) (2018) 946–957. Doi:10.29020/nybg.ejpam.v11i4.3321.
- [13] S. Güzide, The relation between soft topological space and soft ditopological space, *Communications Series A1: Mathematics and Statistics* 67 (2) (2018) 209–219. Doi:10.1501/Commua1.0000000875.
- [14] Q. Hu, L. Zhang, S. An and D. Zhang, On robust fuzzy rough set models, *IEEE Trans. Fuzzy Syst.* 20 (2012) 636–651.
- [15] D. Martinetti, I. Montes, S. Díaz and S. Montes, A study on the transitivity of probabilistic and fuzzy relations, *Fuzzy Sets and Systems* 184 (2011) 156–170.
- [16] A. Pradera, G. Belia, H. Bustince and B. De Baets, A review of the relationships between implication, negation and aggregation functions from the point of view of material implication, *Inform. Sci.* 329 (2016) 357–380.
- [17] J. Qiao and B. Q. Hua, Granular variable precision L-fuzzy rough sets based on residuated lattices, *Fuzzy Sets and Systems* 336 (2018) 148–166.
- [18] B. Rosman and S. Ramamoorthy, Learning spatial relationships between objects, *Int. J. Robot. Res.* 30 (11) (2011) 1328–1342.
- [19] S. Sahoo and M. Pal, Intuitionistic fuzzy tolerance graphs with application, *J. Appl. Math. Comput.* 55 (2017) 495–511.
- [20] C. Y. Wang and B. Q. Hu, Granular variable precision fuzzy rough sets with general fuzzy relations, *Fuzzy Sets and Systems* 275 (2015) 39–57.
- [21] X. Wang, X. Cao, C. Wu and J. Chen, Indicators of fuzzy relations, *Fuzzy Sets and Systems* 216 (2013) 91–107.
- [22] S. Wibowo and S. Grandhi, Fuzzy multicriteria analysis for performance evaluation of internet of things based supply chains, *Symmetry* 2018, 10, 603; doi:10.3390/sym10110603.
- [23] R. R. Yager, Fuzzy relations between Dempster–Shafer belief structures, *Knowledge-Based Systems* 105 (2016) 60–67.

- [24] K. Yin, P. Wang and X. Jin, Dynamic intuitionistic fuzzy multi-attribute group decision-making based on power geometric weighted average operator and prediction Model, *Symmetry* 2018, 10, 536; doi:10.3390/sym10110536m
- [25] L. A. Zadeh, Fuzzy sets, *Information and Control* 8 (1965) 338–353.
- [26] L.A. Zadeh, Similarity relations and fuzzy orderings, *Inform. Sci.* 3 (1971) 177–200.
- [27] L. A. Zadeh, A note on similarity-based definitions of possibility and probability, *Inform. Sci.* 267 (2014) 334–336.
- [28] W. R. Zhang, Bipolar fuzzy sets and relations: a computational framework for cognitive modeling and multiagent decision analysis, in: *Proceedings of IEEE Conference 1994* 305–309.
- [29] W. R. Zhang, Bipolar fuzzy sets, in: *Proceedings of FUZZ-IEEE 1998* 835–840.
- [30] W. R. Zhang and Yin Yang, *Bipolar relativity: A unifying theory of nature, agents and causality with applications in quantum computing, cognitive informatics and life sciences*, Information Sciences Reference, Hershey, 2011.
- [31] W. R. Zhang, L. Zhang and Yin Yang, bipolar logic and bipolar fuzzy logic, *Inform. Sci.* 165 (3–4) (2004) 265–287.

F. SALAMA (fatma.salama@science.tanta.edu.eg)

Mathematics Department, Faculty of Science, Tanta University, Tanta, Egypt