


On fuzzy strongly semiregular spaces

G. THANGARAJ, M. PONNUSAMY



Reprinted from the
Annals of Fuzzy Mathematics and Informatics
Vol. 26, No. 3, December 2023

On fuzzy strongly semiregular spaces

G. THANGARAJ, M. PONNUSAMY

Received 28 August 2023; Revised 10 September 2023; Accepted 10 November 2023

ABSTRACT. In this paper, the notion of fuzzy strongly semiregular space is introduced and studied. It is established that fuzzy strongly semiregular spaces are fuzzy quasi-regular spaces and fuzzy almost P -spaces. The conditions under which fuzzy strongly semiregular spaces become fuzzy Baire spaces, are also obtained. It is observed that fuzzy strongly semiregular and fuzzy hyperconnected spaces are fuzzy semi- P -spaces and fuzzy ∂ -spaces. It is found that fuzzy globally disconnected and fuzzy quasi-regular spaces are fuzzy strongly semiregular spaces. A condition under which fuzzy G_δ -sets become fuzzy Baire dense sets in fuzzy strongly semiregular spaces is also obtained.

2020 AMS Classification: 54A40, 03E72

Keywords: Fuzzy G_δ -set, Fuzzy F_σ -set, Fuzzy Baire dense set, Fuzzy residual set, Fuzzy σ -nowhere dense set, Fuzzy almost P -space, Fuzzy Baire space, Fuzzy hyperconnected space, Fuzzy globally disconnected space.

Corresponding Author: G. Thangaraj (g.thangaraj@rediffmail.com),

1. INTRODUCTION

The potential of fuzzy notion, introduced by Zadeh [1] in 1965 to describe vagueness mathematically in its very abstractness, was realized by many researchers. In 1968, Chang [2] introduced the notion of fuzzy topological spaces. Subsequently, many researchers have been worked in this area and other related areas which have applications in different fields based on this concept. In recent years, there has been a growing trend to introduce and study different kinds of fuzzy topological spaces.

In 1937, Stone [3] had introduced the idea of semiregular spaces in classical topology and obtained their characterizations. In 1969, Singal and Arya [4] introduced a new class of separation axiom (named almost-regular space) in topological spaces which is weaker than regularity but it is equivalent to semiregular spaces due to Stone. Nayar [5] has carried out a study on semi regular spaces in 2008. In 1981,

Azad [6] defined the notions of fuzzy semi-open set, fuzzy semi-closed set, fuzzy regular-open set, fuzzy regular-closed set, fuzzy regular spaces and fuzzy semi regular spaces. The notion of fuzzy semiregular spaces in the sense of Sostak was introduced and studied by Kim and Ko [7]. Fuzzy semi regularity in fuzzy topological spaces was also studied by Ajmal and Tyagi [8]. In 1998, Noiri and Jafari [9] introduced and studied the notion of strong semi-regularity in classical topology.

The aim of this paper is to introduce and study the notion of fuzzy strongly semi regular spaces. In Section 3, several characterizations of fuzzy strongly semi regular spaces are established. In Section 4, it is established that fuzzy strongly semiregular spaces are fuzzy quasi-regular spaces and fuzzy almost P-spaces. The conditions under which fuzzy strongly semiregular spaces become fuzzy Baire spaces, are also obtained. It is observed that fuzzy strongly semiregular and fuzzy hyperconnected spaces are fuzzy semi-P-spaces and fuzzy ∂ -spaces. It is found that fuzzy globally disconnected and fuzzy quasi-regular spaces are fuzzy strongly semiregular spaces and fuzzy G_δ -sets in fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semi regular spaces are fuzzy second category sets.

In recent years, the topological space theory has been embedding in the soft set theory to obtain some interesting applications [10, 11]. Many authors [12, 13] redefined the classical topological concepts via soft topological structure. Recently, Şenel et al. [14] applied the concept of octahedron sets to multi-criteria group decision making problems. Lee et al. [15] formed the concrete category of cubic relations and morphisms between them and studied some of its categorical structures in the sense of the topological universe. On these lines, there is a need and scope of investigation considering different types of fuzzy spaces such as fuzzy semi regular spaces and fuzzy strongly semi regular spaces for applying some fuzzy topological concepts to information science and decision-making problems. Throughout this paper, \mathbb{N} denotes the set of all positive integers.

2. PRELIMINARIES

Some basic notions and results used in the sequel, are given in order to make the exposition self-contained. In this work by (X, T) or simply by X , we will denote a fuzzy topological space due to Chang (1968). Let X be a non-empty set and I the unit interval $[0, 1]$. A fuzzy set λ in X is a mapping from X into I . The fuzzy set 0_X is defined as $0_X(x) = 0$ for all $x \in X$ and the fuzzy set 1_X is defined as $1_X(x) = 1$ for all $x \in X$. The *fuzzy interior*, the *fuzzy closure* and the *complement* of a fuzzy set λ in a fuzzy topological space (X, T) are defined respectively as follows:

- (i) $int(\lambda) = \bigvee \{ \mu : \mu \leq \lambda, \mu \in T \}$,
- (ii) $cl(\lambda) = \bigwedge \{ \mu : \lambda \leq \mu, 1 - \mu \in T \}$,
- (iii) $\lambda'(x) = 1 - \lambda(x)$ for all $x \in X$.

For a family $(\lambda_j)_{j \in J}$ of fuzzy sets in (X, T) , where J denotes an index set, the *union* $\psi = \bigvee_{j \in J} \lambda_j$ and the *intersection* $\delta = \bigwedge_{j \in J} \lambda_j$, are defined respectively as follows: for each $x \in X$,

- (iv) $\psi(x) = \sup_{j \in J} \{ \lambda_j(x) : x \in X \}$,
- (v) $\delta(x) = \inf_{j \in J} \{ \lambda_j(x) / x \in X \}$.

Lemma 2.1 (Lemma 3.2, [6]). *For a fuzzy set λ of a fuzzy topological space X ,*

- (1) $1 - \text{int}(\lambda) = \text{cl}(1 - \lambda)$,
- (2) $1 - \text{cl}(\lambda) = \text{int}(1 - \lambda)$.

Definition 2.2. A fuzzy set λ in a fuzzy topological space (X, T) is called a:

- (i) *fuzzy regular-open set* in X , if $\lambda = \text{intcl}(\lambda)$ and *fuzzy regular-closed set* in X , if $\lambda = \text{clint}(\lambda)$ [6],
- (ii) *fuzzy semi-open set* in X , if $\lambda \leq \text{clint}(\lambda)$ and *fuzzy semi-closed set* in (X, T) , if $\text{intcl}(\lambda) \leq \lambda$ [6],
- (iii) *fuzzy G_δ -set* in X , if $\lambda = \bigwedge_{j=1}^\infty \lambda_j$, where $\lambda_j \in T$ for $j \in J$ and *fuzzy F_σ -set* in X , if $\lambda = \bigvee_{j=1}^\infty \mu_j$, where $1 - \mu_j \in T$ for $j \in J$ [16],
- (iv) *fuzzy semi- G_δ -set* in X if $\lambda = \bigwedge_{j=1}^\infty \lambda'_j$, where each λ'_j is a fuzzy semi-open set in X and *fuzzy semi- F_σ -set* in X , if $\lambda = \bigvee_{j=1}^\infty \mu'_j$, where each μ'_j is a fuzzy semi-closed set in X [17],
- (v) *fuzzy dense set* in X , if there exists no fuzzy closed set μ in X such that $\lambda < \mu < 1$, i.e., $\text{cl}(\lambda) = 1$ [18],
- (vi) *fuzzy nowhere dense set* in X , if there exists no non-zero fuzzy open set μ in X such that $\mu < \text{cl}(\lambda)$, i.e., $\text{intcl}(\lambda) = 0$ [18],
- (vii) *fuzzy first category set* in X , if $\lambda = \bigvee_{j=1}^\infty \lambda_j$, where each λ_j is a fuzzy nowhere dense set in X and any other fuzzy set in X is said to be of *fuzzy second category* [18],
- (viii) *fuzzy residual set* in X , if $1 - \lambda$ is a fuzzy first category set in X [19],
- (ix) *fuzzy σ -nowhere dense set* in X , if λ is a fuzzy F_σ -set with $\text{int}(\lambda) = 0$ [20],
- (x) *fuzzy somewhere dense set* in X , if there exists a non-zero fuzzy open set μ in X such that $\mu < \text{cl}(\lambda)$, i.e., $\text{intcl}(\lambda) \neq 0$ [21] and $1 - \lambda$ is called a *fuzzy complement of fuzzy somewhere dense set* (briefly, fuzzy cs dense set) in X [22],
- (xi) *fuzzy δ -open set* in X , if $\lambda = \bigvee_{j \in J} \lambda_j$, where each λ_j is a fuzzy regular open set in X and *fuzzy δ -closed set* in X , if $1 - \lambda$ is a fuzzy δ -open set in X [23],
- (xii) *fuzzy locally closed set* (resp. *fuzzy A -set*) in X , if $\lambda = \mu \wedge \delta$, where μ is a fuzzy open set and δ is a fuzzy closed set (resp. fuzzy regular closed) in X [24],
- (xiii) *fuzzy Baire set* in X , if $\lambda = \mu \wedge \delta$, where μ is a non-zero fuzzy open set and δ is a fuzzy residual set in X [25],
- (xiv) *fuzzy pseudo-open set* in X , if $\lambda = \mu \vee \delta$, where μ is a non-zero fuzzy open set and δ is a fuzzy first category set in X [26],
- (xv) *fuzzy simply open set* in X , if $\text{Bd}(\lambda) = \text{cl}(\lambda) \wedge \text{cl}(1 - \lambda)$ is a fuzzy nowhere dense set in X [27],
- (xvi) *fuzzy Baire dense set* in X , if for a non-zero fuzzy open set μ in X , $\lambda \wedge \mu$ is a fuzzy second category set in X [28].

Definition 2.3. A fuzzy topological space (X, T) is called a:

- (i) *fuzzy regular space*, if for each $\lambda \in T$, there is $(\lambda_j)_{j \in J} \subset T$ such that $\text{cl}(\lambda_j) \leq \lambda$ for each $j \in J$ and $\lambda = \bigvee_{j \in J} \lambda_j$ [6],
- (ii) *fuzzy semi regular*, if the collection of all fuzzy regular open sets in X forms a base for T [6],
- (iii) *fuzzy Baire space*, if $\text{int}(\bigvee_{j=1}^\infty \lambda_j) = 0$, where each λ_j is a fuzzy nowhere dense set in X [19],

- (iv) *fuzzy quasi-regular space*, if for each $\lambda \in T$, there exists a fuzzy regular closed set μ in (X, T) such that $\mu \leq \lambda$ [29],
- (v) *fuzzy P-space*, if for each fuzzy G_δ -set in X , $G_\delta \in T$ [30],
- (vi) *fuzzy almost P-space*, if for each non-zero fuzzy G_δ -set λ in X , $\text{int}(\lambda) \neq 0$ [31],
- (vii) *fuzzy semi-P-space*, if each fuzzy G_δ -set in X is a fuzzy semi-open set in X [32],
- (viii) *fuzzy globally disconnected space*, if each fuzzy semi-open set in X is a fuzzy open set in X [33],
- (ix) *fuzzy hyperconnected space* if every non-null fuzzy open set in X is fuzzy dense in X [34],
- (x) *fuzzy ∂ -space*, if each fuzzy G_δ -set in X is a fuzzy simply open set in X [35],
- (xi) *fuzzy strongly regular space*, if for each fuzzy semi-open set λ in X , there exist $(\lambda_j)_{j \in J} \subset T$ such that $\lambda = \bigvee_{j \in J} \lambda_j$ and $\text{cl}(\lambda_j) \leq \lambda$ [36].
- (xii) *it fuzzy first category space*, if $1_X = \bigvee_{n=1}^{\infty} \lambda_n$, where $(\lambda_n)_{n \in \mathbb{N}}$ is a family of fuzzy nowhere dense set in X . A fuzzy topological space which is not of fuzzy first category is said to be of *fuzzy second category* [18].

Theorem 2.4 (Theorem 5.6, [6]). *In a fuzzy topological space,*

- (1) *the closure of a fuzzy open set is a fuzzy regular closed set,*
- (2) *the interior of a fuzzy closed set is a fuzzy regular open set.*

Theorem 2.5 (Proposition 3.5, [17]). *If λ is a fuzzy residual set in a fuzzy topological space (X, T) , then λ is a fuzzy semi- G_δ -set in (X, T) .*

Theorem 2.6 (Proposition 4.3, [19]). *Let (X, T) be a fuzzy topological space. Then the following are equivalent:*

- (1) *X is a fuzzy Baire space,*
- (2) *$\text{int}(\lambda) = 0$ for every fuzzy first category set λ in X ,*
- (3) *$\text{cl}(\mu) = 1$ for every fuzzy residual set μ in X .*

Theorem 2.7 (Proposition 3.1, [20]). *If λ is a fuzzy σ -nowhere dense set in a fuzzy topological space (X, T) , then $1 - \lambda$ is a fuzzy residual set in X .*

Theorem 2.8 (Proposition 3.5, [28]). *If λ is a fuzzy somewhere dense set in a fuzzy topological space (X, T) , then there exists a fuzzy regular closed set η in X such that $\eta \leq \text{cl}(\lambda)$.*

Theorem 2.9 (Proposition 3.12, [25]). *If λ is a fuzzy Baire set in a fuzzy topological space (X, T) , then $\lambda = \mu \wedge \delta$, where $\mu \in T$ and δ is a fuzzy semi- G_δ -set in X .*

Theorem 2.10 (Proposition 3.16, [27]). *If each fuzzy simply open set λ is a fuzzy open set in a fuzzy topological space (X, T) , then λ is a fuzzy semi-open set in X .*

Theorem 2.11 (Proposition 4.1, [31]). *If a fuzzy topological space (X, T) is a fuzzy almost P-space, then X is a fuzzy second category space.*

Theorem 2.12 (Proposition 3.6, [33]). *If λ is a fuzzy residual set in a fuzzy globally disconnected space (X, T) , then λ is a fuzzy G_δ -set in X .*

Theorem 2.13 (Proposition 3.5, [37]). *If λ is a fuzzy residual set in a fuzzy topological space (X, T) , then there exists a fuzzy G_δ -set μ in X such that $\mu \leq \lambda$.*

Theorem 2.14 (Proposition 3.1, [31]). *A fuzzy topological space (X, T) is a fuzzy almost P -space if and only if the only fuzzy F_σ -set λ in X such that $cl(\lambda) = 1$ is 1_X .*

Theorem 2.15 (Proposition 3.11, [38]). *If $clint(\mu) = 1$ for each fuzzy G_δ -set μ in a fuzzy topological space (X, T) , then X is a fuzzy ∂ -space.*

Theorem 2.16 (Proposition 5.1, [39]). *If a fuzzy topological space (X, T) is a fuzzy hyperconnected and fuzzy P -space, then X is a fuzzy Baire space.*

Theorem 2.17 (Proposition 5.6, [39]). *If λ is a fuzzy G_δ -set in a fuzzy Baire and fuzzy P -space (X, T) , then λ is a fuzzy second category set in X .*

Theorem 2.18 (Proposition 4.6, [40]). *If λ is a fuzzy Baire set in a fuzzy globally disconnected, fuzzy Baire and fuzzy P -space (X, T) , then λ is a fuzzy Baire dense set in X .*

3. FUZZY STRONGLY SEMIREGULAR SPACES

Motivated by the works of Jafari and Noiri [9] on strongly semiregular spaces in classical topology, the notion of fuzzy strongly semiregular space is introduced as follows.

Definition 3.1. A fuzzy topological space (X, T) is called a *fuzzy strongly semiregular space*, if for each fuzzy semi-open set λ in X , there exists a fuzzy regular closed set μ in X such that $\mu \leq \lambda$.

Example 3.2. Let $X = \{a, b, c\}$ and let $\alpha, \beta, \gamma, \delta$ and θ be the fuzzy sets in X defined as follows:

$$\alpha(a) = 0.5, \alpha(b) = 0.6, \alpha(c) = 0.4; \beta(a) = 0.6, \beta(b) = 0.4, \beta(c) = 0.5;$$

$$\gamma(a) = 0.5, \gamma(b) = 0.4, \gamma(c) = 0.6; \delta(a) = 0.7, \delta(b) = 0.6, \delta(c) = 0.8;$$

$$\theta(a) = 0.6, \theta(b) = 0.5, \theta(c) = 0.6.$$

Then $T = \{0, \alpha, \beta, \gamma, \alpha \vee \beta, \alpha \vee \gamma, \beta \vee \gamma, \alpha \wedge \beta, \beta \wedge \gamma, \alpha \vee (\beta \wedge \gamma), \alpha \vee \beta \vee \gamma, 1\}$ is a fuzzy topology on X . By computation, one can find that:

$$\begin{aligned} cl(\alpha) &= 1 - \gamma = \alpha, \quad cl(\beta) = 1, \quad cl(\gamma) = 1 - \alpha = \gamma, \quad cl(\alpha \vee \beta) = 1, \\ cl(\alpha \vee \gamma) &= 1 - (\alpha \wedge \beta) = \alpha \vee \gamma, \quad cl(\beta \vee \gamma) = 1, \quad cl(\alpha \wedge \beta) = 1 - (\alpha \vee \gamma) = \alpha \wedge \beta, \\ cl(\beta \wedge \gamma) &= 1 - [\alpha \vee (\beta \wedge \gamma)] = \beta \wedge \gamma, \quad cl(\alpha \vee [\beta \wedge \gamma]) = 1 - (\beta \wedge \gamma) = \alpha \vee (\beta \wedge \gamma), \\ cl(\alpha \vee \beta \vee \gamma) &= 1; \\ int(1 - \alpha) &= \gamma, \quad int(1 - \beta) = 0, \quad int(1 - \gamma) = \alpha; \quad int[1 - (\alpha \vee \beta)] = 0, \\ int(1 - [\alpha \vee \gamma]) &= \alpha \wedge \beta, \quad int(1 - [\beta \vee \gamma]) = 0, \quad int(1 - [\alpha \wedge \beta]) = \alpha \vee \gamma, \\ int(1 - [\beta \wedge \gamma]) &= \alpha \vee (\beta \wedge \gamma), \quad int[1 - (\alpha \vee [\beta \wedge \gamma])] = \beta \wedge \gamma, \quad int(1 - [\alpha \vee \beta \vee \gamma]) = 0. \end{aligned}$$

Also by computation, one can find that $clint(\theta) = 1$ and $clint(\delta) = 1$ and thus δ and θ are the fuzzy semi-open sets in X . By computation, one can find that the fuzzy regular closed sets in X are $1 - \alpha, 1 - \gamma, 1 - (\alpha \vee \gamma), 1 - (\alpha \wedge \beta), 1 - (\beta \wedge \gamma)$ and $1 - [\alpha \vee \beta \vee \gamma]$ and the fuzzy semi-open sets are all the fuzzy open sets and δ and θ and for each fuzzy semi-open set λ in X , there exists a fuzzy regular closed set $\eta [= 1 - \alpha, 1 - \gamma, 1 - (\alpha \vee \gamma), 1 - (\alpha \wedge \beta), 1 - (\beta \wedge \gamma)$ and $1 - (\alpha \vee \beta \vee \gamma)]$ in X such that $\eta \leq \lambda$. So X is a fuzzy strongly semiregular space.

Proposition 3.3. *If a fuzzy topological space (X, T) is a fuzzy strongly semi-regular space, then for each fuzzy semi-closed set δ in X , there exists a fuzzy regular open set η in X such that $\delta \leq \eta$.*

Proof. Suppose δ is a fuzzy semi-closed set in X . Then $1 - \delta$ is a fuzzy semi-open set in X . As X is a fuzzy strongly semiregular space, there exists a fuzzy regular closed set μ in X such that $\mu \leq 1 - \delta$. Thus $\delta \leq 1 - \mu$. Let $\eta = 1 - \mu$. Then η is a fuzzy regular open set in X and $\delta \leq \eta$. \square

Proposition 3.4. *If δ is a fuzzy closed set in a fuzzy strongly semiregular space (X, T) , then there exists a fuzzy regular open set η in X such that $\delta \leq \eta$.*

Proof. Suppose δ is a fuzzy closed set in X . Then clearly, δ is a fuzzy semi closed set in X . Since X is a fuzzy strongly semi regular space, by Proposition 3.3, there exists a fuzzy regular open set η in X such that $\delta \leq \eta$. \square

Proposition 3.5. *If λ is a fuzzy open set in a fuzzy strongly semiregular space (X, T) , then there exists a fuzzy regular closed set μ in X such that $\mu \leq \lambda$.*

Proof. Suppose λ is a fuzzy open set in X . Then clearly, λ is a fuzzy semi-open set in X . Since X is a fuzzy strongly semiregular space, there exists a fuzzy regular closed set μ in X such that $\mu \leq \lambda$. \square

Remark 3.6. In view of Propositions 3.4 and 3.5, one will have the following result.

If λ is a fuzzy open set in a fuzzy strongly semiregular space (X, T) , then there exists a fuzzy regular closed set μ and a fuzzy regular open set η in X such that $\mu \leq \lambda \leq \eta$ [for, $\mu \leq \lambda \leq cl(\lambda) \leq \eta$].

Proposition 3.7. *If λ is a fuzzy residual set in a fuzzy strongly semiregular space (X, T) , there exist a family $(\delta_n)_{n \in \mathbb{N}}$ of fuzzy regular closed sets in X such that $\bigwedge_{n=1}^{\infty} \delta_n \leq \lambda$.*

Proof. Suppose λ is a fuzzy residual set in X . Then, by Theorem 2.5, λ is a fuzzy semi- G_δ - set in X and thus $\lambda = \bigwedge_{n=1}^{\infty} \mu_n$, where $(\mu_n)_{n \in \mathbb{N}}$ is a family of fuzzy semi-open sets in X . Since X is a fuzzy strongly semiregular space, by Proposition 3.5, there exists a fuzzy regular closed set δ_n in X such that $\delta_n \leq \mu_n$. So $\bigwedge_{n=1}^{\infty} \delta_n \leq \bigwedge_{n=1}^{\infty} \mu_n$ and $\bigwedge_{n=1}^{\infty} \delta_n \leq \lambda$. \square

Corollary 3.8. *If λ is a fuzzy residual set in a fuzzy strongly semiregular space (X, T) , then there exist a fuzzy closed set μ in X such that $\mu \leq \lambda$.*

Proof. Suppose λ is a fuzzy residual set in X . Then by Proposition 3.7, there exist a family $(\delta_n)_{n \in \mathbb{N}}$ of fuzzy regular closed sets in X such that $\bigwedge_{n=1}^{\infty} \delta_n \leq \lambda$. Since fuzzy regular closed sets are fuzzy closed sets in a fuzzy topological space, $(\delta_n)_{n \in \mathbb{N}}$ is a family of fuzzy closed sets in X . Let $\mu = \bigwedge_{n=1}^{\infty} \delta_n$. Then μ is a fuzzy closed set in X and $\mu \leq \lambda$. \square

Proposition 3.9. *If λ is a fuzzy residual set in a fuzzy strongly semiregular space (X, T) , then $int(\lambda) \neq 0$.*

Proof. Suppose λ is a fuzzy residual set in X . Then by Corollary 3.8, there exist a fuzzy closed set μ in X such that $\mu \leq \lambda$. This implies that $\text{int}(\mu) \leq \text{int}(\lambda)$. Thus by Theorem 2.4, $\text{int}(\mu)$ is a fuzzy regular open set in X . So $\text{int}(\mu)$ is a fuzzy open set in X . Hence $\text{int}(\mu) \leq \text{int}(\lambda)$ implies that $\text{int}(\lambda) \neq 0$. \square

Corollary 3.10. *If λ is a fuzzy residual set in a fuzzy strongly semiregular space (X, T) , then λ is a fuzzy somewhere dense set in X .*

Proof. Suppose λ is a fuzzy residual set in a fuzzy strongly semiregular space X . Then by Proposition 3.9, $\text{int}(\lambda) \neq 0$. Since $\text{int}(\lambda) \leq \text{intcl}(\lambda)$, $\text{int}(\lambda) \neq 0$. Thus $\text{intcl}(\lambda) \neq 0$. So λ is a fuzzy somewhere dense set in X . \square

Proposition 3.11. *If μ is a fuzzy first category set in a fuzzy strongly semiregular space (X, T) , there exist a family $(\gamma_n)_{n \in \mathbb{N}}$ of fuzzy regular open sets in X such that $\mu \leq \bigvee_{n=1}^{\infty} \gamma_n$.*

Proof. Suppose μ is a fuzzy first category set in X . Then $1 - \mu$ is a fuzzy residual set in X . Since X is a fuzzy strongly semiregular space, by Proposition 3.7, there exist a family $(\delta_n)_{n \in \mathbb{N}}$ of fuzzy regular closed sets in X such that $\bigwedge_{n=1}^{\infty} \delta_n \leq 1 - \mu$. Thus $\mu \leq 1 - \bigwedge_{n=1}^{\infty} \delta_n = \bigvee_{n=1}^{\infty} (1 - \delta_n)$. Let $\gamma_n = 1 - \delta_n$. Then $(\gamma_n)_{n \in \mathbb{N}}$ is a family of fuzzy regular open sets in X and $\mu \leq \bigvee_{n=1}^{\infty} \gamma_n$. \square

Corollary 3.12. *If μ is a fuzzy first category set in a fuzzy strongly semiregular space (X, T) , then there exist a fuzzy open set γ in X such that $\mu \leq \gamma$.*

Proof. Suppose μ is a fuzzy first category set in X . Then by Proposition 3.11, there exist a family $(\gamma_n)_{n \in \mathbb{N}}$ of fuzzy regular open sets in X such that $\mu \leq \bigvee_{n=1}^{\infty} \gamma_n$. Since fuzzy regular open sets are fuzzy open sets, $(\gamma_n)_{n \in \mathbb{N}}$ is a family of fuzzy open sets in X and $\mu \leq \gamma$. \square

The following propositions show that fuzzy first category sets, fuzzy σ -nowhere dense sets are not fuzzy dense sets in fuzzy strongly semiregular spaces.

Proposition 3.13. *If μ is a fuzzy first category set in a fuzzy strongly semiregular space (X, T) , then μ is not a fuzzy dense set in X .*

Proof. Suppose μ is a fuzzy first category set in X . Then by Corollary 3.12, there exist a fuzzy open set γ in X such that $\mu \leq \gamma$. Thus $\text{cl}(\mu) \leq \text{cl}(\gamma)$. By Theorem 2.4, $\text{cl}(\gamma)$ is a fuzzy regular closed set in X . So $\text{cl}(\gamma)$ is a fuzzy closed set in X . Let $\eta = \text{cl}(\gamma)$. Then there exist a fuzzy closed set η in X such that $\mu \leq \text{cl}(\mu) \leq \eta$. Thus μ is not a fuzzy dense set in X . \square

Proposition 3.14. *If λ is a fuzzy σ -nowhere dense set in a fuzzy strongly semiregular space (X, T) , then λ is not a fuzzy dense set in X .*

Proof. Suppose λ is a fuzzy σ -nowhere dense set in X . Then by Theorem 2.7, $1 - \lambda$ is a fuzzy residual set in X . Since X is a fuzzy strongly semiregular space, by Proposition 3.9, $\text{int}(1 - \lambda) \neq 0$. Thus by Lemma 2.1, $1 - \text{cl}(\lambda) \neq 0$. So $\text{cl}(\lambda) \neq 1$. Hence it follows that the fuzzy σ -nowhere dense set λ is not a fuzzy dense set in X . \square

Remark 3.15. In view of the Proposition 3.14, one will have the following result.

Fuzzy F_σ -sets with zero interior in fuzzy strongly semiregular spaces are not fuzzy dense sets.

Proposition 3.16. *If λ is a fuzzy residual set in a fuzzy strongly semiregular space (X, T) , then there exists a fuzzy regular closed set η in X such that $\eta \leq cl(\lambda)$.*

Proof. Suppose λ is a fuzzy residual set in X . Since X is a fuzzy strongly semiregular space, by Corollary 3.10, λ is a fuzzy somewhere dense set in X . Then by Theorem 2.8, there exists a fuzzy regular closed set η in X such that $\eta \leq cl(\lambda)$. \square

In classical topology, each semi-open set is the union of regular closed sets in a strongly semi-regular space whereas each fuzzy semi-open set lies between fuzzy regular closed sets in fuzzy strongly semiregular spaces. For, consider the following proposition.

Proposition 3.17. *If λ is a fuzzy semi-open set in a fuzzy strongly semiregular space (X, T) , then there exist fuzzy regular closed sets η_1 and η_2 in X such that $\eta_1 \leq \lambda \leq \eta_2$.*

Proof. Suppose λ is a fuzzy semi-open set in X . Since X is a fuzzy strongly semiregular space, there exists a fuzzy regular closed set η_1 in X such that $\eta_1 \leq \lambda$. Then, $\eta_1 \leq \lambda \leq cl(\lambda)$. Since the closure of a fuzzy semi-open set is a fuzzy regular closed set, $cl(\lambda)$ is a fuzzy regular closed set in X . Let $\eta_2 = cl(\lambda)$. Then $\eta_1 \leq \lambda \leq \eta_2$. Thus fuzzy semi-open sets lies between fuzzy regular closed sets in fuzzy strongly semiregular spaces. \square

Proposition 3.18. *If η is a fuzzy regular closed set in a fuzzy strongly semiregular space (X, T) , then there exists a fuzzy regular closed set γ in X such that $\gamma \leq \eta$.*

Proof. Suppose η is a fuzzy regular closed set in X . Then $clint(\eta) = \eta$. Since $int[int(\eta)] = int(\eta)$, $int(\eta) \leq \eta = clint(\eta) = cl(int[int(\eta)])$. Thus $int(\eta) \leq clint[int(\eta)]$. So $int(\eta)$ is a fuzzy semi-open set in X . Since X is a fuzzy strongly semiregular space, there exists a fuzzy regular closed set γ in X such that $\gamma \leq int(\eta)$. Hence $\gamma \leq int(\eta) \leq \eta$. \square

Corollary 3.19. *If η is a fuzzy regular closed set in a fuzzy strongly semiregular space (X, T) , then $\eta \geq \bigvee_{n=1}^{\infty} \gamma_n$, where $(\gamma_n)_{n \in \mathbb{N}}$ is a decreasing sequence of fuzzy regular closed sets in X .*

Proof. Suppose η is a fuzzy regular closed set in X . Then by Proposition 3.18, there exists a family $(\gamma_n)_{n \in \mathbb{N}}$ of fuzzy regular closed sets in X such that $\dots \leq \gamma_3 \leq \gamma_2 \leq \gamma_1 \leq \eta$. Thus $\eta \geq \bigvee_{n=1}^{\infty} \gamma_n$, where $(\gamma_n)_{n \in \mathbb{N}}$ is a decreasing sequence of fuzzy regular closed sets in X . \square

Proposition 3.20. *If θ is a fuzzy regular open set in a fuzzy strongly semiregular space (X, T) , then there exists a fuzzy regular open set ρ in X such that $\theta \leq \rho$.*

Proof. Suppose θ is a fuzzy regular open set in X . Then $1 - \theta$ is a fuzzy regular closed set in X . Since X is a fuzzy strongly semiregular space, by Proposition 3.18, there exists a fuzzy regular closed set γ in X such that $\gamma \leq 1 - \theta$. Thus $\theta \leq 1 - \gamma$. Let $\rho = 1 - \gamma$. Then ρ is a fuzzy regular open set in X and $\theta \leq \rho$. \square

Corollary 3.21. *If θ is a fuzzy regular open set in a fuzzy strongly semiregular space (X, T) , then $\theta \leq \bigwedge_{n=1}^{\infty} \delta_n$, where $(\delta_n)_{n \in \mathbb{N}}$ is an increasing sequence of fuzzy regular open sets in X .*

Proof. Suppose θ is a fuzzy regular open set in X . Then by Proposition 3.20, there exist a family $(\delta_n)_{n \in \mathbb{N}}$ of fuzzy regular open sets in X such that $\theta \leq \delta_1 \leq \delta_2 \leq \delta_3 \cdots$. Thus $\theta \leq \bigwedge_{n=1}^{\infty} \delta_n$, where $(\delta_n)_{n \in \mathbb{N}}$ is an increasing sequence of fuzzy regular open sets in X . \square

Proposition 3.22. *If λ is a fuzzy G_δ -set in a fuzzy strongly semiregular space (X, T) , then there exists a fuzzy closed set μ in X such that $\mu \leq \lambda$.*

Proof. Suppose λ is a fuzzy G_δ -set in X . Then $\lambda = \bigwedge_{n=1}^{\infty} \lambda_n$, where $(\lambda_n)_{n \in \mathbb{N}} \subset T$. Since X is a fuzzy strongly semiregular space, by Proposition 3.5, for each $n \in \mathbb{N}$, there exist fuzzy regular closed sets μ_n in X such that $\mu_n \leq \lambda_n$. Thus $\bigwedge_{n=1}^{\infty} \mu_n \leq \bigwedge_{n=1}^{\infty} \lambda_n$. Since each fuzzy regular closed set is a fuzzy closed set, $(\mu_n)_{n \in \mathbb{N}}$ is a family of fuzzy closed sets in X . So $\bigwedge_{n=1}^{\infty} \mu_n$ is a fuzzy closed set in X . Let $\mu = \bigwedge_{n=1}^{\infty} \mu_n$. Then for the fuzzy G_δ -set λ , there exists a fuzzy closed set μ in X such that $\mu \leq \lambda$. \square

Proposition 3.23. *If δ is a fuzzy F_σ -set in a fuzzy strongly semiregular space (X, T) , then there exists a fuzzy open set γ in X such that $\delta \leq \gamma$.*

Proof. Suppose δ is a fuzzy F_σ -set in X . Then $1 - \delta$ is a fuzzy G_δ -set in X . Since X is a fuzzy strongly semiregular space, by Proposition 3.22, there exists a fuzzy closed set μ in X such that $\mu \leq 1 - \delta$. Thus $\delta \leq 1 - \mu$. Let $\gamma \leq 1 - \mu$. Then clearly, $\gamma \in T$. Thus there exists a fuzzy open set γ in X such that $\delta \leq \gamma$. \square

Proposition 3.24. *If a fuzzy simply open set λ is a fuzzy open set in a fuzzy strongly semiregular space (X, T) , then there exists a fuzzy regular closed set μ in X such that $\mu \leq \lambda$.*

Proof. Suppose λ is a fuzzy simply open set in X . Then by the hypothesis, $\lambda \in T$. Thus by Theorem 2.10, λ is a fuzzy semi-open set in X . Since X is a fuzzy strongly semiregular space, there exists a fuzzy regular closed set μ in X such that $\mu \leq \lambda$. \square

Proposition 3.25. *If λ is a fuzzy Baire set in a fuzzy strongly semiregular space (X, T) , then there exist a fuzzy open set μ and a fuzzy closed set γ in X such that $\mu \wedge \gamma \leq \lambda$.*

Proof. Suppose λ is a fuzzy Baire set in X . Then by Theorem 2.9, $\lambda = \mu \wedge \delta$, where $\mu \in T$ and δ is a fuzzy semi- G_δ -set in X . Since δ is a fuzzy semi- G_δ -set, $\delta = \bigwedge_{n=1}^{\infty} \delta_n$, where $(\delta_i)_{i \in \mathbb{N}}$ is a family of fuzzy semi-open sets in X . Thus $\lambda = \mu \wedge (\bigwedge_{n=1}^{\infty} \delta_n)$. Since X is a fuzzy strongly semiregular space, for each $n \in \mathbb{N}$, there exist a fuzzy regular closed set γ_n in X such that $\gamma_n \leq \delta_n$. So $\bigwedge_{n=1}^{\infty} \gamma_n \leq \bigwedge_{n=1}^{\infty} \delta_n$. Hence $\bigwedge_{n=1}^{\infty} \gamma_n \leq \delta$. Let $\gamma = \bigwedge_{n=1}^{\infty} \gamma_n$. Then $\gamma \leq \delta$ and $\mu \wedge \gamma \leq \mu \wedge \delta = \lambda$. Since each fuzzy regular closed set is a fuzzy closed set, $(\gamma_n)_{n \in \mathbb{N}}$ is a family of fuzzy closed sets in X . Thus $\bigwedge_{n=1}^{\infty} \gamma_n$ is a fuzzy closed set in X . So γ is a fuzzy closed set in X . Hence by the hypothesis, there exist a fuzzy open set μ and a fuzzy closed set γ in X such that $\mu \wedge \gamma \leq \lambda$. \square

Remark 3.26. It should be noted that if λ is a fuzzy Baire set in a fuzzy strongly semiregular space (X, T) , then λ need not be a fuzzy semi-open set in X . For, in the above proof, $\gamma = \bigwedge_{n=1}^{\infty} \delta_n$, where $(\delta_n)_{n \in \mathbb{N}} \subset T$ and then $\lambda = \mu \wedge \delta$, where $\mu \in T$, need not be a fuzzy semi-open set in fuzzy topological spaces, whereas in ordinary topological setting the intersection of a semi-open set with an open set is a semi-open set (See Lemma 1, [41]).

Corollary 3.27. *If λ is a fuzzy Baire set in a fuzzy strongly semiregular space (X, T) , then there exist a fuzzy locally closed set η in X such that $\eta \leq \lambda$.*

Proof. The proof follows from the definition of fuzzy locally closed set and Proposition 3.25. □

Proposition 3.28. *If λ is a fuzzy residual set in a fuzzy strongly semiregular space (X, T) , then there exist a fuzzy G_δ -set μ and a fuzzy closed set δ in X such that $\delta \leq \mu \leq \lambda$.*

Proof. Suppose λ is a fuzzy residual set in X . Then by Theorem 2.13, there exists a fuzzy G_δ -set μ in X such that $\mu \leq \lambda$. Thus by Proposition 3.22, there exists a fuzzy closed set δ in X such that $\delta \leq \mu$. So $\delta \leq \mu \leq \lambda$. □

Proposition 3.29. *If γ is a fuzzy first category set in a fuzzy strongly semiregular space (X, T) , then there exist a fuzzy F_σ -set η and a fuzzy open set λ in X such that $\gamma \leq \eta \leq \lambda$.*

Proof. Suppose γ is a fuzzy first category set in a fuzzy strongly semiregular space X . Then clearly, $1 - \gamma$ is a fuzzy residual set in X . Thus by Proposition 3.28, there exist a fuzzy G_δ -set μ and a fuzzy closed set δ in X such that $\delta \leq \mu \leq 1 - \gamma$. So $1 - \delta \geq 1 - \mu \geq \gamma$. Let $\eta = 1 - \mu$ and $\lambda = 1 - \delta$. Then η is a fuzzy F_σ -set and λ is a fuzzy open set in X . Thus by the hypothesis, there exist a fuzzy F_σ -set η and a fuzzy open set λ in X such that $\gamma \leq \eta \leq \lambda$. □

Proposition 3.30. *If μ is a fuzzy first category set in a fuzzy strongly semiregular space (X, T) , then there exists a fuzzy δ -open γ in X such that $\mu \leq \gamma$.*

Proof. Suppose μ is a fuzzy first category set in X . Since (X, T) is a fuzzy strongly semiregular space, by Proposition 3.11, there exist a family $(\gamma_n)_{n \in \mathbb{N}}$ of fuzzy regular open sets in X such that $\mu \leq \bigvee_{n=1}^{\infty} \gamma_n$. Let $\gamma = \bigvee_{n=1}^{\infty} \gamma_n$. Then γ is a fuzzy δ -open in X . Thus $\mu \leq \gamma$. □

Proposition 3.31. *If λ is a fuzzy residual set in a fuzzy strongly semiregular space (X, T) , then there exist a fuzzy δ -closed set η in X such that $\eta \leq \lambda$.*

Proof. Suppose λ is a fuzzy residual set in X . Then $1 - \lambda$ is a fuzzy first category set in X . Since X is a fuzzy strongly semiregular space, by Proposition 3.30, there exists a fuzzy δ -open γ in X such that $1 - \lambda \leq \gamma$. Thus $1 - \gamma \leq \lambda$. Let $\eta = 1 - \gamma$. Then η is a fuzzy δ -closed set in X . Thus $\eta \leq \lambda$. □

Remark 3.32. It is to be mentioned that any fuzzy regular open set is fuzzy δ -open and any fuzzy δ -open set is a fuzzy open set in a fuzzy topological space. Furthermore, if a fuzzy set λ is fuzzy semi-open in a fuzzy topological space X , then $cl(\lambda) = cl\delta(\lambda)$ (See [38]).

Proposition 3.33. *If a fuzzy topological space (X, T) is a fuzzy strongly semi-regular space, then for each fuzzy semi-closed set δ in X , there exists a fuzzy δ -open set η in X such that $\delta \leq \eta$.*

Proof. Let δ be a fuzzy semi-closed set in X . Since X is a fuzzy strongly semiregular space, by Proposition 3.3, there exists a fuzzy regular open set η in X such that $\delta \leq \eta$. Since each fuzzy regular open set is a fuzzy δ -open set, η is a fuzzy δ -open in X and $\delta \leq \eta$. \square

Proposition 3.34. *If α is a fuzzy pseudo open-set in a fuzzy strongly semiregular space (X, T) , then there exist a fuzzy open set β in X such that $\alpha \leq \beta$.*

Proof. Suppose α is a fuzzy pseudo open-set in X . Then $\alpha = \mu \vee \gamma$, where μ is a non-zero fuzzy open set and γ is a fuzzy first category set in X . Since X is a fuzzy strongly semiregular space, by Proposition 3.29, there exist a fuzzy F_σ -set η and a fuzzy open set λ in X such that $\gamma \leq \eta \leq \lambda$. Thus $\mu \vee \gamma \leq \mu \vee \eta \leq \mu \vee \lambda$ and $\alpha \leq \mu \vee \lambda$. Let $\beta = \mu \vee \lambda$. Then clearly, $\alpha \leq \beta$. Furthermore, $\beta \in T$. \square

Proposition 3.35. *If λ is a fuzzy semi-open set in a fuzzy strongly semi-regular space (X, T) , then there exists a fuzzy regular open set η in X such that $\eta \leq \lambda$.*

Proof. Suppose λ is a fuzzy semi-open set in X . Since X is a fuzzy strongly semiregular space, there exists a fuzzy regular closed set μ in X such that $\mu \leq \lambda$. Then $\text{int}(\mu) \leq \mu \leq \lambda$. Since μ is a fuzzy regular closed set in X , $\mu = \text{clint}(\mu)$. On the other hand, $\text{intcl}(\text{int}(\mu)) = \text{int}(\text{clint}(\mu)) = \text{int}(\mu)$. Thus $\text{int}(\mu)$ is a fuzzy regular open set in X . Let $\eta = \text{int}(\mu)$. Then clearly, $\mu \leq \lambda$. Moreover, η is a fuzzy regular open set in X . \square

Corollary 3.36. *If λ is a fuzzy open set in a fuzzy strongly semiregular space (X, T) , then there exists a fuzzy regular open set η in X such that $\eta \leq \lambda$.*

Proof. Since each fuzzy open set is a fuzzy semi-open set in a fuzzy topological space, the proof follows from Proposition 3.35. \square

Proposition 3.37. *If λ is a fuzzy set in a fuzzy strongly semiregular space (X, T) , then there exist fuzzy regular open sets η_1 and η_2 in X such that $\eta_1 \leq \lambda \leq \eta_2$.*

Proof. Suppose λ is a fuzzy set in a fuzzy strongly semiregular space (X, T) . Then clearly, $\text{int}(\lambda) \in T$. Thus by Corollary 3.36, there exists a fuzzy regular open set η_1 in X such that $\eta_1 \leq \text{int}(\lambda)$. Since $\text{cl}(\lambda)$ is a fuzzy closed set in X , by Proposition 3.4, there exists a fuzzy regular open set η_2 in X such that $\text{cl}(\lambda) \leq \eta_2$. Thus $\eta_1 \leq \text{int}(\lambda) \leq \lambda \leq \text{cl}(\lambda) \leq \eta_2$. So $\eta_1 \leq \lambda \leq \eta_2$. \square

Proposition 3.38. *If λ is a fuzzy σ -nowhere dense set in a fuzzy strongly semiregular space (X, T) , then there exists a fuzzy open γ in X such that $\lambda \leq \gamma$.*

Proof. Suppose λ is a fuzzy σ -nowhere dense set in X . Then by Theorem 2.7, $1-\lambda$ is a fuzzy residual set in X . Since X is a fuzzy strongly semiregular space, by corollary 3.8, there exist a fuzzy closed set μ in X such that $\mu \leq 1-\lambda$. Thus $\lambda \leq 1-\mu$. Let $\gamma = 1-\mu$. Then clearly, $\lambda \leq \gamma$ and $\sigma \in T$. \square

4. FUZZY STRONGLY SEMIREGULAR SPACES AND OTHER FUZZY TOPOLOGICAL SPACES

The inter-relations between fuzzy strongly regular spaces, fuzzy strongly semiregular spaces, fuzzy quasi-regular spaces and fuzzy almost P-spaces are established in this section.

The conditions under which fuzzy Baire sets in fuzzy strongly semi regular spaces become fuzzy Baire dense sets are obtained.

Proposition 4.1. *If a fuzzy topological space (X, T) is a fuzzy strongly semiregular space, then (X, T) is a fuzzy quasi-regular space.*

Proof. Let $\lambda \in T$. Since each fuzzy open set is a fuzzy semi-open set, λ is a fuzzy semi-open set in X . Since X is a fuzzy strongly semiregular space, there exists a fuzzy regular closed set μ in X such that $\mu \leq \lambda$. Thus X is a fuzzy quasi-regular space. \square

Proposition 4.2. *If a fuzzy topological space (X, T) is a fuzzy globally disconnected and fuzzy quasi regular space, then X is a fuzzy strongly semi regular space.*

Proof. Suppose X is a fuzzy globally disconnected and fuzzy quasi regular space, and let λ be a fuzzy semi-open set in X . Since X is a fuzzy globally disconnected space, $\lambda \in T$. Since X is a fuzzy quasi regular space, there exists a fuzzy regular closed set μ in X such that $\mu \leq \lambda$. Then X is a fuzzy strongly semiregular space. \square

Proposition 4.3. *If a fuzzy first category space (X, T) is a fuzzy strongly semiregular space, then $\bigvee_{n=1}^{\infty} \eta_n = 1$, where $(\eta_n)_{n \in \mathbb{N}}$ is a family of fuzzy regular open sets in X .*

Proof. Suppose a fuzzy first category space X is a fuzzy strongly semiregular space. It is clear that 1 is a fuzzy first category set in X . Then $\bigvee_{n=1}^{\infty} (\lambda_n) = 1$, where $(\lambda_n)_{n \in \mathbb{N}}$ is a family of fuzzy nowhere dense sets in X . Thus $\text{intcl}(\lambda_n) = 0$, i.e., $\text{intcl}(\lambda_n) \leq \lambda_n$ for each $n \in \mathbb{N}$. So $(\lambda_n)_{n \in \mathbb{N}}$ is a family of fuzzy semi-closed sets in X . By the hypothesis and Proposition 3.3, for each $n \in \mathbb{N}$, there exist fuzzy regular open set η_n in X such that $\lambda_n \leq \eta_n$. This implies that $\bigvee_{n=1}^{\infty} \lambda_n \leq \bigvee_{n=1}^{\infty} \eta_n$ and $1 \leq \bigvee_{n=1}^{\infty} \eta_n$. Hence $\bigvee_{n=1}^{\infty} \eta_n = 1$, where $(\eta_n)_{n \in \mathbb{N}}$ is a family of fuzzy regular open sets in X . \square

The following proposition gives a condition for fuzzy strongly semiregular spaces to become fuzzy Baire spaces by means of fuzzy G_δ -sets.

Proposition 4.4. *If each fuzzy G_δ -set is a fuzzy dense set in a fuzzy strongly semiregular space (X, T) , then X is a fuzzy Baire space.*

Proof. Let λ be a fuzzy residual set in X . Since X is a fuzzy strongly semiregular space, by Proposition 3.28, there exist a fuzzy G_δ -set μ and a fuzzy closed set δ in X such that $\delta \leq \mu \leq \lambda$. Then $\text{cl}(\delta) \leq \text{cl}(\mu) \leq \text{cl}(\lambda)$. By the hypothesis, μ is a fuzzy dense set in X and thus $\text{cl}(\mu) = 1$. So $1 \leq \text{cl}(\lambda)$, i.e., $\text{cl}(\lambda) = 1$. Hence by Theorem 2.6, X is a fuzzy Baire space. \square

The following proposition gives a condition for fuzzy strongly semiregular spaces to become fuzzy Baire spaces by means of fuzzy F_σ -sets.

Proposition 4.5. *If $\text{int}(\mu) = 0$ for each fuzzy F_σ -set μ in a fuzzy strongly semiregular space (X, T) , then X is a fuzzy Baire space.*

Proof. Let γ be a fuzzy first category set in X . Since (X, T) is a fuzzy strongly semiregular space, by Proposition 3.29, there exist a fuzzy F_σ -set η and a fuzzy open set λ in X such that $\gamma \leq \eta \leq \lambda$. Then $\text{int}(\gamma) \leq \text{int}(\eta) \leq \text{int}(\lambda) = \lambda$. Thus By the hypothesis, $\text{int}(\eta) = 0$. So $\text{int}(\gamma) = 0$. Hence by Theorem 2.6, (X, T) is a fuzzy Baire space. \square

Proposition 4.6. *If (X, T) is a fuzzy strongly semiregular space, then X is a fuzzy almost P -space*

Proof. Let λ be a fuzzy G_δ -set in X . Since X is a fuzzy strongly semiregular space, by Proposition 3.22, there exists a fuzzy closed set μ in X such that $\mu \leq \lambda$. Then $\text{int}(\mu) \leq \text{int}(\lambda)$. By Lemma 2.1, $\text{int}(\mu)$ is a fuzzy regular open set and thus a fuzzy open set in X . So $\text{int}(\lambda) \neq 0$. Hence X is a fuzzy almost P -space. \square

Proposition 4.7. *If (X, T) is a fuzzy strongly semiregular space, then the only fuzzy F_σ -set λ such that $\text{cl}(\lambda) = 1$ is 1_X .*

Proof. Suppose X is a fuzzy strongly semiregular space. Then by Proposition 4.6, X is a fuzzy almost P -space. Thus by Theorem 2.14, the only fuzzy F_σ -set λ such that $\text{cl}(\lambda) = 1$ is 1_X . \square

Proposition 4.8. *If λ is a fuzzy G_δ -set in a fuzzy strongly semiregular and fuzzy hyperconnected space (X, T) , then λ is a fuzzy semi-open set in X .*

Proof. Suppose λ is a fuzzy G_δ -set in X . Since X is a fuzzy strongly semiregular space, by Proposition 4.6, X is a fuzzy almost P -space. Then $\text{int}(\lambda) \neq 0$. Since X is a fuzzy hyperconnected space and $\text{int}(\lambda) \in T$, $\text{int}(\lambda)$ is a fuzzy dense set in X . Thus $\text{clint}(\lambda) = 1$. So $\lambda \leq \text{clint}(\lambda)$. Hence λ is a fuzzy semi-open set in X . \square

Proposition 4.9. *If a fuzzy topological space (X, T) is a fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semi regular space, then X is a fuzzy P -space.*

Proof. Let λ be a fuzzy G_δ -set in X . Since X is a fuzzy strongly semiregular and fuzzy hyperconnected space, by Proposition 4.8, λ is a fuzzy semi-open set in X . Since X is a fuzzy globally disconnected space, $\lambda \in T$. Then X is a fuzzy P -space. \square

Proposition 4.10. *If a fuzzy topological space (X, T) is a fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semi regular space, then X is a fuzzy Baire space.*

Proof. The proof follows from Proposition 4.8 and Theorem 2.16. \square

Proposition 4.11. *If a fuzzy topological space (X, T) is a fuzzy hyperconnected fuzzy strongly semi regular and fuzzy P -space, then X is a fuzzy Baire space.*

Proof. Let λ be a fuzzy G_δ -set in X . Since X is a fuzzy P -space, $\lambda \in T$. Since X is a fuzzy hyperconnected space, λ is a fuzzy dense set in X . Then λ is a fuzzy dense set in X . Thus, by Proposition 4.4, X is a fuzzy Baire space. \square

The following propositions show that fuzzy strongly semiregular and fuzzy hyperconnected spaces are fuzzy semi- P -spaces and fuzzy ∂ -spaces.

Proposition 4.12. *If a fuzzy topological space (X, T) is a fuzzy strongly semiregular and fuzzy hyperconnected space, then X is a fuzzy semi-P-space .*

Proof. Let λ be a fuzzy G_δ -set in X . Since X is a fuzzy strongly semiregular and fuzzy hyperconnected space, by Proposition 4.8, λ is a fuzzy semi-open set in X . Then (X, T) is a fuzzy semi-P-space. \square

Proposition 4.13. *If a fuzzy topological space (X, T) is a fuzzy strongly semiregular and fuzzy hyperconnected space, then X is a fuzzy ∂ -space.*

Proof. Let λ be a fuzzy G_δ -set in X . Since X is a fuzzy strongly semiregular space, by Proposition 4.6, X is a fuzzy almost P-space. Then $\text{int}(\lambda) \neq 0$. Since X is a fuzzy hyperconnected space and $\text{int}(\lambda) \in T$, $\text{int}(\lambda)$ is a fuzzy dense set in X . Thus $\text{clint}(\lambda) = 1$. So by Theorem 2.15, X is a fuzzy ∂ -space. \square

The following proposition shows that fuzzy G_δ -sets are having non-zero interiors in fuzzy strongly semiregular spaces.

Proposition 4.14. *If λ is a non-zero fuzzy G_δ -set in a fuzzy strongly semiregular space (X, T) , then $\text{int}(\lambda) \neq 0$.*

Proof. Assume that λ is a non-zero fuzzy G_δ -set with $\text{int}(\lambda) = 0$. Then $1 - \lambda$ is a fuzzy F_σ -set in X such that $\text{cl}(1 - \lambda) = 1$. Since X is a fuzzy strongly semi regular space, by Proposition 4.6, $1 - \lambda = 1$. Thus $\lambda = 0$. This is a contradiction. So $\text{int}(\lambda) \neq 0$. \square

Corollary 4.15. *If λ is a fuzzy G_δ -set in a fuzzy strongly semiregular space (X, T) , then λ is a fuzzy somewhere dense set in X .*

Corollary 4.16. *If μ is a fuzzy F_σ -set in a fuzzy strongly semiregular space (X, T) , then μ is a fuzzy cs dense set in X .*

Remark 4.17. In view of corollary 4.16, one will have the following result.

Fuzzy G_δ -sets in fuzzy strongly semiregular spaces are fuzzy somewhere dense sets.

Proposition 4.18. *If a fuzzy topological space (X, T) is a fuzzy strongly regular space, then X is a fuzzy strongly semiregular space.*

Proof. Let λ be a fuzzy semi-open set in X . Since X is a fuzzy strongly regular space, there exist a family $(\delta_j)_{j \in J}$ of fuzzy open sets in X such that $\lambda = \bigvee_{j \in J} \delta_j$ and $\text{cl}(\delta_j) \leq \lambda$ for each $j \in J$. Then by Lemma 2.1, $(\text{cl}(\delta_j))_{j \in J}$ is a family of fuzzy regular closed sets in X . Thus X is a fuzzy strongly semiregular space. \square

The following proposition shows that fuzzy G_δ -sets in fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semi regular spaces are not fuzzy first category sets.

Proposition 4.19. *If λ is a fuzzy G_δ -set in a fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semiregular space (X, T) , then λ is a fuzzy second category set in X .*

Proof. Suppose λ is a fuzzy G_δ -set in X . Since X is a fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semiregular space, by Propositions 4.9 and 4.10, X is a fuzzy Baire and fuzzy P-space. Then by Theorem 2.17, λ is a fuzzy second category set in X . \square

The following proposition gives a condition for fuzzy G_δ -sets in fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semi regular spaces to become fuzzy Baire dense sets.

Proposition 4.20. *Let (X, T) be a fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semiregular space. If λ is a fuzzy G_δ -set in X such that $\lambda \leq \mu$ for each $\mu \in T$, then λ is a fuzzy Baire dense set in X .*

Proof. Suppose λ is a fuzzy G_δ -set in X such that $\lambda \leq \mu$ for each $\mu \in T$. Since X is a fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semiregular space, by Propositions 4.9 and 4.10, X is a fuzzy Baire and fuzzy P-space. Then Proposition 4.19, λ is a fuzzy second category set in X . Then by the hypothesis, $\lambda \wedge \mu = \lambda$. Note that λ is a non-zero fuzzy open set in X . Thus $\lambda \wedge \mu$ is a fuzzy second category set in X . So λ is a fuzzy Baire dense set in X . \square

The following proposition give conditions under which fuzzy Baire sets become fuzzy Baire dense sets in fuzzy strongly semi regular spaces.

Proposition 4.21. *If λ is a fuzzy Baire set in a fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semi regular space (X, T) , then λ is a fuzzy Baire dense set in X .*

Proof. Suppose λ is a fuzzy Baire set in X . Since X is a fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semi regular space, by Propositions 4.9 and 4.10, X is a fuzzy Baire and fuzzy P-space. Then X is a fuzzy globally disconnected, fuzzy Baire and fuzzy P-space. Thus by Theorem 2.18, λ is a fuzzy Baire dense set in X . \square


Proposition 4.22. *If (X, T) is a fuzzy strongly semiregular space, then X is a fuzzy second category space.*

Proof. The proof follows from Proposition 4.6 and Theorem 2.11. \square

Proposition 4.23. *If μ is a fuzzy F_σ -set in a fuzzy strongly semiregular and fuzzy hyperconnected space (X, T) , then λ is a fuzzy nowhere dense set in X .*

Proof. Suppose μ is a fuzzy F_σ -set in X . Then $1 - \mu$ is a fuzzy G_δ -set in X . Since X is a fuzzy strongly semiregular space, by Proposition 4.6, X is a fuzzy almost P-space. Thus $\text{int}(1 - \mu) \neq 0$. Since X is a fuzzy hyperconnected space and $\text{int}(1 - \mu) \in T$, $\text{int}(1 - \mu)$ is a fuzzy dense set in X . So $\text{clint}(1 - \mu) = 1$. This implies that $1 - \text{intcl}(\mu) = 1$, i.e., $\text{intcl}(\mu) = 0$. Hence λ is a fuzzy nowhere dense set in X . \square

Remark 4.24. The inter-relations between fuzzy strongly regular space, fuzzy strongly semi regular spaces, fuzzy quasi-regular spaces and fuzzy almost P -spaces can be stated as follows:



5. CONCLUSION

In this paper, the concept of fuzzy strongly semiregular space is introduced by means of fuzzy semi-open sets and fuzzy regular closed sets. Several characterizations of fuzzy strongly semi regular spaces are established. It is obtained that fuzzy residual sets, fuzzy G_δ -sets are fuzzy somewhere dense sets and fuzzy first category sets are not fuzzy dense sets and fuzzy F_σ -sets with zero interior are not fuzzy dense sets in fuzzy strongly semiregular spaces. Further it is established that fuzzy locally closed sets are subsets of fuzzy Baire sets in fuzzy strongly semiregular spaces. It is obtained that fuzzy G_δ -sets are fuzzy semi-open sets in fuzzy strongly semiregular and fuzzy hyperconnected spaces. It is shown that fuzzy globally disconnected and fuzzy quasi-regular spaces are fuzzy strongly semiregular spaces. It is obtained that fuzzy G_δ -sets in fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semi regular spaces are fuzzy second category sets. A condition under which fuzzy G_δ -sets become fuzzy Baire dense sets in fuzzy strongly semiregular spaces is also obtained. It is found that fuzzy Baire sets in fuzzy globally disconnected, fuzzy hyperconnected and fuzzy strongly semi regular spaces are fuzzy Baire dense sets. The inter-relations between fuzzy strongly semiregular spaces, fuzzy quasi-regular spaces and fuzzy almost P -spaces are also established in this paper.

Acknowledgements. The authors are grateful to the referees for their valuable suggestions.

REFERENCES

- [1] L. A. Zadeh, Fuzzy Sets, Information and Control 8 (1965) 338–353.
- [2] C. L. Chang, Fuzzy topological spaces, J. Math. Anal. Appl. 24 (1968) 182–190.
- [3] M. H. Stone, Applications of Boolean rings to general topology, Trans. Amer. Math. Soc. 41 (1937) 375–481.
- [4] M. K. Singal and S. P. Arya, On almost regular spaces, Glasnik. Mat. Ser. III4 (24) (1969) 89–99.
- [5] Bhamini M. Nayar, Some classes of generalized completely regular spaces, Inter. J. Pure and Appl. Math. 44 () (2008) 609–626.
- [6] K. K. Azad, On fuzzy semi continuity, fuzzy almost continuity and fuzzy weakly continuity, J. Math. Anal. Appl. 82 (1981) 14–32.
- [7] Yong Chan Kim and Jung Mi Ko, Fuzzy semi-regular spaces and fuzzy δ -continuous functions, Inter. J. Fuzzy Logic and Intelligent. Sys. 1 (1) (2001) 69–74.
- [8] Naseem Ajmal and B. K. Tyagi, Regular fuzzy spaces and fuzzy almost regular spaces, Matematiamki Vesnik 40 (102) (1988) 97–108.
- [9] Takashi Noiri and Saeid Jafari, On strongly semi-regular spaces, Demonstrate Mathematica XXXIV (1) (2001) 187–190.
- [10] Güzide Şenel, Lee Jeong Gon and Kul Hur, Advanced soft relation and soft mapping, Inte J. Comp. Intell. Sys. 14 (1) (2021) 461–470. Doi: <https://dx.doi.org/10.2991/ijcis.d.201221.001>.

- [11] Güzide Şenel, A new approach to Hausdorff space theory via the soft sets, *Math. Probl. Engin.* (9) (2016) 1–6. Doi: 10.1155/2016/2196743.
- [12] Güzide Şenel and Naim Çağman, Soft topological subspaces, *Ann. Fuzzy Math. Inform.* 10 (4) (2015) 525–535.
- [13] Güzide Şenel and Naim Çağman, Soft closed sets on soft bitopological Space, *J. New Results in Science* 3 (5) (2014) 57–66.
- [14] Güzide Şenel, Jeong-Gon Lee and Kul Hur, Distance and similarity measures for octahedron sets and their application to MCGDM problems, *Mathematics* 2020 8 (10) 1690. <https://doi.org/10.3390/math8101690>.
- [15] Jeong-Gon Lee, Güzide Şenel, Jong-il Baek, Sang-Hyeon Han and Hul Kur, Neighborhood structures and continuities via cubic sets, *Axioms* 2022 11 (8) 406. <https://doi.org/10.3390/axioms11080406>.
- [16] G. Balasubramanian, Maximal fuzzy topologies, *Kybernetika* 31 (5) (1995) 459–464.
- [17] G. Thangaraj and R. Palani, A short note on fuzzy residual sets and fuzzy functions, *Inter. Jour. Adv. Math.* 2017 (2) (2017) 1–9.
- [18] G. Thangaraj and G. Balasubramanian, On somewhat fuzzy continuous functions, *J. Fuzzy Math.* 11 (2) (2003) 725–736.
- [19] G. Thangaraj and S. Anjalmose, On fuzzy Baire spaces. *J. Fuzzy Math.* 21 (3) (2013) 667–676.
- [20] G. Thangaraj and E. Poongothai, On fuzzy σ -Baire spaces, *Int. J. Fuzzy Math. Sys.* 3 (4) (2013) 275–283.
- [21] G. Thangaraj, Resolvability and irresolvability in fuzzy topological spaces, *News Bull. Cal. Math. Soc.* 31 (46) (2008) 11–14.
- [22] G. Thangaraj, and S. Senthil, On somewhere fuzzy continuous functions, *Annl. Fuzzy Math. Inform.* 15 (2) (2018) 181–198.
- [23] Z. Petricevic, On fuzzy semi-regularization, separation properties and mappings, *Indian J. Pure Appl. Math.* 22 (12) (1991) 971–982.
- [24] P. K. Gain, R. P. Chakraborty and M. Pal, Characterization of some fuzzy subsets of fuzzy ideal topological spaces and decomposition of fuzzy continuity, *Inter. J. Fuzzy Math. and Sys.* 2 (2) (2012) 149–161.
- [25] G. Thangaraj and R. Palani, On fuzzy Baire sets, *Jour. Manag. Sci. Human. JOMSAH* 4 (2) (2017) 151–158.
- [26] G. Thangaraj and K. Dinakaran, On fuzzy pseudo-continuous functions, *IOSR. J. Math.* 15 (5) (2017) 12–20.
- [27] G. Thangaraj and K. Dinakaran, On fuzzy simply continuous functions, *J. Fuzzy Math.* 25 (1) (2017) 99–124.
- [28] G. Thangaraj and S. Lokeshwari, On fuzzy Baire dense sets and fuzzy Baire resolvable spaces, *Adv. Fuzzy Sets and Sys.* 23 (2-3) (2018) 93–116.
- [29] G. Thangaraj and M. Ponnusamy, On fuzzy quasi-regular spaces, *Inter. Conf. on Math. Methods and Computation (Dec 2022)*, Jamal Mohamed College (Auto), Tiuchirallpalli, Tamilnadu, India.
- [30] G. Thangaraj and G. Balasubramanian, On fuzzy basically disconnected spaces, *J. Fuzzy Math.* 9 (1) (2001) 103–110.
- [31] G. Thangaraj and C. Anbazhagan, On fuzzy almost P-spaces, *Inter. J. Innov. Sci. Engin. and Tech.* 2 (4) (2015) 389–407.
- [32] G. Thangaraj and A. Vinothkumar, On fuzzy semi-P-spaces and related concepts, *Pure and Appl. Math. J.* 11 (1) (2022) 20–27.
- [33] G. Thangaraj and S. Muruganatham, On fuzzy globally disconnected spaces, *Journal Tri. Math. Soc.* 21 (2019) 37–46.
- [34] Miguel Caldas, Govindappa Navalagi and Ratnesh Saraf, On fuzzy weakly semi-open functions, *Proyecciones* 21 (1) (2002) 51–63.
- [35] G. Thangaraj and J. Premkumar, On fuzzy ∂ -Spaces, *Inter. J. Appl. Engin. Research* 13 (15) (2018) 12417–12421.
- [36] Guo Shuangbing and Dang Faning, Fuzzy almost semi continuous functions and fuzzy strongly regular topological spaces, *Universite Savoie Mont Blanc* <https://projects.istic.univ-smb.fr/papers> 37–45.

- [37] G. Thangaraj and R. Palani, Somewhat fuzzy continuity and fuzzy Baire spaces, *Annl. Fuzzy Math. Inform.* 12 (1) (2016) 75–82.
- [38] Seok Jong Lee and Sang Min Yun, Fuzzy δ -topology and compactness, *Commun. Korean Math. Soc.* 27 (2) (2012) 357–368.
- [39] G. Thangaraj and C. Anbazhagan, Some remarks on fuzzy P-spaces, *Gen. Math. Notes* 26 (1) (2015) 8–16.
- [40] G. Thangaraj and S. Lokeshwari, Some remarks on fuzzy Baire dense sets, *Zeichen Journal* 7 (7) (2021) 255–272.
- [41] T. Noiri, On semi-continuous mapping, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur* 8 (54) (1973) 132–136.

G. THANGARAJ (g.thangaraj@rediffmail.com)

Department of Mathematics, Thiruvalluvar University,
Vellore - 632 115, Tamil Nadu, India

M.PONNUSAMY (ponsmsc87@gmail.com)

Research scholar, Department of Mathematics, Thiruvalluvar University,
Vellore - 632 115, Tamil Nadu, India